


Amended claims (clean version)

1. A compound of Formula (VI),



Formula (VI)

- 5 or a pharmaceutically acceptable salt or ester thereof;
 wherein:

X^1 is CH or N;

- X^2 , X^3 , X^4 , X^5 , X^6 and X^7 independently are NR^4 , CH_2 , CHQ or $C(Q)_2$, provided
 that: (i) zero, one or two of X^2 , X^3 , X^4 , X^5 , X^6 and X^7 are NR^4 ; (ii) when X^1 is N,
 10 both of X^2 and X^7 are not NR^4 ; (iii) when X^1 is N, X^3 and X^6 are not NR^4 ; and
 (iv) when X^1 is CH and two of X^2 , X^3 , X^4 , X^5 , X^6 and X^7 are NR^4 , the two NR^4 are
 located at adjacent ring positions or are separated by two or more other ring
 positions;

- A and V independently are H, halo, azido, $-CN$, $-CF_3$, $-CONR^1R^2$, $-NR^1R^2$, $-SR^2$,
 15 $-OR^2$, $-R^3$, $-W$, $-L-W$, $-W^0$, or $-L-N(R)-W^0$;

each Q is independently halo, azido, $-CN$, $-CF_3$, $-CONR^1R^2$, $-NR^1R^2$, $-SR^2$, $-OR^2$,
 $-R^3$, $-W$, $-L-W$, $-W^0$, or $-L-N(R)-W^0$;

- in each $-NR^1R^2$, R^1 and R^2 together with N may form an optionally substituted
 azacyclic ring, optionally containing one additional heteroatom selected from N,
 20 O and S as a ring member;

R¹ is H or C1-C6 alkyl, optionally substituted with one or more halogens, or =O;

R is H, or C1-C10 alkyl, C1-C10 heteroalkyl, C2-C10 alkenyl, or C2-C10 heteroalkenyl, each of which may be optionally substituted with one or more
 5 halogens, =O, or an optionally substituted 3-7 membered carbocyclic or heterocyclic ring;

R³ is an optionally substituted C1-C10 alkyl, C2-C10 alkenyl, C5-C10 aryl, or C6-C12 arylalkyl, or a heteroform of one of these, each of which may be optionally substituted with one or more halogens, =O, or an optionally
 10 substituted 3-6 membered carbocyclic or heterocyclic ring;

each R⁴ is independently H, or C1-C6 alkyl; or R⁴ may be -W, -L-W or -L-N(R)-W⁰;

each R is independently H or C1-C6 alkyl;

R⁷ is H and R⁸ is C1-C10 alkyl, C1-C10 heteroalkyl, C2-C10 alkenyl, or C2-C10
 15 heteroalkenyl, each of which may be optionally substituted with one or more halogens, =O, or an optionally substituted 3-7 membered carbocyclic or heterocyclic ring; or in —NR⁷R⁸, R⁷ and R⁸ together with N may form an optionally substituted azacyclic ring, optionally containing an additional heteroatom selected from N, O and S as a ring member;

20 m is 0, 1, 2, 3 or 4;

n is 0, 1, 2, 3, 4, or 5;


L is a C1-C10 alkylene, C1-C10 heteroalkylene, C2-C10 alkenylene or C2-C10 heteroalkenylene linker, each of which may be optionally substituted with one or more substituents selected from the group consisting of halogen, oxo (=O), or
 25 C1-C6 alkyl;

W is an optionally substituted 4-7 membered azacyclic ring, optionally containing an additional heteroatom selected from N, O and S as a ring member; and

W⁰ is an optionally substituted 3-4 membered carbocyclic ring, or a C1-C6 alkyl
 30 group substituted with from 1 to 4 fluorine atoms,

for use in treating multiple sclerosis in a subject.

2. A compound of Formula (VIII),



Formula (VIII)

or a pharmaceutically acceptable salt or ester thereof;

wherein:

A and V independently are H, halo, azido, -CN, -CF₃, -CONR¹R², -NR¹R², -SR²,

-OR², -R³, -W, -L-W, -W⁰, or -L-N(R)-W⁰;

each Q is independently halo, azido, -CN, -CF₃, -CONR¹R², -NR¹R², -SR², -OR², -R³, -W, -L-W, -W⁰, or -L-N(R)-W⁰;

in each —NR¹R², R¹ and R² together with N may form an optionally substituted azacyclic ring, optionally containing an additional heteroatom selected from N,

O and S as a ring member;

R¹ is H or C1-C6 alkyl, optionally substituted with one or more halogens, or =O;

R² is H, or C1-C10 alkyl, C1-C10 heteroalkyl, C2-C10 alkenyl, or C2-C10 heteroalkenyl, each of which may be optionally substituted with one or more halogens, =O, or an optionally substituted 3-7 membered carbocyclic or heterocyclic ring;

R³ is an optionally substituted C1-C10 alkyl, C2-C10 alkenyl, C5-C10 aryl, or C6-C12 arylalkyl, or a heteroform of one of these, each of which may be

optionally substituted with one or more halogens, =O, or an optionally substituted 3-6 membered carbocyclic or heterocyclic ring;
 each R⁴ is independently H, or C1-C6 alkyl; or R⁴ may be -W, -L-W or -L-N(R)-W⁰;

5 each R is independently H or C1-C6 alkyl;

R⁷ is H and R⁸ is C1-C10 alkyl, C1-C10 heteroalkyl, C2-C10 alkenyl, or C2-C10 heteroalkenyl, each of which may be optionally substituted with one or more halogens, =O, or an optionally substituted 3-7 membered carbocyclic or heterocyclic ring; or in —NR⁷R⁸, R⁷ and R⁸ together with N may form an

10 optionally substituted azacyclic ring, optionally containing an additional heteroatom selected from N, O and S as a ring member;

m is 0, 1, 2, 3 or 4;

n is 0, 1, 2, 3, 4 or 5;

p is 0, 1, 2 or 3;

15 L is a C1-C10 alkylene, C1-C10 heteroalkylene, C2-C10 alkenylene or C2-C10 heteroalkenylene linker, each of which may be optionally substituted with one or more substituents selected from the group consisting of halogen, oxo (=O), or C1-C6 alkyl;

W is an optionally substituted 4-7 membered azacyclic ring, optionally

20 containing an additional heteroatom selected from N, O and S as a ring member; and


W⁰ is an optionally substituted 3-4 membered carbocyclic ring, or a C1-C6 alkyl group substituted with from 1 to 4 fluorine atoms,
 for use in treating multiple sclerosis in a subject.

25

3. A compound for use according to claim 1 or 2, wherein R⁷ is H and R⁸ is a C₁₋₄ alkyl substituted with an optionally substituted aromatic heterocyclic ring.

4. A compound for use according to claim 3, wherein the optionally substituted aromatic heterocyclic ring is selected from pyridine, pyrimidine, pyrazine, imidazole, pyrrolidine, and thiazole.

5 5. A compound of Formula (VII)



Formula (VII)

or a pharmaceutically acceptable salt or ester thereof;

10 A and V independently are H, halo, azido, -CN, -CF₃, -CONR¹R², -NR¹R², -SR², -OR², -R³, -W, -L-W, -W⁰, or -L-N(R)-W⁰;

each Q is independently halo, azido, -CN, -CF₃, -CONR¹R², -NR¹R², -SR², -OR², -R³, -W, -L-W, -W⁰, or -L-N(R)-W⁰;

15 in each —NR¹R², R¹ and R² together with N may form an optionally substituted azacyclic ring, optionally containing an additional heteroatom selected from N, O and S as a ring member;


R¹ is H or C1-C6 alkyl, optionally substituted with one or more halogens, or =O;







20 R² is H, or C1-C10 alkyl, C1-C10 heteroalkyl, C2-C10 alkenyl, or C2-C10 heteroalkenyl, each of which may be optionally substituted with one or more halogens, =O, or an optionally substituted 3-7 membered carbocyclic or heterocyclic ring;







- R³ is an optionally substituted C1-C10 alkyl, C2-C10 alkenyl, C5-C10 aryl, or C6-C12 arylalkyl, or a heteroform of one of these, each of which may be optionally substituted with one or more halogens, =O, or an optionally substituted 3-6 membered carbocyclic or heterocyclic ring;
- 5 each R⁴ is independently H, or C1-C6 alkyl; or R⁴ may be -W, -L-W or -L-N(R)-W⁰;
- each R is independently H or C1-C6 alkyl;
- m is 0, 1, 2, 3 or 4;
- n is 0, 1, 2, 3, 4 or 5;
- 10 p is 0, 1, 2 or 3;
- L is a C1-C10 alkylene, C1-C10 heteroalkylene, C2-C10 alkenylene or C2-C10 heteroalkenylene linker, each of which may be optionally substituted with one or more substituents selected from the group consisting of halogen, oxo (=O), or C1-C6 alkyl;
- 15 W is an optionally substituted 4-7 membered azacyclic ring, optionally containing an additional heteroatom selected from N, O and S as a ring member; and
- W⁰ is an optionally substituted 3-4 membered carbocyclic ring, or a C1-C6 alkyl group substituted with from 1 to 4 fluorine atoms,
- 20 for use in treating multiple sclerosis in a subject.
6. A compound for use according to claim 5, wherein p is 0 or 1.
7. A compound for use according to any of claims 1 to 6, wherein A and V
- 25 are independently H or halo.
8. A compound for use according to any of claims 1 to 7, wherein R⁴ is H or C1-4 alkyl.







9. A compound for use according to any of claims 1 to 8, wherein m and n are each 0.


10. A compound selected from the group consisting of:











or a pharmaceutically acceptable salt or ester thereof,

for use in treating multiple sclerosis in a subject.

11. A method of monitoring treatment efficiency of a subject having multiple sclerosis, the method comprising:

comparing a level of expression of least one gene involved in the RNA polymerase I pathway in a cell of the subject who has been treated with said compound of any one of claims 1-10 to a level of expression of said at least one gene in a cell of the subject prior to being treated with said compound,


(i) wherein a decrease above a predetermined threshold in said level of expression of said at least one gene following said treating with said compound relative to said level of expression of said at least one gene prior to said treating with said compound indicates that said compound is efficient for treating the subject;

(ii) wherein an increase above a predetermined threshold in said level of expression of said at least one gene following said treating with said compound relative to said level of expression of said at least one gene prior to said treating with said compound indicates that said compound is not efficient for treating the subject; or

(iii) wherein when a level of expression of said at least one gene following said treating with said compound is identical or changed below a predetermined threshold as compared to prior to said treating with said compound then the treatment is not efficient for treating the subject

thereby monitoring treatment efficiency of the subject having multiple sclerosis.

12. A compound for use according to any one of claims 1-10, or the method of claim 11, wherein said compound is 2-(4-methyl-1,4-diazepan-1-yl)-N-((5-methylpyrazin-2-yl)methyl)-5-oxo-5H-benzo[4,5]thiazolo[3,2-a][1,8]naphthyridine-6-carboxamide as depicted in Formula



13. A compound for use according to any of claims 1-10 or the method of claim 11 or 12, wherein said multiple sclerosis is a relapsing-remitting course of multiple sclerosis.

14. A compound for use according to any of claims 1-10 or the method of claim 11 or 12, wherein said multiple sclerosis is a progressive course of multiple sclerosis.

15. A compound for use according to any of claims 1-10 or the method of claim 11 or 12, wherein said multiple sclerosis is a benign multiple sclerosis.