


New Claims

1. A compound of Formula (I),



5 Formula (I)

- or a pharmaceutically acceptable salt or ester thereof;
wherein ----- indicates an optionally unsaturated bond;
each of B, X, A or V is absent if Z¹, Z², Z³ and Z⁴, respectively, is N; and
10 each of B, X, A and V is independently H, halo, azido, —CN, —CF₃, —
CONR¹R², —NR¹R², —SR², —OR², —R³, —W, —L-W, —W⁰, —L-N(R) —W⁰, A² or A³, when
each of Z¹, Z², Z³ and Z⁴, respectively, is C;
Z is O, S, CR⁴₂, NR⁴CR⁴, CR⁴NR⁴, CR⁴, NR⁴ or N;
each of Z¹, Z², Z³ and Z⁴ is independently C or N, provided any three N are non-
15 adjacent;
Z⁵ is C; or Z⁵ may be N when Z is N;
Y is an optionally substituted 5-6 membered carbocyclic or heterocyclic ring;
U¹ is —C(=T)N(R)—, —C(=T)N(R)O—, —C(=T)—, —SO₂N(R)—, —SO₂N(R)N(R⁰)—, —SO₂—, or
—SO₃—, where T is O, S, or NH; or U¹ may be a bond when Z⁵ is N or U² is H;
20 U² is H, or C3-C7 cycloalkyl, C1-C10 alkyl, C1-C10 heteroalkyl, C2-C10 alkenyl
or C2-C10 heteroalkenyl group, each of which may be optionally substituted

with one or more halogens, =O, or an optionally substituted 3-7 membered carbocyclic or heterocyclic ring; or U^2 is -W, -L-W, -L-N(R)- W^0 , A^2 or A^3 ; in each -NR¹R², R¹ and R² together with N may form an optionally substituted azacyclic ring, optionally containing an additional heteroatom selected from N, O and S as a ring member;

R¹ is H or C1-C6 alkyl, optionally substituted with one or more halogens, or =O;

R² is H, or C1-C10 alkyl, C1-C10 heteroalkyl, C2-C10 alkenyl, or C2-C10 heteroalkenyl, each of which may be optionally substituted with one or more halogens, =O, or an optionally substituted 3-7 membered carbocyclic or heterocyclic ring;

R³ is an optionally substituted C1-C10 alkyl, C2-C10 alkenyl, C5-C10 aryl, or C6-C12 arylalkyl, or a heteroform of one of these, each of which may be optionally substituted with one or more halogens, =O, or an optionally substituted 3-6 membered carbocyclic or heterocyclic ring;

each R⁴ is independently H, or C1-C6 alkyl; or R⁴ may be -W, -L-W or -L-N(R)- W^0 ;

each R and R⁰ is independently H or C1-C6 alkyl;

L is a C1-C10 alkylene, C1-C10 heteroalkylene, C2-C10 alkenylene or C2-C10 heteroalkenylene linker, each of which may be optionally substituted with one or more substituents selected from the group consisting of halogen, oxo (=O), or C1-C6 alkyl;

W is an optionally substituted 4-7 membered azacyclic ring, optionally containing an additional heteroatom selected from N, O and S as a ring member;


W^0 is an optionally substituted 3-4 membered carbocyclic ring, or a C1-C6 alkyl group substituted with from 1 to 4 fluorine atoms;

provided one of U^2 , V, A, X and B is a secondary amine A^2 or a tertiary amine A^3 , wherein

the secondary amine A^2 is -NH- W^0 , and

the tertiary amine A^3 is a fully saturated and optionally substituted six-membered or seven-membered azacyclic ring optionally containing an additional heteroatom selected from N, O or S as a ring member, or the tertiary amine A^3 is a partially unsaturated or aromatic optionally substituted five-membered azacyclic ring, optionally containing an additional heteroatom selected from N, O or S as a ring member;
 for use in treating an autoimmune disease in a subject.

2. A compound of Formula (II),



Formula (II)

or a pharmaceutically acceptable salt or ester thereof;
 wherein ----- indicates an optionally unsaturated bond;
 each of A and X is independently H, halo, azido, -CN, -CF₃, -CONR¹R², -NR¹R²,
 -SR², -OR², -R³, -W, -L-W, -W⁰, -L-N(R)-W⁰, A² or A³;
 Z is O, S, CR⁴₂, NR⁴CR⁴, CR⁴NR⁴ or NR⁴;
 Y is an optionally substituted 5-6 membered carbocyclic or heterocyclic ring;
 U¹ is -C(=T)N(R)-, -C(=T)N(R)O-, -C(=T)-, -SO₂N(R)-, -SO₂N(R)N(R⁰)-, -SO₂-, or
 -SO₃-, where T is O, S, or NH; or U¹ may be a bond when U² is H;
 U² is H, or C3-C7 cycloalkyl, C1-C10 alkyl, C1-C10 heteroalkyl, C2-C10 alkenyl
 or C2-C10 heteroalkenyl group, each of which may be optionally substituted
 with one or more halogens, =O, or an optionally substituted 3-7 membered
 carbocyclic or heterocyclic ring; or U² is -W, -L-W, -L-N(R)-W⁰, A² or A³;

in each $-NR^1R^2$, R^1 and R^2 together with N may form an optionally substituted azacyclic ring, optionally containing an additional heteroatom selected from N, O and S as a ring member;

R^1 is H or C1-C6 alkyl, optionally substituted with one or more halogens, or
5 $=O$;

R^2 is H, or C1-C10 alkyl, C1-C10 heteroalkyl, C2-C10 alkenyl, or C2-C10 heteroalkenyl, each of which may be optionally substituted with one or more halogens, $=O$, or an optionally substituted 3-7 membered carbocyclic or heterocyclic ring;

10 R^3 is an optionally substituted C1-C10 alkyl, C2-C10 alkenyl, C5-C10 aryl, or C6-C12 arylalkyl, or a heteroform of one of these, each of which may be optionally substituted with one or more halogens, $=O$, or an optionally substituted 3-6 membered carbocyclic or heterocyclic ring;

each R^4 is independently H, or C1-C6 alkyl; or R^4 may be $-W$, $-L-W$ or $-L-N(R)-$
15 W^0 ;

each R and R^0 is independently H or C1-C6 alkyl;

L is a C1-C10 alkylene, C1-C10 heteroalkylene, C2-C10 alkenylene or C2-C10 heteroalkenylene linker, each of which may be optionally substituted with one or more substituents selected from the group consisting of halogen, oxo ($=O$), or
20 C1-C6 alkyl;

W is an optionally substituted 4-7 membered azacyclic ring, optionally containing an additional heteroatom selected from N, O and S as a ring member;

W^0 is an optionally substituted 3-4 membered carbocyclic ring, or a C1-C6 alkyl
25 group substituted with from 1 to 4 fluorine atoms;

provided that one of U^2 , A, and X is a secondary amine A^2 or a tertiary amine A^3 , wherein


the secondary amine A^2 is $-NH-W^0$, and

the tertiary amine A^3 is a fully saturated and optionally substituted six-
30 membered or seven-membered azacyclic ring optionally containing an

additional heteroatom selected from N, O or S as a ring member, or the tertiary amine A³ is a partially unsaturated or aromatic optionally substituted five-membered azacyclic ring optionally containing an additional heteroatom selected from N, O or S as a ring member;

5 for use in treating an autoimmune disease in a subject.

3. A compound of Formula (III),



Formula (III)

10

or a pharmaceutically acceptable salt or ester thereof;

wherein ---- indicates an optionally unsaturated bond; and

each of B, X, A or V is absent if Z¹, Z², Z³ and Z⁴, respectively, is N; and

each of B, X, A and V is independently H, halo, azido, -CN, -CF₃, -CONR¹R², -

15 NR¹R², -SR², -OR², -R³, -W, -L-W, -W⁰, -L-N(R)-W⁰, A² or A³, when each of Z¹, Z², Z³ and Z⁴, respectively, is C;

each of Z¹, Z², Z³ and Z⁴ is independently C or N, provided any three N are non-adjacent;

Y is an optionally substituted 5-6 membered carbocyclic or heterocyclic ring;

20 U¹ is -C(=T)N(R)-, -C(=T)N(R)O-, -C(=T)-, -SO₂N(R)-, -SO₂N(R)N(R⁰)-, -SO₂-, or -SO₃-, where T is O, S, or NH; or U¹ may be a bond when Z⁵ is N or U² is H;

U² is H, or C3-C7 cycloalkyl, C1-C10 alkyl, C1-C10 heteroalkyl, C2-C10 alkenyl or C2-C10 heteroalkenyl group, each of which may be optionally substituted

with one or more halogens, =O, or an optionally substituted 3-7 membered carbocyclic or heterocyclic ring; or U² is -W, -L-W, -L-N(R) -W⁰, A² or A³; in each -NR¹R², R¹ and R² together with N may form an optionally substituted azacyclic ring, optionally containing an additional heteroatom selected from N, O and S as a ring member;

R¹ is H or C1-C6 alkyl, optionally substituted with one or more halogens, or =O;

R² is H, or C1-C10 alkyl, C1-C10 heteroalkyl, C2-C10 alkenyl, or C2-C10 heteroalkenyl, each of which may be optionally substituted with one or more halogens, =O, or an optionally substituted 3-7 membered carbocyclic or heterocyclic ring;

R³ is an optionally substituted C1-C10 alkyl, C2-C10 alkenyl, C5-C10 aryl, or C6-C12 arylalkyl, or a heteroform of one of these, each of which may be optionally substituted with one or more halogens, =O, or an optionally substituted 3-6 membered carbocyclic or heterocyclic ring;

each R⁴ is independently H, or C1-C6 alkyl; or R⁴ may be -W, -L-W or -L-N(R)-W⁰;

each R and R⁰ is independently H or C1-C6 alkyl;

L is a C1-C10 alkylene, C1-C10 heteroalkylene, C2-C10 alkenylene or C2-C10 heteroalkenylene linker, each of which may be optionally substituted with one or more substituents selected from the group consisting of halogen, oxo (=O), or C1-C6 alkyl;

W is an optionally substituted 4-7 membered azacyclic ring, optionally containing an additional heteroatom selected from N, O and S as a ring member;


W⁰ is an optionally substituted 3-4 membered carbocyclic ring, or a C1-C6 alkyl group substituted with from 1 to 4 fluorine atoms;

provided that one of U², V, A, X and B is a secondary amine A² or a tertiary amine A³, wherein

the secondary amine A² is -NH-W⁰, and

the tertiary amine A³ is a fully saturated and optionally substituted six-membered or seven-membered azacyclic ring optionally containing an additional heteroatom selected from N, O or S as a ring member, or the tertiary amine A³ is a partially unsaturated or aromatic optionally substituted five-membered azacyclic ring, optionally containing an additional heteroatom selected from N, O or S as a ring member,
 5 for use in treating an autoimmune disease in a subject.

4. A compound of Formula (IV),



Formula (IV)

or a pharmaceutically acceptable salt or ester thereof;
 wherein ----- indicates an optionally unsaturated bond;
 15 each of B, X, A or V is absent if Z¹, Z², Z³ and Z⁴ respectively, is N; and
 each of B, X, A and V is independently H, halo, azido, -CN, -CF₃, -CONR¹R², —
 NR¹R², -SR², -OR², -R³, -W, -L-W, -W⁰, -L-N(R)-W⁰, A² or A³, when each of Z¹,
 Z², Z³ and Z⁴, respectively, is C;
 each of Z¹, Z², Z³ and Z⁴ is independently C or N, provided any three N are non-
 20 adjacent;
 Y is an optionally substituted 5-6 membered carbocyclic or heterocyclic ring;
 in each NR¹R², R¹ and R² together with N may form an optionally substituted
 azacyclic ring, optionally containing an additional heteroatom selected from N,
 O and S as a ring member;

R¹ is H or C1-C6 alkyl, optionally substituted with one or more halogens, or =O;

R² is H, or C1-C10 alkyl, C1-C10 heteroalkyl, C2-C10 alkenyl, or C2-C10 heteroalkenyl, each of which may be optionally substituted with one or more
 5 halogens, =O, or an optionally substituted 3-7 membered carbocyclic or heterocyclic ring;

R³ is an optionally substituted C1-C10 alkyl, C2-C10 alkenyl, C5-C10 aryl, or C6-C12 arylalkyl, or a heteroform of one of these, each of which may be optionally substituted with one or more halogens, =O, or an optionally
 10 substituted 3-6 membered carbocyclic or heterocyclic ring;

R⁴ is -W, -L-W or -L-N(R)-W⁰; and


each R is independently H or C1-C6 alkyl;

L is a C1-C10 alkylene, C1-C10 heteroalkylene, C2-C10 alkenylene or C2-C10 heteroalkenylene linker, each of which may be optionally substituted with one
 15 or more substituents selected from the group consisting of halogen, oxo (=O), or C1-C6 alkyl;

W is an optionally substituted 4-7 membered azacyclic ring, optionally containing an additional heteroatom selected from N, O and S as a ring member; and

20 W⁰ is an optionally substituted 3-4 membered carbocyclic ring, or a C1-C6 alkyl group substituted with from 1 to 4 fluorine atoms,
 for use in treating an autoimmune disease in a subject.

5. A compound of Formula (V),



Formula (V)

or a pharmaceutically acceptable salt or ester thereof;

5 ---- indicates an optionally unsaturated bond;

A and V independently are H, halo, azido, -CN, -CF₃, -CONR¹R², -NR¹R², -SR², -OR², -R³, -W, -L-W, -W⁰, -L-N(R)-W⁰, A² or A³;

Z is O, S, CR₄², NR⁴CR⁴, CR⁴NR⁴ or NR⁴;

Y is an optionally substituted 5-6 membered carbocyclic or heterocyclic ring;

10 U¹ is -C(=T)N(R)-, -C(=T)N(R)O-, -C(=T)-, -SO₂N(R)-, -SO₂N(R)N(R⁰)-, -SO₂-, or -SO₃-, where T is O, S, or NH; or U¹ may be a bond when U² is H;

U² is H, or C3-C7 cycloalkyl, C1-C10 alkyl, C1-C10 heteroalkyl, C2-C10 alkenyl or C2-C10 heteroalkenyl group, each of which may be optionally substituted with one or more halogens, =O, or an optionally substituted 3-7 membered carbocyclic or heterocyclic ring; or U² is -W, -L-W or -L-N(R)-W⁰, A² or A³;

15 in each -NR¹R², R¹ and R² together with N may form an optionally substituted azacyclic ring, optionally containing an additional heteroatom selected from N, O and S as a ring member;


R¹ is H or C1-C6 alkyl, optionally substituted with one or more halogens, or

20 =O;

R² is H, or C1-C10 alkyl, C1-C10 heteroalkyl, C2-C10 alkenyl, or C2-C10 heteroalkenyl, each of which may be optionally substituted with one or more halogens, =O, or an optionally substituted 3-7 membered carbocyclic or heterocyclic ring;

- R^3 is an optionally substituted C1-C10 alkyl, C2-C10 alkenyl, C5-C10 aryl, or C6-C12 arylalkyl, or a heteroform of one of these, each of which may be optionally substituted with one or more halogens, =O, or an optionally substituted 3-6 membered carbocyclic or heterocyclic ring;
- 5 each R^4 is independently H, or C1-C6 alkyl; or R^4 may be -W, -L-W or -L-N(R)-W⁰;
- each R and R^0 is independently H or C1-C6 alkyl;
- L is a C1-C10 alkylene, C1-C10 heteroalkylene, C2-C10 alkenylene or C2-C10 heteroalkenylene linker, each of which may be optionally substituted with one
- 10 or more substituents selected from the group consisting of halogen, oxo (=O), or C1-C6 alkyl;
- W is an optionally substituted 4-7 membered azacyclic ring, optionally containing an additional heteroatom selected from N, O and S as a ring member;
- 15 W⁰ is an optionally substituted 3-4 membered carbocyclic ring, or a C1-C6 alkyl group substituted with from 1 to 4 fluorine atoms;
- provided one of U², A, and V is a secondary amine A² or a tertiary amine A³, wherein
- the secondary amine A² is -NH-W⁰, and
- 20 the tertiary amine A³ is a fully saturated and optionally substituted six-membered or seven-membered azacyclic ring optionally containing an additional heteroatom selected from N, O or S as a ring member, or the tertiary amine A³ is a partially unsaturated or aromatic optionally substituted five-membered azacyclic ring optionally containing an additional heteroatom
- 25 selected from N, O or S as a ring member,
- for use in treating an autoimmune disease in a subject.

6. A compound of Formula (VI),



Formula (VI)

or a pharmaceutically acceptable salt or ester thereof;

5 wherein:

X^1 is CH or N;

X^2 , X^3 , X^4 , X^5 , X^6 and X^7 independently are NR^4 , CH_2 , CHQ or $C(Q)_2$, provided that: (i) zero, one or two of X^2 , X^3 , X^4 , X^5 , X^6 and X^7 are NR^4 ; (ii) when X^1 is N, both of X^2 and X^7 are not NR^4 ; (iii) when X^1 is N, X^3 and X^6 are not NR^4 ; and

10 (iv) when X^1 is CH and two of X^2 , X^3 , X^4 , X^5 , X^6 and X^7 are NR^4 , the two NR^4 are located at adjacent ring positions or are separated by two or more other ring positions;

A and V independently are H, halo, azido, -CN, -CF₃, -CONR¹R², -NR¹R², -SR², -OR², -R³, -W, -L-W, -W⁰, or -L-N(R)-W⁰;

15 each Q is independently halo, azido, -CN, -CF₃, -CONR¹R², -NR¹R², -SR², -OR², -R³, -W, -L-W, -W⁰, or -L-N(R)-W⁰;

in each —NR¹R², R¹ and R² together with N may form an optionally substituted azacyclic ring, optionally containing one additional heteroatom selected from N, O and S as a ring member;

20 R¹ is H or C1-C6 alkyl, optionally substituted with one or more halogens, or =O;

R is H, or C1-C10 alkyl, C1-C10 heteroalkyl, C2-C10 alkenyl, or C2-C10 heteroalkenyl, each of which may be optionally substituted with one or more

halogens, =O, or an optionally substituted 3-7 membered carbocyclic or heterocyclic ring;

R³ is an optionally substituted C1-C10 alkyl, C2-C10 alkenyl, C5-C10 aryl, or C6-C12 arylalkyl, or a heteroform of one of these, each of which may be

5 optionally substituted with one or more halogens, =O, or an optionally substituted 3-6 membered carbocyclic or heterocyclic ring;

each R⁴ is independently H, or C1-C6 alkyl; or R⁴ may be -W, -L-W or -L-N(R)-W⁰;

each R is independently H or C1-C6 alkyl;

10 R⁷ is H and R⁸ is C1-C10 alkyl, C1-C10 heteroalkyl, C2-C10 alkenyl, or C2-C10 heteroalkenyl, each of which may be optionally substituted with one or more halogens, =O, or an optionally substituted 3-7 membered carbocyclic or heterocyclic ring; or in —NR⁷R⁸, R⁷ and R⁸ together with N may form an optionally substituted azacyclic ring, optionally containing an additional

15 heteroatom selected from N, O and S as a ring member;

m is 0, 1, 2, 3 or 4;

n is 0, 1, 2, 3, 4, or 5;


L is a C1-C10 alkylene, C1-C10 heteroalkylene, C2-C10 alkenylene or C2-C10 heteroalkenylene linker, each of which may be optionally substituted with one

20 or more substituents selected from the group consisting of halogen, oxo (=O), or C1-C6 alkyl;

W is an optionally substituted 4-7 membered azacyclic ring, optionally containing an additional heteroatom selected from N, O and S as a ring member; and

25 W⁰ is an optionally substituted 3-4 membered carbocyclic ring, or a C1-C6 alkyl group substituted with from 1 to 4 fluorine atoms,
for use in treating an autoimmune disease in a subject.

7. A compound of Formula (VIII),



Formula (VIII)

or a pharmaceutically acceptable salt or ester thereof;

5 wherein:

A and V independently are H, halo, azido, -CN, -CF₃, -CONR¹R², -NR¹R², -SR², -OR², -R³, -W, -L-W, -W⁰, or -L-N(R)-W⁰;

each Q is independently halo, azido, -CN, -CF₃, -CONR¹R², -NR¹R², -SR², -OR², -R³, -W, -L-W, -W⁰, or -L-N(R)-W⁰;

10 in each —NR¹R², R¹ and R² together with N may form an optionally substituted azacyclic ring, optionally containing an additional heteroatom selected from N, O and S as a ring member;

R¹ is H or C1-C6 alkyl, optionally substituted with one or more halogens, or =O;

15 R² is H, or C1-C10 alkyl, C1-C10 heteroalkyl, C2-C10 alkenyl, or C2-C10 heteroalkenyl, each of which may be optionally substituted with one or more halogens, =O, or an optionally substituted 3-7 membered carbocyclic or heterocyclic ring;

R³ is an optionally substituted C1-C10 alkyl, C2-C10 alkenyl, C5-C10 aryl, or
20 C6-C12 arylalkyl, or a heteroform of one of these, each of which may be optionally substituted with one or more halogens, =O, or an optionally substituted 3-6 membered carbocyclic or heterocyclic ring;

each R^4 is independently H, or C1-C6 alkyl; or R^4 may be -W, -L-W or -L-N(R)- W^0 ;

each R is independently H or C1-C6 alkyl;

R^7 is H and R^8 is C1-C10 alkyl, C1-C10 heteroalkyl, C2-C10 alkenyl, or C2-C10

5 heteroalkenyl, each of which may be optionally substituted with one or more halogens, =O, or an optionally substituted 3-7 membered carbocyclic or heterocyclic ring; or in $—NR^7R^8$, R^7 and R^8 together with N may form an optionally substituted azacyclic ring, optionally containing an additional heteroatom selected from N, O and S as a ring member;

10 m is 0, 1, 2, 3 or 4;

n is 0, 1, 2, 3, 4 or 5;

p is 0, 1, 2 or 3;

L is a C1-C10 alkylene, C1-C10 heteroalkylene, C2-C10 alkenylene or C2-C10 heteroalkenylene linker, each of which may be optionally substituted with one


15 or more substituents selected from the group consisting of halogen, oxo (=O), or C1-C6 alkyl;

W is an optionally substituted 4-7 membered azacyclic ring, optionally containing an additional heteroatom selected from N, O and S as a ring member; and

20 W^0 is an optionally substituted 3-4 membered carbocyclic ring, or a C1-C6 alkyl group substituted with from 1 to 4 fluorine atoms,

for use in treating an autoimmune disease in a subject.

8. A compound of Formula (VII)



Formula (VII)

or a pharmaceutically acceptable salt or ester thereof;

- 5 A and V independently are H, halo, azido, -CN, -CF₃, -CONR¹R², -NR¹R², -SR², -OR², -R³, -W, -L-W, -W⁰, or -L-N(R)-W⁰;

each Q is independently halo, azido, -CN, -CF₃, -CONR¹R², -NR¹R², -SR², -OR², -R³, -W, -L-W, -W⁰, or -L-N(R)-W⁰;

- in each —NR¹R², R¹ and R² together with N may form an optionally substituted
10 azacyclic ring, optionally containing an additional heteroatom selected from N, O and S as a ring member;

R¹ is H or C1-C6 alkyl, optionally substituted with one or more halogens, or =O;

- R² is H, or C1-C10 alkyl, C1-C10 heteroalkyl, C2-C10 alkenyl, or C2-C10
15 heteroalkenyl, each of which may be optionally substituted with one or more halogens, =O, or an optionally substituted 3-7 membered carbocyclic or heterocyclic ring;

- R³ is an optionally substituted C1-C10 alkyl, C2-C10 alkenyl, C5-C10 aryl, or C6-C12 arylalkyl, or a heteroform of one of these, each of which may be
20 optionally substituted with one or more halogens, =O, or an optionally substituted 3-6 membered carbocyclic or heterocyclic ring;

each R⁴ is independently H, or C1-C6 alkyl; or R⁴ may be -W, -L-W or -L-N(R)-W⁰;

each R is independently H or C1-C6 alkyl;

m is 0, 1, 2, 3 or 4;

n is 0, 1, 2, 3, 4 or 5;

p is 0, 1, 2 or 3;

- 5 L is a C1-C10 alkylene, C1-C10 heteroalkylene, C2-C10 alkenylene or C2-C10 heteroalkenylene linker, each of which may be optionally substituted with one or more substituents selected from the group consisting of halogen, oxo (=O), or C1-C6 alkyl;

10 W is an optionally substituted 4-7 membered azacyclic ring, optionally containing an additional heteroatom selected from N, O and S as a ring member; and

W⁰ is an optionally substituted 3-4 membered carbocyclic ring, or a C1-C6 alkyl group substituted with from 1 to 4 fluorine atoms,
for use in treating an autoimmune disease in a subject.

15

9. A compound selected from the group consisting of:

