

Quantum Mechanics

Diez B. Borja

May 7, 2022

Chapter 1

The wave function

1.1 The Schrodinger equation

Imagine a particle of mass m , constrained to move along the x -axis, subject to some specified force $F(x, t)$. The program of *classical* mechanics is to determine the position of the particle at any given time: $x(t)$. Once we know that, we can figure out the velocity ($v = dx/dt$), the momentum ($p = mv$), the kinetic energy ($T = (1/2)mv^2$), or any other dynamical variable of interest. And how do we go about determining $x(t)$? We apply Newton's second law: $F = ma$. (For *conservative* systems—the only kinds we shall consider, and, fortunately, the only kinds that occur at the microscopic level—the force can be expressed as the derivative of a potential energy function¹, $F = -\partial V/\partial x$, and Newton's law reads $m d^2x/dt^2 = -\partial V/\partial x$) This, together with appropriate initial conditions (typically the position and velocity at $t = 0$), determines $x(t)$.

Quantum mechanics approaches this same problem quite differently. In this case what we're looking for is the particle's **wave function**, $\Psi(x, t)$, and we get it by solving the **Schrodinger equation**:

$$\boxed{i\hbar \frac{\partial \Psi}{\partial t} = -\frac{\hbar^2}{2m} \frac{\partial^2 \Psi}{\partial x^2} + V\Psi} \quad (1.1) \quad \text{Schrodinger equation}$$

This equation plays a role logically analogous to Newton's second law: Given suitable initial conditions (typically, $\Psi(x, 0)$), the Schrodinger equation determines $\Psi(x, t)$ for all future time, just as, in classical mechanics, Newton's law determines $x(t)$ for all future time.

1.2 The statistical interpretation

What exactly is this "wave function"? Born's **statistical interpretation** of the wave function, says that $|\Psi(x, t)|^2$ gives the **probability** of finding the particle at point x , at time t , or more precisely

$$\boxed{\int_a^b |\Psi(x, t)|^2 dx = \text{probability of finding the particle between } a \text{ and } b, \text{ at time } t} \quad (1.2) \quad \text{Statistical interpretation}$$

¹Magnetic forces are an exception, but let's not worry about them just yet. By the way, we shall assume for the moment that the motion is nonrelativistic ($v \ll c$)

Probability is the *area* under the graph of $|\Psi(x, t)|^2$

The statistical interpretation introduces a kind of **indeterminacy** into quantum mechanics, for even if you know everything the theory has to tell you about the particle /to wit: its wave function, still you cannot predict with certainty the outcome of a simple experiment to measure its position– all quantum mechanics has to offer is *statistical* information about the *possible* results.

We say that the wave function **collapses**, upon measurement, to spike some point. There are, then, two entirely distinct kinds of physical processes: "ordinary" ones, in which the wave function evolves in a leisurely fashion under the Schrodinger equation, and "measurements", in which Ψ suddenly and discontinuously collapses ².

1.3 Probability

I will give some definitions. The total number is

$$N = \sum_{j=0}^{\infty} N(j) \quad (1.3)$$

The probability of getting j is

$$P(j) = \frac{N(j)}{N} \quad (1.4)$$

In particular, the sum of all the probabilities is 1:

$$\sum_{j=0}^{\infty} P(j) = 1 \quad (1.5)$$

In general, the average value of some *function* of j es given by

$$\boxed{\langle f(j) \rangle = \sum_{j=0}^{\infty} f(j)P(j)} \quad (1.6) \quad \text{Average value of a function}$$

In quantum mechanics the average is usually the quantity of interest; in that context it has come to be called the **expectation value**.

Now we need a numerical measure of the amount of "spread" in a distribution, with respect to the average. The most obvious way to do this would be to find out how far each individual deviates from the average,

$$\Delta j = j - \langle j \rangle \quad (1.7)$$

and compute the average of Δj . Trouble is, of course, that you get zero, by the nature of the average, Δj is as often negative as positive. To avoid this irritating problem you might decide

²The role of measurement in quantum mechanics is so critical and so bizarre that you may well be wondering what precisely constitutes a measurement. Does it have to do with the interaction between a microscopic (quantum) system and a macroscopic (classical) measuring apparatus (as Bohr insisted), or is it intervention of a conscious "observer" (as Wigner proposed)? For the moment let's take the naive view: A measurement is the kind of thing that a scientist does in the laboratory, with rules, stopwatches, and so on

to average the *absolute value* of Δj . But absolute values are nasty to work with; instead, we get around the sign problem by *squaring* before averaging:

$$\sigma^2 \equiv \langle (\Delta j)^2 \rangle \quad (1.8) \quad \text{Variance}$$

This quantity is known as the **variance** of the distribution; σ itself is called the **standard deviation**.

There is a useful little theorem on variances:

$$\sigma = \sqrt{\langle j^2 \rangle - \langle j \rangle^2} \quad (1.9)$$

1.4 Continuous variables