

SEPTEMBER 25, 2024

# Nonlinear Electrodynamics

---

**Borja Diez**

*Universidad Arturo Prat*

*E-mail:* [borjadiez1014@gmail.com](mailto:borjadiez1014@gmail.com)

ABSTRACT: Personal notes on nonlinear electrodynamics.

---

## Contents

### 1 Nonlinear Electrodynamics

---

1

## 1 Nonlinear Electrodynamics

As is defined in the Plebanski book [XX], nonlinear electrodynamics are described by the following action principle

$$S_E[g, A, P] = -\frac{1}{4\pi} \int d^4x \sqrt{-g} \left( \frac{1}{2} F_{\mu\nu} P^{\mu\nu} - \mathcal{H}(\mathcal{P}, \mathcal{Q}) \right) \quad (1.1)$$

which depends on the metric  $g_{\mu\nu}$ , the gauge potential  $A_\mu$ , and the antisymmetric tensor  $P_{\mu\nu}$ . Here the structural function  $\mathcal{H}$  describes the precise nonlinear electrodynamics and depends, in general, on the two Lorentz scalars that can be constructed with  $P^{\mu\nu}$  [XX]. As usual, the field strength is related to the gauge potential as  $F = dA$ , ensuring the Faraday equations

$$dF = 0. \quad (1.2)$$

On the other hand, the variation of action (1.1) with respect to the gauge potential leads to the Maxwell equations

$$d \star P = 0, \quad (1.3)$$

where  $\star$  stands for the Hodge dual, whereas varying (1.1) with respect to the antisymmetric tensor  $P^{\mu\nu}$  yields the constitutive relations

$$F_{\mu\nu} = \frac{\partial \mathcal{H}}{\partial \mathcal{P}} P_{\mu\nu} + \frac{\partial \mathcal{H}}{\partial \mathcal{Q}} \star P_{\mu\nu}. \quad (1.4)$$

Notice that Maxwell electrodynamics is recovered for  $\mathcal{H} = \mathcal{P}$ , giving linear constitutive relations. Lastly, the corresponding energy-momentum tensor reads

$$4\pi T_{\mu\nu}^E = F_{\mu\alpha} P_\nu{}^\alpha - g_{\mu\nu} \left( \frac{1}{2} F_{\alpha\beta} P^{\alpha\beta} - \mathcal{H} \right). \quad (1.5)$$

The main motivation for the action principle (1) is that now Maxwell equations (3) remain linear as the Faraday ones (2) while the nonlinearity is encoded into the constitutive relations (4). Consequently, Maxwell equations (3) can be now understood just like the Faraday ones (2), i.e., implying the local existence of a vector potential  $P = dA$ . Therefore, from the point of view of the action principle (1), a solution to nonlinear electrodynamics can be understood as a pair of vector potentials  $A$  and  $A$  compatible with the constitutive relations (4). Additionally, since in four dimensions both Faraday (2) and Maxwell (3) equations define conservation laws,

## References