

# Package ‘CropPhenology’

September 19, 2017

**Type** Package

**Title** Extract phenologic metrics from timeseries vegetation index data

**Version** 0.1.0

**Author** Sofanit Araya, Bertram Ostendorf, Megan Lewis and Greg Lyle

**Maintainer** Sofanit Araya <sofanitgirma.araya@adelaide.edu.au>

**Description** CropPhenology-package

**License** GPL (>=2)

**Encoding** UTF-8

**URL** <https://github.com/SofanitAraya/CropPhenology>,  
<http://cropphenology.wix.com/package>

**Repository** github

**Depends** foreign, raster, sp (>= 1.0-13), maptools, shapefiles,  
rgdal, Rcpp, rgdal, rgeos

**SystemRequirements** R (> 3.0)

**LazyLoad** true

**LazyData** true

**RoxygenNote** 6.0.1

## R topics documented:

CropPhenology . . . . .	2
MultiPointsPlot . . . . .	4
PhenoMetrics . . . . .	5
<b>Index</b>	<b>7</b>

## Description

Extract phenological metrics from time series vegetation index data

## Details

### Introduction

Multi temporal vegetation index data can be used to get information on seasonal vegetation growth dynamics. This information indicates vegetation phenological growth stages and conditions of environmental factors influencing the vegetation growth. In cropping regions the crop growth dynamics observed from multi temporal vegetation index data has been used in applications such as crop type detection (Zhong et.al. 2011, Roerink et.al. 2011), regional crop yield estimation (Hill et.al. 2003) and many more related studies. Moreover, the long term vegetation dynamics can provide information about influential environmental factors such as soil property mapping (Araya et.al. 2016). Plotting a time series of vegetation index values across time creates a curve that summarises the vegetation dynamics (Figure 1). Extraction of seasonal parameters is an essential step for analysing such vegetation dynamics curve. CropPhenology package has been developed to extract phenological parameters from time series vegetation index data in cropping regions.




Figure 1 – Illustration for vegetation dynamics derived from multi temporal vegetation index data, and phenological metrics derived from vegetation dynamics using CropPhenology package

### Overview of data processing

CropPhenology has two functions: PhenoMetrics and MultiPointsPlots. PhenoMetrics:- takes the path for the time series vegetation index data and the vector file that defines the Area of Interest (AOI). It extracts fifteen phenological metrics (Figure 2) which represent the seasonal growth condition of the crop at each pixel for the season. The output is presented as a raster stack of phenological metrics or a table of phenological metrics for point AOI. Table 1 summarises the defined metrics and their descriptions.




Figure 2 –NDVI dynamics curve with representation of phenological metrics

Metrics	Definition on the NDVI curve, Formula, and description
OnsetV (in NDVI value)	NDVI value measured at the start of continuous positive slope over a threshold between successive NDVI values. The threshold is defined as user defined percentage above the minimum NDVI value before the Maximum value.
OnsetT (in MODIS image period)	MODIS acquisition time when OnsetV is derived.
MaxV (in NDVI value)	Maximum NDVI value achieved during the season $MaxV = \text{Maximum (NDVI1 : NDVI23)}$
MaxT (in MODIS imaging period)	MODIS acquisition time when MaxV is derived.
OffsetV (in NDVI value)	NDVI value measured at the lowest slope below a threshold between successive NDVI values. The threshold is defined as the user defined percentage of the minimum NDVI value after maximum. Values are higher than 0.2.
OffsetT (in MODIS imaging period)	MODIS acquisition period when OffsetV is derived.
LengthGS (in MODIS imaging period)	The duration of time that the crop takes to go through all the stages of crop growth $LengthGS = OffsetT - OnsetT$
BeforeMaxT (in MODIS image period)	The length of time from OnsetT to the MaxT $BeforeMaxT = MaxT - OnsetT$
AfterMaxT (in MODIS image period)	The length of time from MaxT and OffsetT $AfterMaxT = OffsetT - MaxT$
GreenUpSlope	The rate at which NDVI increases from the OnsetV to MaxV over the time difference between MaxT and OnsetT $GreenUpSlope = \frac{(MaxV - OnsetV)}{(MaxT - OnsetT)}$
BrownDownSlope	The rate at which NDVI decreases from MaxV to OffsetV over the difference between OffsetT and MaxT. $BrownDownSlope = \frac{(MaxV - OffsetV)}{(OffsetT - MaxT)}$
TINDVI (Accumulated NDVI value)	Area under the NDVI curve between OnsetT and OffsetT. TINDVI is estimated using trapezoidal numerical integration.
TINDVIBeforeMax (Accumulated NDVI value)	Numerical integration of NDVI between OnsetT and MaxT. This metric indicates the pre-anthesis crop growth.
TINDVIAfterMax (Accumulated NDVI value)	Numerical integration of NDVI between MaxT and OffsetT. This metric indicates the post-anthesis growth.
Asymmetry (in NDVI value)	The symmetry of the NDVI curve. It measures which part of the growing season attain relatively higher accumulated NDVI values. $Asymmetry = TINDVIBeforeMax - TINDVIAfterMax$

Table 1 – summary of descriptions of the phenological metrics defined in CropPhenology

**MultiPointsPlots:-** provides the user with the ability to visualise the NDVI curve by plotting the temporal sequences of NDVI values of user selected raster pixels (maximum of five). This allows the user to observe the spatial and temporal differences in relative dynamics of the vegetation index for the selected points. Figure 3 shows example for the output of MultiPointsPlots.




Figure 3 – Illustration for NDVI dynamics from 3 locations plotted together using MultiPointsPlots

## References

- Araya, S., Lyle, G., Lewis, M., and Ostendorf, B. 2016. Phenologic metrics derived from MODIS NDVI as indicators for Plant Available Water-holding Capacity. *Ecological Indicators* **60**:1263-1272. <http://dx.doi.org/10.1016/j.ecolind.2015.09.012>
- Hill, M. J. and Donald, G. E. 2003. Estimating spatio-temporal patterns of agricultural productivity in fragmented landscapes using AVHRR NDVI time series. *Remote Sensing of Environment* **84**:367-384. DOI:10.1016/s0034-4257(02)00128-1
- Roerink, G. J., Danes, M. H. G. I., Prieto, O. G., De Wit, A. J. W., and Van Vliet, A. J. H. 2011. Deriving plant phenology from remote sensing. in *6th International Workshop on the Analysis of Multi-temporal Remote Sensing Images*, Trento, Italy, **Pages 261-264**
- Zhong, L., Hawkins, T., Biging, G., Gong, P., 2011. A phenology-based approach to map crop types in the San Joaquin Valley, California. *International Journal of Remote Sensing*, **32**, 7777-7804.

---

MultiPointsPlot

*Time series curves for Multiple points in the Region of Interest*

---

## Description

MultiPointsPlot function takes the ID for the pixels within the region of interest and returns, the timeseries curves from these points, plotted together. The Id numbers can be obtained from the txt file (AllPixels.txt) outputs.

## Usage

```
MultiPointsPlot(path, N, Id1, Id2, Id3, Id4, Id5)
```

## Arguments

path	- the path where AllPixel.txt saved
N	- number of interested points
Id1	- ID number for point 1
Id2	- Id number for point 2
Id3	- ID number for point 3
Id4	- ID number for point 4
Id5	- ID number for point 5

**Details**

This function allows plotting time series curves from multiple points together in a single plot which helps understanding the growth variability across the field. The maximum number of pixels allowed plotting together are 5 points.

**Value**

Multiple time series curves together at the plot panel

**See Also**

PhenoMetrics()

---

PhenoMetrics	<i>Phenologic metrics from time series vegetation index data</i>
--------------	--

---

**Description**

This function extracts 15 phenologic metrics from time series vegetation index data, as raster and Ascii files. The function takes path of the vegetation index data and the boolean Value for BolAOI (True- if there is AOI polygon, FALSE- if the parameters are calculated for the whole region).

**Usage**

```
PhenoMetrics(Path, BolAOI, Percentage, Smoothing)
```

**Arguments**

Path	- Text value - the path where the time series images saved
BolAOI	- Logical value - if there is any area of interest or not
Percentage	- Optional Numeric Value - percentage of minimum NDVI value at which the Onset and Offset is defined. The 'Percentage' parameter is optional; if not provided, a Default value of 10 will be taken.
Smoothing	- Optional logical value - if the user chooses to use smoothed curve or raw/unsmoothed curve. If "Smoothing" is set to TRUE, the moving average filter will be applied to the vegetation index curve. The default value, if not provided, is FALSE, then the unsmoothed raw data be used for the analysis.

**Value**

PhenoStack.img - a raster stack of 15 images in the order of OnsetV, OnsetT, MaxV, MaxT, OffsetV, OffsetT, LengthGS, BeforeMaxT, AfterMaxT, GreenUpSlope, BrownDownSlope, TINDVI, TINDVIBeforeMax, TINDVIAfterMax, Asymmetry

**See Also**

MultiPointsPlot (Path, N, Id1, Id2...IdN)

**Examples**

```
#EXAMPLE - 1
```

```
#PhenoMetrics(system.file("extdata/data1", package="CropPhenology"), FALSE, 15, TRUE)
```

```
#EXAMPLE - 2
```

```
# PhenoMetrics(system.file("extdata/data2", package="CropPhenology"), TRUE)
```

# Index

- \*Topic **Curve**
  - MultiPointsPlot, [4](#)
- \*Topic **Phenology**
  - PhenoMetrics, [5](#)
- \*Topic **Time-series**
  - PhenoMetrics, [5](#)
- \*Topic **curves**
  - MultiPointsPlot, [4](#)
- \*Topic **from**
  - MultiPointsPlot, [4](#)
- \*Topic **image**
  - PhenoMetrics, [5](#)
- \*Topic **multiple**
  - MultiPointsPlot, [4](#)
- \*Topic **package**
  - CropPhenology, [2](#)
- \*Topic **points**
  - MultiPointsPlot, [4](#)
- \*Topic **remote**
  - PhenoMetrics, [5](#)
- \*Topic **satellite**
  - PhenoMetrics, [5](#)
- \*Topic **sensing**
  - PhenoMetrics, [5](#)
- \*Topic **time-series**
  - MultiPointsPlot, [4](#)

CropPhenology, [2](#)

MultiPointsPlot, [4](#)

PhenoMetrics, [5](#)