Mathematical Modeling

Task Sheet 1: Complementarity and Two-Phase Simplex

Personal study notes

Contents

1 How to read the certificates and tableaus **1**

2 Task 2: complementary slackness **2**
2.1 Task 2(a): a feasible candidate and a matching upper bound 2
2.2 Task 2(b): reject the supplied point, then solve the dual 2
2.3 Task 2(c): mixed row signs and an optimal segment 3

3 Task 3: dual problems and two-phase simplex **3**
3.1 Task 3(a): a surplus variable enters during Phase I 4
3.2 Task 3(b): a nonpositive variable and negative right-hand sides 5

4 Checks and questions for understanding **6**

5 Sources and editorial remarks **7**

1 How to read the certificates and tableaus

Feasibility comes before optimality. A proposed point must satisfy every row and every variable-domain restriction. Complementary products alone cannot certify an inadmissible point.

For a maximization problem, multiply each $\leq$ row by a nonnegative weight and each $\geq$ row by a nonpositive weight. Summing produces $y^T Ax \leq y^T b$. With $x \geq 0$, the column conditions $A^T y \geq c$ then give $c^T x \leq b^T y$: the dual minimizes a valid upper bound. For minimization with $Ax \geq b$, $x \geq 0$, nonnegative weights and $A^T y \leq c$ instead give the lower bound $b^T y \leq c^T x$. If a variable is nonpositive, its column comparison reverses when multiplied by that variable. The signs follow from these comparisons, rather than from a rule to reverse every symbol indiscriminately.

Once primal and dual points are feasible, equal objective values prove optimality. Complementarity helps find the dual point: a strict primal row forces its weight to zero, and a nonzero primal coordinate forces its dual column inequality to be an equality. For mixed rows, use their nonnegative slacks and the corresponding signed weights consistently. These ideas are developed in Degeneracy, duality, and dual simplex, in the sections on weighted bounds and complementary slackness.

For each simplex calculation below, use a maximization problem $\max d^T v$ with $Mv = b$, $v \geq 0$. An ordered basis B produces

$$\bar{M} = M_B^{-1} M, \qquad \bar{b} = M_B^{-1} b, \qquad q_j = d_B^T \bar{m}_j - d_j, \qquad z_0 = d_B^T \bar{b}.$$

Tableau rows mean $\bar{M}v = \bar{b}$ and $z + q^T v = z_0$. A negative q_k offers improvement; increasing v_k by θ changes the basic coordinates to $\bar{b} - \theta \bar{m}_k$. Only positive entries of $\bar{m}_k$ limit the step, with $\theta^* = \min_{i:\bar{m}_{ik}>0} \bar{b}_i / \bar{m}_{ik}$. Choose a most negative q_k, breaking entering ties by the smaller column index. The displayed paths have no leaving tie. The derivations appear in The simplex method, in the sections on canonical equations, one simplex step, and initialization.

2 Task 2: complementary slackness

Korzystając z zasady komplementarności sprawdzić czy podane rozwiązanie jest rozwiązaniem optymalnym. Podać rozwiązanie dualne.

2.1 Task 2(a): a feasible candidate and a matching upper bound

Task statement.

$$\max(4x_1 + 5x_2 - 8x_3), \qquad \begin{cases} 4x_1 + x_2 - 2x_3 \le 1, \\ 2x_1 + x_2 - 2x_3 \le 6, \\ -x_1 + x_2 - x_3 \le 2, \\ x_1 + x_2 - x_3 \le 3, \\ x_1, x_2, x_3 \ge 0, \end{cases} \qquad x = (0, 3, 1).$$

The candidate is nonnegative, and its row values are $(1, 1, 2, 2)^T$. The slacks are $(0, 5, 0, 1)^T$, so it is feasible, with value 7. There is one nonnegative dual weight per upper-bound row:

$$\begin{aligned} \min \quad & y_1 + 6y_2 + 2y_3 + 3y_4, \\ 4y_1 + 2y_2 - y_3 + y_4 &\ge 4, \\ y_1 + y_2 + y_3 + y_4 &\ge 5, \\ -2y_1 - 2y_2 - y_3 - y_4 &\ge -8, \qquad y \ge 0. \end{aligned}$$

Strict primal rows force $y_2 = y_4 = 0$. Since $x_2, x_3 > 0$, the last two dual columns must be tight:

$$y_1 + y_3 = 5, \qquad -2y_1 - y_3 = -8.$$

Thus $y = (3, 0, 2, 0)^T$. The remaining column equals $10 \ge 4$; all dual signs hold. Its value is $3 + 2 \cdot 2 = 7$, matching the primal. The dual surplus is $(6, 0, 0)^T$, so all complementary products vanish. Both points are optimal.

This also proves uniqueness of the primal optimum. The positive first-column surplus forces $x_1 = 0$; the positive weights y_1, y_3 force rows 1 and 3 to be tight. Solving $x_2 - 2x_3 = 1$, $x_2 - x_3 = 2$ gives $(x_2, x_3) = (3, 1)$.

2.2 Task 2(b): reject the supplied point, then solve the dual

Task statement.

$$\min(6x_1 + 2x_2 - 3x_3), \qquad \begin{cases} x_1 - x_2 - x_3 \ge -1, \\ -3x_2 - 2x_3 \ge -4, \\ x_1 - x_2 + x_3 \ge 4, \\ x_1, x_2, x_3 \ge 0, \end{cases} \qquad x = (2, 0, -2).$$

The supplied point has $x_3 = -2 < 0$ and its third row is $0 < 4$. It is infeasible, hence not optimal. We retain the stated point; replacing its sign would answer a different candidate question.

The task also requests a dual solution. Nonnegative weights on the three lower-bound rows yield

$$\begin{aligned} \max \quad & -y_1 - 4y_2 + 4y_3, \\ y_1 + y_3 &\le 6, \\ -y_1 - 3y_2 - y_3 &\le 2, \\ -y_1 - 2y_2 + y_3 &\le -3, \qquad y \ge 0. \end{aligned}$$

To find a feasible primal witness, the second row gives $3x_2 + 2x_3 \le 4$, hence $x_3 \le 2$. The third row gives $x_1 \ge 4 + x_2 - x_3$, and therefore

$$6x_1 + 2x_2 - 3x_3 \ge 24 + 8x_2 - 9x_3 \ge 6.$$

The point $\hat{x} = (2, 0, 2)^T$ attains 6; its row values are $(0, -4, 4)^T$, so it is feasible. Its positive first-row surplus forces $y_1 = 0$. Its positive coordinates $\hat{x}_1, \hat{x}_3$ force

$$y_3 = 6, \qquad -2y_2 + y_3 = -3,$$

giving $y = (0, 9/2, 6)^T$. Its second column is $-39/2 \le 2$; the other columns are tight. The dual value is $-18 + 24 = 6$. Thus this is an optimal dual solution, and $\hat{x}$ is the actual primal optimum. Equality in the elementary bound above forces $x_2 = 0$, $x_3 = 2$, and $x_1 = 2$, proving primal uniqueness.

2.3 Task 2(c): mixed row signs and an optimal segment

Task statement.

$$\max(2x_1 + x_2 - 3x_3 + x_4), \qquad \begin{cases} -x_1 + x_2 + x_3 + x_4 \ge 3, \\ 2x_1 + x_2 + 2x_3 + 2x_4 \le 10, \\ x_1 + x_2 - x_3 + x_4 \le 2, \\ x_1, x_2, x_3, x_4 \ge 0, \end{cases} \qquad x = (0, 5/2, 1/2, 0).$$

The row values are $(3, 7/2, 2)^T$. The nonnegative slacks, in the order "left minus right" for row 1 and "right minus left" for rows 2–3, are $(0, 13/2, 0)^T$. The candidate is feasible and has value 1.

For an upper bound, row 1 needs a nonpositive weight, whereas rows 2–3 need nonnegative weights:

$$\begin{aligned} \min \quad & 3y_1 + 10y_2 + 2y_3, \\ -y_1 + 2y_2 + y_3 &\ge 2, \\ y_1 + y_2 + y_3 &\ge 1, \\ y_1 + 2y_2 - y_3 &\ge -3, \\ y_1 + 2y_2 + y_3 &\ge 1, \\ y_1 \le 0, \qquad & y_2, y_3 \ge 0. \end{aligned}$$

The strict second row forces $y_2 = 0$. Positive x_2, x_3 require $y_1 + y_3 = 1$ and $y_1 - y_3 = -3$, hence $y = (-1, 0, 2)^T$. The four dual column values are $(3, 1, -3, 1)^T$, so the surplus is $(1, 0, 0, 0)^T$. Its value is $-3 + 4 = 1$: both points are optimal.

The positive first-column surplus forces every primal optimum to have $x_1 = 0$. Nonzero y_1, y_3 force rows 1 and 3 to be tight, giving $x_3 = 1/2$ and $x_2 + x_4 = 5/2$. The second row then equals $7/2 + x_4$, which remains below 10 for $0 \le x_4 \le 5/2$. Thus the entire optimal set is

$$x(t) = (0, 5/2 - t, 1/2, t)^T, \qquad 0 \le t \le 5/2.$$

The supplied point is one endpoint, not the unique optimum. A zero dual surplus alone would not prove this; the displayed nonzero feasible changes do.

3 Task 3: dual problems and two-phase simplex

Podaj ZD(y). Rozwiąż metodą 2-fazową. Here ZD denotes the dual problem. Artificial variables provide a feasible basis for an auxiliary system; they are not allowed to change the final model. Phase I maximizes minus their sum, which is never positive. An optimum of zero allows their removal and a return to the original objective. A negative optimum would prove the original constraints infeasible.

3.1 Task 3(a): a surplus variable enters during Phase I

Task statement.

$$\max(-x_1 - 2x_2 + x_3), \qquad \begin{cases} x_1 + x_3 \le 12, \\ x_1 + 3x_2 + x_3 \ge 4, \\ x_1 + 2x_2 - x_3 \ge 6, \\ x_1, x_2, x_3 \ge 0. \end{cases}$$

Dual from an upper bound. The first row receives $y_1 \ge 0$ and the two lower-bound rows receive $y_2, y_3 \le 0$. The dual is

$$\begin{aligned} \min \quad & 12y_1 + 4y_2 + 6y_3, \\ y_1 + y_2 + y_3 &\ge -1, \\ 3y_2 + 2y_3 &\ge -2, \\ y_1 + y_2 - y_3 &\ge 1, \\ y_1 &\ge 0, \qquad y_2, y_3 \le 0. \end{aligned}$$

Initialization. Add slack s_1, surpluses s_2, s_3, and artificials a_2, a_3:

$$\begin{aligned} x_1 + x_3 + s_1 &= 12, \\ x_1 + 3x_2 + x_3 - s_2 + a_2 &= 4, \\ x_1 + 2x_2 - x_3 - s_3 + a_3 &= 6. \end{aligned}$$

All eight variables are nonnegative. Use column order $(x_1, x_2, x_3, s_1, s_2, s_3, a_2, a_3)$, initial basis $B = (4, 7, 8)$, and Phase I objective $w = -a_2 - a_3$. Surplus columns point in the negative unit direction and cannot supply nonnegative initial basic values for positive right-hand sides.

Phase I: initial tableau. Substitution gives $w = -10 + 2x_1 + 5x_2 - s_2 - s_3$; the indicator row must already account for both basic artificial variables.

B	x_1	x_2	x_3	s_1	s_2	s_3	a_2	a_3	$\bar{b}$
s_1	1	0	1	1	0	0	0	0	12
a_2	1	3	1	0	−1	0	1	0	4
a_3	1	2	−1	0	0	−1	0	1	6
q	−2	−5	0	0	1	1	0	0	−10

Enter x_2. Ratios are $4/3$ and $6/2 = 3$, so a_2 leaves with $\theta = 4/3$. Divide its row by 3 and eliminate column x_2:

B	x_1	x_2	x_3	s_1	s_2	s_3	a_2	a_3	$\bar{b}$
s_1	1	0	1	1	0	0	0	0	12
x_2	1/3	1	1/3	0	−1/3	0	1/3	0	4/3
a_3	1/3	0	−5/3	0	2/3	−1	−2/3	1	10/3
q	−1/3	0	5/3	0	−2/3	1	5/3	0	−10/3

Now s_2 has the most negative indicator. Its only positive entry is $2/3$ in row a_3, so a_3 leaves with $\theta = (10/3)/(2/3) = 5$. There is no rule restricting entry to original decision variables: a surplus is a legitimate variable in the equivalent system.

B	x_1	x_2	x_3	s_1	s_2	s_3	a_2	a_3	$\bar{b}$
s_1	1	0	1	1	0	0	0	0	12
x_2	1/2	1	−1/2	0	0	−1/2	0	1/2	3
s_2	1/2	0	−5/2	0	1	−3/2	−1	3/2	5
q	0	0	0	0	0	0	1	1	0

Phase I has value zero and both artificials are nonbasic. Delete their columns; the remaining basis $B = (4, 2, 5)$ already belongs to the original system.

Phase II: restore the original objective. Recompute the indicator row for the original coefficients:

$$d = (-1, -2, 1, 0, 0, 0)^T.$$

B	x_1	x_2	x_3	s_1	s_2	s_3	$\bar{b}$
s_1	1	0	1	1	0	0	12
x_2	1/2	1	−1/2	0	0	−1/2	3
s_2	1/2	0	−5/2	0	1	−3/2	5
q	0	0	0	0	0	1	−6

All indicators are nonnegative, so no Phase II pivot is needed. The primal solution is $x^* = (0, 3, 0)^T$, with slacks/surpluses $(12, 5, 0)^T$ and maximum -6. Indeed, the original third constraint immediately gives $-x_1 - 2x_2 + x_3 \le -6$. The dual point $y^* = (0, 0, -1)^T$ has the required signs, satisfies every column with equality, and has value -6.

There are alternative optima: $x(t) = (t, 3 - t/2, 0)^T$ is feasible and has value -6 for every $0 \le t \le 6$. This can also be read from the zero x_1 indicator and its positive allowable step from the final basis.

3.2 Task 3(b): a nonpositive variable and negative right-hand sides

Task statement.

$$\min(2x_1 - 2x_2 + x_3), \qquad \begin{cases} x_1 - x_2 + x_3 \ge -1, \\ x_1 + 2x_2 + 2x_3 \ge -4, \\ x_1 - x_2 - x_3 \ge 4, \\ x_1 \ge 0, \quad x_2 \ge 0, \quad x_3 \le 0. \end{cases}$$

Dual in the original variables. All lower-bound rows have nonnegative weights. The first two variables are nonnegative, so their column comparisons are $\le$; the nonpositive x_3 requires $\ge$. Explicitly,

$$\begin{aligned} \max \quad & -y_1 - 4y_2 + 4y_3, \\ y_1 + y_2 + y_3 &\le 2, \\ -y_1 + 2y_2 - y_3 &\le -2, \\ y_1 + 2y_2 - y_3 &\ge 1, \qquad y \ge 0. \end{aligned}$$

The last comparison ensures that $(y_1 + 2y_2 - y_3)x_3 \le x_3$ because $x_3 \le 0$.

Transform and initialize. Set $t = -x_3 \ge 0$, so the original objective becomes $F = 2x_1 - 2x_2 - t$. We maximize its negative $z = -F = -2x_1 + 2x_2 + t$. Multiply the first two constraint rows by -1 to obtain positive right-hand sides; their inequality directions reverse:

$$\begin{aligned} -x_1 + x_2 + t + s_1 &= 1, \\ -x_1 - 2x_2 + 2t + s_2 &= 4, \\ x_1 - x_2 + t - s_3 + a_3 &= 4. \end{aligned}$$

Here s_1, s_2 are slacks of the reversed rows, s_3 a surplus, and a_3 artificial. Use column order $(x_1, x_2, t, s_1, s_2, s_3, a_3)$ and initial basis $B = (4, 5, 7)$. Phase I maximizes $w = -a_3$. The two negative original right-hand sides do not require artificials after the row reversal; their slack columns provide a feasible start.

Phase I. The initial tableau is

B	x_1	x_2	t	s_1	s_2	s_3	a_3	$\bar{b}$
s_1	-1	1	1	1	0	0	0	1
s_2	-1	-2	2	0	1	0	0	4
a_3	1	-1	1	0	0	-1	1	4
q	-1	1	-1	0	0	1	0	-4

Both x_1 and t have indicator -1; the smaller column index chooses x_1. Its only positive coefficient is in row a_3, so a_3 leaves with $\theta = 4$:

B	x_1	x_2	t	s_1	s_2	s_3	a_3	$\bar{b}$
s_1	0	0	2	1	0	-1	1	5
s_2	0	-3	3	0	1	-1	1	8
x_1	1	-1	1	0	0	-1	1	4
q	0	0	0	0	0	0	1	0

The zero Phase I value and nonbasic artificial allow its column to be removed.

Phase II. For the negative original objective use $d = (-2, 2, 1, 0, 0, 0)^T$:

B	x_1	x_2	t	s_1	s_2	s_3	$\bar{b}$
s_1	0	0	2	1	0	-1	5
s_2	0	-3	3	0	1	-1	8
x_1	1	-1	1	0	0	-1	4
q	0	0	-3	0	0	2	-8

Enter t. The ratios are $5/2$, $8/3$, and 4, so s_1 leaves with $\theta = 5/2$. Normalize its row and eliminate the t column:

B	x_1	x_2	t	s_1	s_2	s_3	$\bar{b}$
t	0	0	1	$1/2$	0	$-1/2$	$5/2$
s_2	0	-3	0	$-3/2$	1	$1/2$	$1/2$
x_1	1	-1	0	$-1/2$	0	$-1/2$	$3/2$
q	0	0	0	$3/2$	0	$1/2$	$-1/2$

All indicators are nonnegative, proving the maximum of z is $-1/2$. Recover $x_3 = -t$ and $F = -z$: the original solution and minimum are

$$x^* = (3/2, 0, -5/2)^T, \qquad F^* = 1/2.$$

Its original row values are $(-1, -7/2, 4)^T$, with surpluses $(0, 1/2, 0)^T$. The positive second surplus forces $y_2 = 0$. Nonzero x_1, x_3 force $y_1 + y_3 = 2$ and $y_1 - y_3 = 1$, yielding $y^* = (3/2, 0, 1/2)^T$. It is dual feasible, including the tight second column, and has value $-3/2 + 2 = 1/2$.

The final objective row also gives $F = 1/2 + (3/2)s_1 + (1/2)s_3$. Optimality therefore forces $s_1 = s_3 = 0$. The remaining equations allow

$$x(\tau) = (3/2 + \tau, \tau, -5/2)^T, \qquad \tau \geq 0,$$

with $s_2 = 1/2 + 3\tau$. This is the entire optimal set, an unbounded ray with a finite constant objective. An unbounded feasible set does not imply an unbounded objective.

4 Checks and questions for understanding

All results above are exact rational calculations. The companion MATLAB script checks the original variable domains, row residuals, primal–dual objective equality, complementary products,

and every displayed tableau. It follows the specified bases and verifies entering choices, ratio tests, step lengths, and the original-coordinate recovery; it is a check of these hand solutions rather than a general LP solver. Its absolute tolerance is 10^{-10} for these small, modestly scaled inputs. No units are specified by the task statements; objective values are reported without invented physical units.

1. Which two independent feasibility tests reject the supplied point in 2(b)?
2. Why is $y_1 \leq 0$ in 2(c), but $y_1 \geq 0$ in 3(b)?
3. Why may a surplus enter the basis in 3(a)?
4. Why must the objective row be rebuilt between phases?
5. Explain separately the row reversal, variable substitution, and objective negation in 3(b). Which changes the final reported objective sign?
6. Use the displayed segment or ray to prove alternatives exist; explain why a zero indicator alone is insufficient.

5 Sources and editorial remarks

Task source: `ProgMatc1.pdf`, titled Programowanie Matematyczne zestaw zadań 1, PDF page 1, group 2, and PDF page 2, group 3, supplied in the `execrices-2026-27/` materials directory. The Polish instructions, coefficients, inequality directions, variable domains, and supplied candidates are retained in Sections 2–3. English explanations, dual derivations, complete tableaus, and descriptions of the optimal sets are added study material. No lecture date is inferred.

Numerical verification. The companion MATLAB script can be run from the repository root with

```
run('matlab/exercises/sheet_01_complementarity_and_two_phase.m')
```

MATLAB execution is not yet verified for this script. The tableau calculations and certificates were independently checked with exact rational arithmetic; this does not constitute MATLAB execution evidence. Actual execution status, release, and output belong in the repository verification record when available.