स्वच्छ भारत
एक कदम स्वच्छता की ओर

सत्यमेव जयते

Phone: 0129-2413985
E-mail:-ppa@nic.in

F.No 3-6/2022-23/IPM/Advisory
भारत सरकार/ Government of India
कृषि एवं किसान कल्याण मंत्रालय / Ministry of Agriculture & Farmers Welfare
कृषि एवं किसान कल्याण विभाग / Department of Agriculture & Farmers Welfare
वनस्पति संरक्षण, संगरोध एवं संग्रह निदेशालय / Directorate of Plant Protection, Quarantine & Storage
एन एच4, फरीदाबाद (हरियाणा) – 121 001/ NH-IV, Faridabad (Haryana) – 121 001

Dated: 25.09.2023

To

The Director of Horticulture
Govt. of Odisha, Punjab and Haryana

Sub: Advisory for management of Rice leaf folder (*Cnaphalocrocis medinalis)* on rice crop -reg.

Sir,

It is to bring to your notice that incidence of Rice leaf folder (*Cnaphalocrocis medinalis*) on rice crop is reported in the Odisha, Punjab and Haryana region. The pest was reported in Faridkot districts of Punjab, Cuttak and Dhrnkanal districts of Odisha and Sonipat district of Haryana at moderate intensity (Area affected 2900 ha in Punjab). Keeping in view the current crop stage and prevailing weather condition, Integrated Pest Management techniques should be adopted. IPM Package of practices of Rice is available at DPPQ&S website in the link https://ppqs.gov.in/ipm-packages. Important IPM for management of Rice Leaf Folder are listed below

1. Low usage of nitrogenous fertilizers decreases leaf damage. Highly fertilized plots seem to attract females for oviposition. Therefore, it is advisable to avoid over-fertilization.
2. Egg predators (crickets) inhabit surrounding grass habitats and move to the field at night for predation. Maintenance of non-rice habitats.
3. Higher damages will occur in shaded areas. Therefore, remove the causes of shading within the field.
4. Removal and destruction of diseased/pest infested plant parts.
5. Use of coir rope in rice crop for dislodging leaf folder larvae etc. on to kerosinized water (1 L of kerosene mixed on 25 kg soil and broadcast in 1ha).
6. Remove grassy weeds from fields and borders.
7. Flood and plough field after harvesting if possible.
8. Rotate rice with a different crop, or have a fallow period.
9. Avoid ratooning
10. Encourage predators such as spider, parasitic wasps, predatory beetles, frogs and dragon fly.
11. Encourage biological control agents like *Trichogramma japonicum* and *T chilonis* may be released @ 1 lakh/ha on appearance of egg masses / moth of leaf folder in the field, Egg Parasitoid *Copidosomopsis nacoleiae*, Larval Parasitoid *Stenobracon nicevillei*, Larval and Pupal Parasitoid *Brachymeria lasus*, *B. excarinata* etc.

12. Encourage the use of biopesticides/ botanicals to manage the pest so as to avoid it to reach to ETL.

13. Pesticides should be used once the pest reaches ETL andas per label claim approved by the registration committee. Detailed information is available on directorate's website at link https://ppqs.gov.in/sites/default/files/1._major_use_of_pesticides_insecticides_as_on_01.06.2023.pdf A list of recommended pesticides and biopesticides for rice leaf Folder is attached at **Annexure-I.**

Further the following strategies should be adopted for spreading awareness among farmers

- Mobilization of extension functionaries at grass route level for regular monitoring of pest status.
- Regular meetings of State Pest Surveillance and advisory unit and District Pest Surveillance and Advisory Unit may be conducted
- Awareness among farming community through newspapers, printing and electronic media, Kisan Gosthi, leaflet and pamphlets.
- The concerned officials at the district/ division/ block level may be advised to take the suitable plant protection measures immediately

The detailed information on pest incidence may also be sent to the directorate through email to ppa@nic.in or appa.ipm-dppqs@gov.in urgently to apprise the same to Department of Agriculture & Farmers' Welfare, MoA&FW, Government of India.

Yours faithfully

(Dr.J.P.Singh)
Plant Protection Adviser

Copy for information:

1. PPS to Joint Secretary(PP), DA& FW, Krishi Bhawan, New Delhi- I 10001
2. The Directors of Agriculture/Horticulture of other rice growing states with the request to keep close watch on pest build up & take appropriate measures as indicated above
3. PPS to PPA for information
4. Officer In-charges, all CIPMCs of rice growing states with directions to conduct regular survey in the area to ensure and necessary follow-up with the state authorities.

Annexure-I

CIBRC Approved registered Pesticides for Rice Leaf Folder

Name	Dosage/ha Formulation(gm/ml)	Dilution in water (Liter)	Waiting period in days
Acephate 75 % SP	666-1000	300-500	15
Acephate 97 % DF	750	500	21
Acephate 95 % SG	592	500	30
Benfuracarb 03 % GR	33000	-	20
Bifenthrin 08.80 % CS	500	500	21
Bifenthrin 10 % EC	500	500	21
Carbofuran 03 % CG	25000	-	-
Carbosulfan 06 % Granules	16700	-	37
Carbosulfan 25% EC	800-1000	500-1000	14
Cartap Hydrochloride 04 % Granules	18750	-	-
Cartap Hydrochloride 50 % SP	1000	500-1000	-
Cartap Hydrochloride 75 % SG	425-500	250-500	35-89
Chlorantraniliprole 18.50 % SC	150	500	47
Chlorantraniliprole 00.40 % GR	10000	-	53
Chlorantraniliprole 50% W/w FS	120	Seed Dresser	-
Chlorpyrifos 10 % Granules	10000	-	30
Chlorpyrifos 20 % EC	1875	500-1000	-
Chlorpyrifos 50 % EC	750-800	500-1000	15
Chlorpyriphos 75 % w/w WG	500-533	500-1000	15
Chlorpyrifos 01.50 % DP	25000	-	07
Chromafenozide 80 % WP	94-125	500	32
Cyclaniliprole 9.3% W/W DC	350-400	500	40
Deltamethrin 11% w/w EC	150	500	13
Deltamethrin 01.80 % EC	625 – 780	500	07
Ethofenoprox 10 % EC	500-750	500	15
Fenpropathrin 30 % EC	333	500	30
Fenpropathrin 10% EW	1000	5000	58
Fipronil 05 % SC	1000-1500	500	32
Fipronil 18.87 % w/w SC	250	500	46
Fipronil 00.30 % GR	16670-25000	-	32
Fipronil 00.60 % w/w GR	10	-	65

Fipronil 80 % WG	50-62.50	375-500	19
Flubendiamide 20 % WG	125	500	30
Flubendiamide 39.35 % w/w SC	50	375-500	40
Flubendiamide 00.70 % GR	12.14-14.28	-	25
Flupyrimin 2% GR	5000-7500	--	77
Isocycloseram 18.1% W/W SC (20 % W/V Sc)	100	500	48
Lambda-cyhalothrin 04.90 % CS	250	500	15
Lambda-cyhalothrin 02.50 % EC	500	400-600	15
Lambda-cyhalothrin 05 % EC	250	400-600	15
Monocrotophos 36 % SL	625	500-1000	-
Phosmet 50% WP	1200	500	45
Quinalphos 25 % Gel	1000	500-1000	-
Quinalphos 05 % Granules	5000	-	-
Quinalphos 20 % AF	1250-1500	750-1000	40
Quinalphos 25 % EC	1000	500-1000	40
Tetraniliprole 18.18 SC	250-300	500	43
Thiacloprid 21.70 % SC	500	500	30
Thiocyclam Hydrogen Oxalate 50% SP	1000	500	30
Thiamethoxam 25 % WG	100	500-750	14
Azoxystrobin 10.0% + Fipronil 5% SC	1250	500	53
Cartap Hydrochloride 50 % + Buprofezin 10 % w/w WP	800	500	20
Cartap Hydrochloride 7.5 % w/w + Emamectin benzoate 0.25 %w/w GR	750	-	35
Fipronil 5% + Buprofezin 20% SC	1000	500	20
Flubendiamide 04 % + Buprofezin 20 % w/w SC	175+700	500	30
Flubendiamide 19.92 % + Thiacloprid 19.92 % w/w SC	250	500	33
Indoxacarb 10.0% + Thiamethoxam 10.0% WG	500	50	14
Phenthoate 45% + Cypermethrin 6% EC	1000	500	At the end of harvest

CIBRC approved biopesticides for management of Rice Leaf Folder

Name	Dosage/ha Formulation (gm/ml)	Dilution in water (Liter)	Waiting period in days
Azadirachtin 0.15% EC w/w Min. Neem Seed Kernel Based	1500-2000	500	05
Azadirachtin 05.00% w/w Min. Neem Extract Concentrates	200	400	05
***Bacillus thuringiensis* var. galleriae 1593 M serotype H 59 5b, 1.3% flowable concentrate Potency 1500 IU/mg**	1-3	1000	
***Bacillus thuringiensis* var. *kurstaki*, serotype H-39, 3B, Strain Z-52**	500-750	-	-
***Bacillus thuringiensis* var. kurstaki 2.5% AS (Spicbio-BTK AS)**	1.50	500-700	
***Beauveria bassiana* 01.15% WP**	2.50 kg/ha	750-850	-