Annual Review

Michalis Panayides


2020-06-10


Game Theory - Syllabus




- ► Normal Form Games
- ► Mixed-Strategy Nash Equilibrium
- ► Alternate Solution Concepts
- ► Extensive-Form Games
- ► Repeated Games (TBC)
- ► Bayesian Games (TBC)
- ► Coalitional Games (TBC)









$$u_i = A_1 \times A_2 \times A_3 \times \cdots \times A_N$$

Rock-Paper-Scissors





$$\begin{bmatrix} (0,0) & (1,-1) & (-1,1) \\ (-1,1) & (0,0) & (1,-1) \\ (1,-1) & (-1,1) & (0,0) \end{bmatrix}$$

$$\begin{bmatrix} (3,3) & (0,5) \\ (5,0) & (1,1) \end{bmatrix}$$

$$\begin{array}{ccc}
\mathsf{C} & \longrightarrow & \begin{bmatrix} (3,-) & (0,-) \\ (5,-) & (1,-) \end{bmatrix}
\end{array}$$

$$\begin{array}{cccc}
(-,3) & (-,5) \\
(-,0) & (-,1)
\end{array}$$

$$D \longrightarrow \begin{bmatrix} (3,3) & (0,5) \\ (5,0) & (1,1) \end{bmatrix}$$

Pareto Optimality

$$\begin{bmatrix}
(3,3) & (0,5) \\
(5,0) & (1,1)
\end{bmatrix}$$

$$(3,3), (0,5), (5,0), (1,1)$$

Pareto Optimality

$$\begin{bmatrix}
(3,3) & (0,5) \\
(5,0) & (1,1)
\end{bmatrix}$$

$$(3,3), (0,5), (5,0), (1,1)$$

(3,3) > (1,1)

Computing the Nash Equilibria

- ► Lemke-Howson Algorithm
- ► Support Enumeration
- ► Iterative removal of strictly dominated strategies

$$P1 \setminus P2$$
 L C R U $(3,0)$ $(2,1)$ $(0,0)$ M $(1,1)$ $(1,1)$ $(5,0)$ D $(0,1)$ $(4,2)$ $(0,1)$

$$\begin{array}{ccccc} P1 \setminus P2 & L & C \\ U & (3,0) & (2,1) \\ M & (1,1) & (1,1) \\ D & (0,1) & (4,2) \end{array}$$

$$\begin{array}{ccccc} P1 \setminus P2 & L & C \\ U & (3,0) & (2,1) \\ M & \textbf{(1,1)} & \textbf{(1,1)} \\ D & (0,1) & \textbf{(4,2)} \end{array}$$

$$\begin{array}{cccc} P1 \setminus P2 & L & C \\ U & (3,0) & (2,1) \\ D & (0,1) & (4,2) \end{array}$$


$$\begin{array}{cccc} P1 \setminus P2 & L & C \\ U & (3,0) & (2,1) \\ D & (0,1) & (4,2) \end{array}$$


$$\begin{array}{ccc} P1 \setminus P2 & C \\ U & (2,1) \\ D & (4,2) \end{array}$$


$$\begin{array}{ccc} P1 \setminus P2 & C \\ U & (2,1) \\ D & (4,2) \end{array}$$


$$P1 \setminus P2$$
 C D $(4,2)$


































PhD - Motivation





PhD - Motivation






PhD - Motivation






Hospital Formulation



Ambulance - Hospital Interface



Hospital - Markov Chain






Figure: C=4, T=3, N=5, M=2

State Probabilities



Waiting Times



Optimal patient distribution

