1. The probability of a leap year selected at random contain 53						
Sunday is:						
		(c) 2/7				
2. A bag contains	s 3 red and 2 b	olue marbles. A r	marble is drawn at			
random. The prol	bability of drav	wing a black ball	is:			
(a) 3/5	(b) 2/5	(c) $0/5$	(d) 1/5			
3. The probabilit	y that it will ra	in tomorrow is 0	.85. What is the			
probability that it	will not rain to	omorrow				
(a) 0.25	(b) 0.145	(c) 3/20	(d) none of these			
4. What is the pr	obability that a	a number selecte	ed from the numbers			
(1, 2, 3,,15) is a multiple	of 4?				
(a) 1/5	(b) 4/5	(c) 2/15	(d) 1/3			
5. What are the	total outcome	s when we throw	three coins?			
		(c) 8				
6. The probability	ty that a prime	number selecte	d at random from the			
numbers (1,2,3, .						
(a) 12/35	(b) 11/35	(c) 13/35	(d) none of these			
7. The sum of th	-					
• •) 0 (d) non				
8. The following	probabilities	are given; choos	e the correct answer			
for that which is a						
(a) 0.15	(b) 2/7	(c) $7/5$	(d) none of these.			
9. If three coins	are tossed sim	nultaneously, tha	n the probability of			
getting at least tw	vo heads, is:					
(a) 1/4	(b) 3/8	(c) ½	(d) 1/8			
10. A letter is ch						
♦ ASSASSINATION	ON�. The pro	bability that the	letter chosen has: (d) none of these.			
(a) 6/13	(b) 7/13	(c) 1	(d) none of these.			
11. A dice is thro			ting an even number.			
(A) 2/3	(B) 1	(C) 5/6 (D) 1/2			
40 - 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1						
12. Two coins are thrown at the same time. Find the probability of						
getting both head		(D) 0				
(A) 3/4 (B) 1/4	(C) 1/2	(D) 0				
13. Two dice are thrown simultaneously. The probability of getting a						

sum of 9 is:

(A) 1/10	(B) 3/10	(C) 1/9	(D) 4	/9		
14. 100 cards are numbered from 1 to 100. Find the probability of getting a prime number.						
(A) 3/4	(B) 27/50	(C) 1/4	(D)	29/100		
_	a blue ball is d			If the probability en the number of		
(A) 5	(B) 10	(C) 15	(D) 20			
taken out at non-defecti	t random from ve bulb is:	ntains 12 defection this box. Then (C)	the probab			
17. Cards marked with numbers 2 to 101 are placed in a box and mixed thoroughly. One card is drawn from this box randomly, then the probability that the number on card is a perfect square. (A) 9/100 (B) 1/10 (C) 3/10 (D) 19/100						
18. What is the probability of getting 53 Mondays in a leap year? (A) 1/7 (B) 53/366 (C) 2/7 (D) 7/366						
19. A card is drawn from a well shuffled deck of 52 cards. Find the probability of getting a king of red suit. (A) 1/26 (B) 3/26 (C) 7/52 (D) 1/13						
20. A game of chance consists of spinning an arrow which is equally likely to come to rest pointing to one of the number 1,2,312, then the probability that it will point to an odd number is: (A) 1/6 (B) 1/12 (C) 7/12 (D) 5/12						
21. A game consists of tossing a one rupee coin 3 times and noting its outcome each time. Aryan wins if all the tosses give the same result i.e. three heads or three tails and loses otherwise. Then the probability that Aryan will lose the game. (A) 3/4 (B) 1/2 (C) 1 (D) 1/4						

22. Riya and Kajal are friends. Probability that both will have the same birthday is the same birthday is: (A) 364/365 (B) 31/365 (C) 1/365 (D) 1/133225					
(A) 304/30	(b) 31/30	(0) 1/300	(D) 1/ 100220		
2. Then th	nber <i>x</i> is chosen at e probability that <i>x</i> (B) 2/5 (C) 3/5	² < 2 is?	numbers -2, -1, 0 , 1,		
a marble is red is 2/3,		from the jar, the p of white marbles in	nd others are white. If probability that it is the jar is:		
Then the p	ber is selected at a probability that it is (B) 4/25 (C) 1/	a multiple of 3 an	50 natural numbers. d 4 is:		
26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?					
a) $\frac{1}{7}$	b) $\frac{5}{42}$	c) $\frac{1}{21}$	d) 4/21		
	scored by batsmar . The standard dev	_	ches are 50, 70, 82,		
	b) 25.49		d) 25.69		
28. Find median and mode of the messages received on 9 consecutive days 15, 11, 9, 5, 18, 4, 18, 13, 17.					
a) 13, 15		c) 18, 15	d) 13, 16		
29. A coin is tossed up 4 times. The probability that tails turn up in 3 cases is					
a) $^{1}/_{2}$	b) $^{1}/_{3}$	c) $^{1}/_{4}$	d) $^{1}/_{6}$		
30. X is a v a) 8	variate between 0 a b) 7		f E(X²) is d) 9		
31. The random variables X and Y have variances 0.2 and 0.5 respectively. Let Z= 5X-2Y. The variance of Z is?					

probability?	•	•	one is not poss	ible in		
c) $P(x) = 0.5$	d) P(x	(x) = -0.5				
33.If E(x) =	2 and E(z) = 4 b) 6	c) 0	•	ufficient data		
34.The cov	ariance of two	independer	nt random variab	le is		
a) 1	b) 0	c) – 1	d) Un	defined		
35.If Σ P(x) a) 0) = k² – 8 then, b) 1	the value o		sufficient data		
36.If P(x) = a) 1	0.5 and x = 4, b) 0.5	then E(x) = (c) 4	? d) 2			
is always?	erete probabilit b) Infinite		on, the sum of all	probabilities		
38.If the pr	·	tting the tar	get is 0.4, find m	ean and		
39.If the probability that a bomb dropped from a place will strike the target is 60% and if 10 bombs are dropped, find mean and variance? a) 0.6, 0.24 b) 6, 2.4 c) 0.4, 0.16 d) 4, 1.6						
a) 2	e mean of toss b) 4 s the mean and	c) 8	d) 1 or standard norm	al distribution?		

c) 5

d) 7

a) 3

b) 4

						id variance and varian	
		e of a rand b) E(X			•	•	d) (E(X))2
43.I a) E(Mean of (X)	a random b) E(X2	variable 2)	e X is giv c) E(X2)	ven by) - (E(X))2	d) (E(X))2
	lean of	a constan b) a		c) a/2		d) 1	
		of a cons b) a				d) 1	
46.F	ind the	mean and	variance	e of X?			
	Х	0	1	2	3	4	
	f(x)	1/9	2/9	3/9	2/9	1/9	
a) 2,	4/3	b) 3			c) 2, 2/3	,	d) 3, 2/3

47. Find the expectation of a random variable X?

	Х	0	1	2	3	
	f(x)	1/6	2/6	2/6	1/6	
a) ().5		b) 1.5		c) 2.5	d) 3.5

48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by

c) np2q

d) npq2

- 49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.**
- a) P(X = x) = nCx px qx
- b) P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx
- 50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?**

- a) \sqrt{np} b) \sqrt{pq} c) (np)2