

SOCIOMAP: Mapping sociogenomic inequities in chronic disease risk— case study on a Nigeria cohort

Authors:

Ariel Ghislain Kemogne Kamdoun^{1#}, Blessing Itoro Afolayan², Sewade Olaolu Ogun³

¹Department of Mathematics and Statistics, University of Calgary Alberta, Canada

²DeustoTech, Faculty of Engineering, University of Deusto, Bilbao, Spain

³GetVocal AI, Paris

[#]Corresponding Author (arielghislain.kemogn@ucalgary.ca)

Abstract

Chronic disease risk is shaped by complex intersections of sociodemographic, cultural, and lifestyle factors, yet most data-driven risk models do not explicitly characterize how disease burden is structured across these dimensions in an interpretable and equity-aware manner. In settings with high sociocultural heterogeneity, this limitation obscures subgroup-specific risk patterns and constrains translational impact. Here, we introduce SOCIOMAP, an interpretable, end-to-end machine learning framework designed to quantify and explain how chronic disease risk varies across sociodemographic strata in a nationally representative Nigerian cohort of approximately 45,000 individuals. We harmonize ICD-10 diagnoses, lifestyle variables, and sociodemographic features to derive binary disease phenotypes and first map stratified prevalence and multimorbidity patterns. We then assess group-wise heterogeneity across tribe, religion, income, and education using chi-squared association analyses. To move beyond association, we train gradient-boosted tree models for disease risk prediction and apply SHAP to identify transparent, subgroup-aware drivers of risk, highlighting contributions from adiposity, socioeconomic status, cultural affiliation proxies, and diet-related variables. Unsupervised embedding and clustering further reveal latent population subgroups enriched for distinct sociodemographic and lifestyle profiles, consistent with non-uniform risk architectures across the cohort. Finally, we operationalize SOCIOMAP through interactive Streamlit tool to support exploratory analysis, risk scoring, and explanation. Together, these results demonstrate that chronic disease risk in Nigeria is systematically structured by intersecting social and lifestyle dimensions and establish SOCIOMAP as a scalable, interpretable framework for equity-centered precision public health and future integration with polygenic and gene–environment interaction analyses.

Introduction

Understanding how genetic susceptibility intersects with social and environmental context is central to advancing equitable healthcare. Over the past decade, large-scale genomic studies have uncovered thousands of loci associated with complex diseases, and polygenic risk scores (PRS) have emerged as a promising tool for individual-level risk stratification [1,2]. However, the clinical and public health utility of these approaches remains uneven. In practice, many genome-informed risk models prioritize genetic effects while treating social, cultural, and environmental factors as secondary adjustments, resulting in limited generalizability and uneven predictive performance across populations [1,2,3,6].

Africa harbours the greatest human genetic diversity globally [3], yet remains profoundly underrepresented in genomic and health data science research [13,14]. Nigeria, the most populous country in Africa, with over 200 million inhabitants and more than 250 ethnolinguistic groups, presents a uniquely informative setting in which to examine how genetic, sociocultural, and environmental factors jointly shape chronic disease risk. At the same time, Nigeria is undergoing a rapid epidemiological transition, with a growing burden of non-communicable diseases (NCDs) such as hypertension, diabetes, and cardiovascular disease [4]. Despite this shift, there is a critical lack of scalable, interpretable analytical frameworks capable of characterizing disease burden in a manner that is population-aware, equity-centered, and actionable for public health decision-making [15,16].

Importantly, health outcomes are not determined by genetic predisposition alone. A substantial body of evidence highlights the role of social determinants, including ethnicity, religion, income, education, dietary practices, and access to healthcare, in influencing disease onset, progression, and outcomes [5,6,46,47]. In multi-ethnic and socioeconomically stratified societies such as Nigeria, these factors are deeply embedded in cultural norms and lived experience, and may mediate, modify, or even dominate biological risk pathways [23-25]. Yet systematic approaches for integrating sociodemographic structure directly into disease risk modeling, particularly in African populations, remain limited, and such variables are often treated as confounders rather than as primary explanatory dimensions [19,27,28].

Here, we present an interpretable, AI-driven framework for the discovery of polygenic-sociodemographic interactions underlying chronic disease risk in Nigeria. Using a nationally representative cohort of approximately 45,000 individuals, we integrate structured phenotypic data, ICD-10-coded diagnoses, lifestyle indicators, and sociodemographic variables into a unified analytical pipeline grounded in principles of explainable artificial intelligence (XAI) and precision public health [15,16,37,38]. Rather than prioritizing predictive performance alone, our framework is explicitly designed to generate transparent, subgroup-aware insights that can inform policy, prevention strategies, and future genomic integration.

Our approach comprises four core components. First, we develop a robust preprocessing and feature engineering pipeline that harmonizes raw clinical and survey data, including the transformation of free-text and coded ICD-10 diagnoses into interpretable binary disease phenotypes [21,22]. Second, we perform comprehensive epidemiologic profiling of disease prevalence and multimorbidity, enabling fine-grained characterization of chronic disease burden across Nigeria's sociocultural landscape [45,46]. Third, we quantify stratified disease heterogeneity across tribe, religion, income, education, and lifestyle factors using chi-squared association testing [40,41], nonlinear embedding (UMAP) [10,11], and unsupervised clustering (KMeans), revealing latent population subgroups with distinct risk architectures. Fourth, we train supervised gradient-boosted tree models (XGBoost) [8] and apply SHAP-based explanations [9,12] to identify interpretable drivers of disease risk at both the population and subgroup levels, enabling systematic risk auditing and alignment with established epidemiological patterns.




Figure 1 | End-to-end interpretable AI protocol for discovering polygenic sociodemographic interactions driving chronic disease inequities in Nigeria. Using a nationally representative Nigerian cohort (~45,000 participants), the pipeline (i) harmonizes phenotype, ICD-10 diagnoses, lifestyle, and sociodemographic variables (tribe, religion, income, education) and converts raw ICD-10/free-text entries into structured binary disease flags; (ii) performs exploratory and stratified epidemiologic profiling via prevalence mapping, comorbidity/co-occurrence heatmaps, and group-wise association testing (χ^2) to quantify disparities across social strata; (iii) trains supervised disease-risk models (XGBoost) and applies SHAP to generate auditable, subgroup-aware feature attributions that highlight modifiable and structural drivers of risk; (iv) identifies latent risk subgroups through unsupervised representation learning (UMAP) and clustering (KMeans) to reveal culturally and lifestyle-aligned health

profiles; and (v) deploys results through interactive Streamlit dashboard enabling real-time risk scoring, explanation, subgroup comparison, and policy-ready reporting. The protocol is designed to be equity-centered, open-source, and extensible for future integration of genotype data, polygenic scores, and gene–environment interaction analyses.

Unlike traditional genomic studies that rely primarily on genotype data, this work focuses on the sociogenomic layer of disease risk, establishing a necessary foundation for the equitable deployment of polygenic models in African settings [19,27]. To facilitate translation, we operationalize the framework through open-source, interactive dashboards built with Dash and Streamlit, enabling real-time risk scoring, subgroup comparison, and explainable model interrogation by researchers, clinicians, and policymakers [30-32].

By embedding local sociocultural context directly into the modeling pipeline, this study contributes to a new generation of equity-centered AI frameworks for health research [16,17,33-35]. More broadly, it provides a scalable blueprint for other Global South contexts facing similar challenges of health disparity, data marginalization, and limited access to interpretable analytic tools (Fig. 1).

Results

We analyzed a nationally representative cohort of 44,927 individuals (see data description section) with harmonized ICD-10 diagnoses, lifestyle indicators, and sociodemographic variables to characterize the distribution and structure of chronic disease risk across Nigeria's diverse population (Fig. 2). Binary disease phenotypes revealed substantial heterogeneity in prevalence, comorbidity, and sociodemographic stratification, indicating that chronic disease burden is non-uniformly distributed across intersecting social dimensions.

Disease prevalence and co-occurrence exhibit structured heterogeneity

Across the cohort, hypertension was the most prevalent chronic condition (9.3%), followed by chronic hepatitis (6.4%) and type 2 diabetes (5.1%) (Fig. 3a; Supplementary Appendix Table S1). Other conditions, including hypertensive heart disease, asthma, epilepsy, and type 1 diabetes, were less frequent but exhibited pronounced variability across population subgroups.

Pairwise disease co-occurrence analysis revealed non-random clustering, particularly among cardiometabolic conditions and between metabolic and liver-related disorders (Fig. 2b; Supplementary Appendix Fig. S4), suggesting shared or interacting risk architectures rather than independent disease processes.

Sociodemographic stratification reveals pronounced disparities

Stratification of disease prevalence by sociodemographic variables revealed marked disparities across tribe, religion, income, and education (Fig. 3b; Fig. 4d). Tribal heterogeneity was especially pronounced. Chronic hepatitis prevalence varied significantly across ethnic groups, with several tribes exhibiting disproportionately elevated burden (Fig. 4d; Supplementary Appendix Table S2).




Figure 2 | Results overview of the sociogenomic analysis of chronic disease risk in Nigeria.

This multi-panel figure summarizes the key empirical findings of the proposed framework across modeling, stratification, explainability, and deployment. **(1)** An end-to-end schematic illustrates the analytical pipeline, from ICD-10 diagnoses, lifestyle, and sociodemographic inputs through preprocessing, exploratory analysis, supervised modeling, unsupervised clustering, and deployment for public health use. **(2)** Disease-level summaries include overall case counts and pairwise comorbidity structure, visualized via prevalence bar charts and a Jaccard similarity heatmap of disease co-occurrence. **(3)** Stratified prevalence and statistical significance heatmaps demonstrate that multiple chronic diseases exhibit significant heterogeneity across tribe, religion, income, and education groups. **(4)** Supervised XGBoost models combined with SHAP reveal interpretable, disease-specific drivers of risk for hypertension and type 2 diabetes at the global, domain, and individual levels, highlighting contributions from BMI, diet, religion, tribe, and socioeconomic status. **(5)** Distributions of key sociodemographic and dietary variables contextualize the population structure underlying observed disparities. **(6)** Snapshots of interactive Dash and Streamlit dashboards illustrate real-time disease risk scoring, subgroup comparison, and explainability for end users. **(7-8)** Additional SHAP-based summaries emphasize subgroup- and individual-level heterogeneity in type 2 diabetes risk. **(12-15)**

Heatmaps of association strength across education, income, religion, and tribe further confirm that chronic disease burden in Nigeria is structured along intersecting social dimensions. Collectively, these results demonstrate that chronic disease risk is non-uniformly distributed across Nigerian populations and can be transparently characterized using interpretable, equity-focused AI.

Type 1 diabetes displayed extreme concentration within a small number of tribes, most notably the Menyang (Meryan) group, where prevalence approached 20%, despite near-zero prevalence in most other groups (Supplementary Appendix Fig. S1d). These contrasts were not mirrored uniformly across religion, income, or education, suggesting localized or group-specific risk factors.

Religious affiliation was associated with differential disease burden for select outcomes, most notably type 1 diabetes (Fig. 3b). Income stratification revealed consistent socioeconomic gradients for hypertension, hypertensive heart disease, epilepsy, and chronic hepatitis, while type 2 diabetes exhibited a non-linear pattern with elevated prevalence in middle-income categories (Fig. 4d). Education level also modulated disease burden, likely reflecting differential access to diagnosis and healthcare utilization (Supplementary Appendix Fig. S5).




Figure 3 | Key observations of the AI-driven sociogenomic analysis of chronic disease risk in Nigeria. This figure summarizes the principal empirical findings from the Nigerian cohort. **(A)** Overall prevalence of major chronic and infectious diseases derived from ICD-10 binary flags, highlighting a high burden of hypertension, malaria, chronic hepatitis, and type 2 diabetes in the population. **(B)** Heatmap of disease prevalence stratified by sociodemographic factors

(tribe, religion, income, and education), demonstrating substantial heterogeneity in disease burden across social groups. **(C)** SHAP-based feature importance from XGBoost models, identifying key drivers of disease risk—including BMI, tribe, religion, diet, income, and lifestyle factors—while enabling interpretable assessment of both biological and social determinants. **(D)** UMAP visualization of unsupervised clustering reveals latent risk subgroups enriched for specific sociodemographic profiles, distinguishing high-risk clusters associated with lower income and certain tribes from lower-risk clusters characterized by higher education and different ethnic backgrounds. Together, these results show that chronic disease risk in Nigeria is structured along intersecting sociocultural and lifestyle dimensions and can be transparently characterized using interpretable AI methods.

Statistical testing confirms non-random stratification

Formal χ^2 association testing confirmed that many stratified patterns were unlikely to arise by chance. Highly significant associations were observed between tribe and multiple diseases, including hypertension, hypertensive heart disease, type 2 diabetes, asthma, and chronic hepatitis (Fig. 4b; Supplementary Appendix Table S2). Income and education were also significantly associated with cardiometabolic disease prevalence.

In contrast, acute myocardial infarction and several cancers showed limited stratification, consistent with their low prevalence in the dataset (Supplementary Appendix Table S2). Simulation-based null experiments demonstrated appropriate Type I error control under high-cardinality stratification following Benjamini–Hochberg correction (Supplementary Appendix Fig. S6).

Predictive modeling and explainability identify subgroup-specific risk drivers

Supervised gradient-boosted tree models achieved moderate discrimination for common diseases such as hypertension and type 2 diabetes, with ROC-AUC values exceeding 0.7 in some cases (Fig. 4a; Supplementary Appendix Table S1). Predictive performance was reduced for rarer diseases, reflecting class imbalance rather than model inadequacy.

SHAP-based explainability revealed consistent and interpretable drivers of disease risk. Body mass index dominated cardiometabolic outcomes, while sociodemographic variables, including religion, income, and tribe, contributed substantially to predictions, often exceeding the influence of individual behavioral variables (Fig. 3c; Fig. 4c; Supplementary Appendix Table S2). Dietary factors, particularly cooking oil type, were strongly associated with hypertension, diabetes, and liver disease risk.

Aggregation of SHAP values by sociodemographic groups revealed marked heterogeneity in feature importance across subpopulations, indicating that similar disease labels arise from distinct underlying risk architectures (Supplementary Appendix Table S2).

Unsupervised analysis reveals latent population risk structure

Unsupervised representation learning using UMAP followed by KMeans clustering identified five latent population clusters characterized by distinct combinations of sociodemographic, lifestyle, and disease features (Fig. 3d; Fig. 4e). One cluster was enriched for lower-income northern tribes with elevated cardiometabolic risk, while another comprised predominantly urban southern populations with higher BMI and increased type 2 diabetes prevalence.

Cluster recovery and robustness were validated across heterogeneous and adversarial simulation scenarios, including null, interaction-driven, access-biased, and distribution-shift regimes (Supplementary Appendix Figs. S1-S3). Disease co-occurrence structure and prevalence estimates remained stable across simulations (Supplementary Appendix Figs. S4-S5).




Figure 4 | Comprehensive results of the sociogenomic analysis of chronic disease risk in Nigeria. This multi-panel figure summarizes predictive performance, statistical associations, explainability, and subgroup discovery across chronic disease outcomes. **(1)** Predictive performance of XGBoost models across ICD-10 disease flags, reported as AUC, showing stronger discrimination for common conditions (e.g., hypertension, type 2 diabetes) and reduced performance for rarer diseases due to class imbalance. **(2)** Heatmap of statistical significance ($-\log_{10}$ p-values) from chi-squared tests assessing differences in disease prevalence across sociodemographic groups (tribe, religion, income, education), highlighting widespread and disease-specific health disparities. **(3)** SHAP-based feature importance for disease risk prediction, decomposed by domain (health, sociodemographic, and dietary factors), illustrating the dominant roles of BMI, income, tribe, religion, and cooking oil type in shaping predicted risk. **(4)** Stratified prevalence of major diseases by sociodemographic factors, showing higher prevalence in certain groups (e.g., hypertension in Tribe A, T2D in low income, chronic hepatitis in Islam). **(5)** Distribution of key sociodemographic and dietary variables, showing the prevalence of different tribes, religions, and cooking oil types. **(6)** Sociodemographic distribution of chronic hepatitis prevalence, showing higher prevalence in certain groups (e.g., Tribe A, Islam, low income, no formal education). **(7)** UMAP + KMeans clustering of individuals by sociodemographic and disease risk, identifying three distinct clusters: Cluster 1 (High Risk - Tribe A, Low Income), Cluster 2 (Urban & Mixed), and Cluster 3 (Low Risk - High Education).

for hypertension and type 2 diabetes. **(4)** Stratified prevalence plots for major diseases demonstrate pronounced gradients across tribe, religion, and income strata, confirming non-uniform disease burden. **(5)** Distribution of key sociodemographic and dietary variables provides population context for downstream analyses. **(6)** Focused stratification of chronic hepatitis prevalence across tribe, religion, income, and education reveals distinct sociocultural patterns. **(8)** UMAP + KMeans clustering identifies latent population subgroups with distinct disease-risk profiles, separating high-risk, low-income clusters from lower-risk, higher-education clusters. Collectively, these results show that chronic disease risk in Nigeria is strongly structured by intersecting social, lifestyle, and health factors and can be transparently characterized using interpretable, equity-focused AI.

Interactive deployment supports translation

All analyses were operationalized through interactive Streamlit dashboard enabling real-time exploration of disease prevalence, risk prediction, explainability, and subgroup structure (Fig. 2f; Fig. 4f). Dashboard outputs were consistent with offline analyses, demonstrating the feasibility of deploying interpretable, equity-centered AI tools for public health use in low-resource settings.

Discussion

This study establishes an interpretable, equity-centered framework for characterizing chronic disease risk in a highly diverse and historically underrepresented population. By integrating structured phenotypic data with sociodemographic context and explainable machine learning, we demonstrate that chronic disease burden in Nigeria is not uniformly distributed, but instead exhibits pronounced stratification across intersecting social, cultural, and lifestyle dimensions. These findings challenge implicit assumptions in population health modeling that disease risk can be adequately summarized without explicit consideration of local social structure and lived context.

A central insight of this work is that non-genetic factors, such as income, education, religion, and dietary practices, are not merely nuisance covariates or confounders, but primary organizing axes of disease risk. Through stratified prevalence analyses, formal χ^2 association testing, and SHAP-based attribution, we show that these dimensions consistently shape both observed disease prevalence and model-predicted risk across multiple chronic conditions. In particular, cardiometabolic diseases such as hypertension and type 2 diabetes display strong heterogeneity across tribal and socioeconomic strata, reinforcing evidence that social determinants play a fundamental role in shaping health outcomes [5,6,46,47]. These results underscore the limitations of population-agnostic risk models and highlight the need for population-aware analytical frameworks in precision public health [15,16].

The use of explainable machine learning enables direct inspection of how social and lifestyle features contribute to disease risk at both population and subgroup levels. Rather than treating machine learning models as opaque predictors, SHAP-based explanations reveal interpretable and policy-relevant drivers, including adiposity, cooking oil type, and income tier, that align with established epidemiological mechanisms while also exposing context-specific patterns [9,12,37,38]. Importantly, group-wise aggregation of SHAP values allows identification of features that disproportionately influence predictions within particular social strata, providing a quantitative framework for auditing equity and informing targeted interventions [16,33-35].

Beyond supervised risk prediction, unsupervised embedding and clustering uncover latent population subgroups with distinct risk architectures that are not fully captured by individual sociodemographic variables alone. These data-driven clusters reflect co-occurring lifestyle behaviors, socioeconomic conditions, and disease profiles, offering a complementary lens for understanding health inequities beyond predefined administrative categories. Such subgroup discovery may be especially valuable in settings where formal classifications fail to capture lived experience or where public health interventions must address overlapping and interacting risk domains.

This work deliberately focuses on the sociogenomic layer of disease risk rather than genotype data alone. While polygenic risk scores have shown promise in some contexts, their uneven portability and performance across populations remain a major barrier to equitable clinical translation [1,19,27,28]. By establishing a rigorous and interpretable framework for modeling social and environmental structure, this study lays essential groundwork for future integration of genetic data, gene–environment interaction analyses, and ancestry-aware polygenic models [13,14]. In this sense, sociogenomic context is not ancillary but foundational to equitable genomic medicine.

Several limitations warrant consideration. First, disease phenotypes were derived from ICD-10 codes and self-reported diagnoses, which may introduce misclassification or underdiagnosis, particularly in low-resource settings [21]. Second, the cross-sectional nature of the data limits causal interpretation, and observed associations should be viewed as descriptive rather than mechanistic [22-25]. Third, although the cohort is nationally representative of Nigeria, findings may not generalize to other African countries with distinct sociopolitical and healthcare contexts. Nevertheless, the analytical framework itself is portable and can be readily adapted to other populations facing similar challenges of data sparsity, heterogeneity, and health inequity. In summary, this study demonstrates how interpretable AI can be used to expose latent structure in chronic disease risk across complex social landscapes. By centering equity, transparency, and population specificity, SOCIOMAP provides a scalable blueprint for precision public health in Nigeria and other Global South settings. More broadly, this work illustrates how contextualized AI can move beyond prediction toward explanation, accountability, and action in global health research.

Methods

Real Data Analysis

Overview of the analytical framework

We developed an integrated analytical framework to identify and interpret polygenic-sociodemographic interactions shaping chronic disease risk in Nigerian populations. The framework combines epidemiological characterization, statistical association testing, supervised and unsupervised machine learning, and model explainability within a unified, equity-focused pipeline. Throughout the analysis, we emphasize interpretability, population heterogeneity, and transparency, reflecting the unique demographic diversity of Nigeria and the need for responsible, explainable AI in public health research.

Data sources and cohort characterization

The analysis was conducted on a nationally representative cohort of approximately 45,000 individuals from Nigeria. Each individual record contains detailed clinical, sociodemographic, lifestyle, behavioral, and anthropometric information. Clinical diagnoses were recorded using both raw and coded ICD-10 entries, while sociodemographic variables included ethnicity (tribe), religion, income tier, and educational attainment. Lifestyle and behavioral indicators captured dietary patterns, physical activity, smoking status, and alcohol use, alongside anthropometric measurements such as body mass index, height, and weight.

Nigeria's population structure, encompassing more than 250 ethnic groups, provides a uniquely rich setting for studying population-specific disease patterns and health inequities. This diversity motivated an analytical strategy explicitly designed to preserve subgroup structure rather than averaging effects across heterogeneous populations.

Construction of disease phenotypes from ICD-10 codes

To enable disease-specific modeling, raw ICD-10 diagnosis strings were transformed into structured binary disease indicators. A curated dictionary mapped ICD-10 root codes to clinically interpretable disease categories, such as hypertension and type 2 diabetes. Diagnosis strings were parsed using prefix matching for ICD-10 codes, supplemented by free-text matching and regular expression-based tokenization to accommodate multiple diagnoses recorded per individual.

Formally, for each individual, diagnosis strings were mapped to disease indicators using ICD-10 prefixes, producing a binary outcome matrix in which each column represents a disease phenotype and each row corresponds to an individual. This representation allowed consistent downstream modeling of multiple chronic disease outcomes within a unified framework.

Mathematically, let d_i denote the diagnosis code string for individual $i \in \{1, \dots, N\}$, where N is the total number of individuals in the cohort. Let $C = \{c_1, c_2, \dots, c_D\}$ be the set of ICD-10 code prefixes corresponding to D disease conditions of interest. We define a binary indicator (or flag) for each individual–disease pair as:

$$flag_{i,c} = 1[\text{startswith}(d_i, c)]$$

Where: $1[\cdot]$ is the indicator function, equal to 1 if the condition inside holds and 0 otherwise; $\text{startswith}(d_i, c)$ returns true if the string d_i begins with the ICD-10 prefix c .

All such flags are assembled into a binary label matrix: $Y \in \{0, 1\}^{N \times D}$; where $Y_{i,j} = flag_{i,c_j}$ indicates whether individual i exhibits a diagnosis corresponding to ICD-10 prefix c_j , and each column corresponds to one of the D disease outcomes under investigation.

Exploratory epidemiological analysis

We first conducted exploratory analyses to characterize disease burden and comorbidity patterns across the cohort. Disease prevalence was estimated empirically for each outcome as the proportion of individuals exhibiting the corresponding diagnosis. To examine multimorbidity, we constructed a disease co-occurrence matrix by computing pairwise overlaps among binary disease indicators, yielding a symmetric representation of shared disease burden.

To investigate potential health disparities, disease prevalence was further stratified by key sociodemographic variables, including ethnicity, religion, and income tier. Subgroup-specific prevalence estimates enabled an initial assessment of social gradients in disease burden and guided subsequent statistical and modeling analyses. These summaries were visualized using prevalence plots and heatmaps to support interpretability and hypothesis generation.

For each disease condition $j \in \{1, \dots, D\}$, where D is the number of binary disease flags, we computed the empirical prevalence across the cohort using:

$$\hat{p}_j = \frac{1}{N} \sum_{i=1}^N Y_{i,j}$$

Where: N is the total number of individuals in the dataset; $Y_{i,j} \in \{0, 1\}$ indicates whether individual i has disease j ; $\hat{p}_j$ represents the proportion of individuals diagnosed with disease j .

To capture comorbidity patterns, we constructed the disease co-occurrence matrix:

$$M = Y^T Y$$

where $M_{j,k}$ denotes the number of individuals diagnosed with both diseases j and k . The matrix $M \in N^{D \times D}$ provides a symmetric representation of disease pairwise associations.

Finally, for each subgroup g , we computed the mean disease prevalence:

$$\hat{p}_j^{(g)} = \frac{1}{|g|} \sum_{i \in g} Y_{i,j}$$

where $|g|$ is the number of individuals in group g , and $\hat{p}_j^{(g)}$ is the subgroup-specific prevalence of disease j . This enabled initial detection of social gradients and disparities in health outcomes.

Statistical testing of sociodemographic associations

To formally assess whether observed differences in disease prevalence across sociodemographic groups reflected statistically meaningful associations, we conducted chi-squared tests of independence between group membership and disease status.

Let $G \in \{1, \dots, K\}$ represent the levels of a discrete socio demographic grouping variable (e.g., the observed tribes), and let $Y \in \{0,1\}$ be a binary indicator of disease status for a particular condition.

For each disease–group combination, we constructed a $2 \times K$ contingency table. The observed cell counts were defined as:

$$T_{k,y} = \sum_{i=1}^N 1[g_i=k \wedge y_i=y]$$

Where: g_i is the group assignment of individual i ; y_i is the disease flag (1 = diagnosed, 0 = not diagnosed); $T_{k,y}$ is the number of individuals in group k with disease status y ; $1[\cdot]$ denotes the indicator function. We then computed the chi-squared test statistic:

$$\chi^2 = \sum_{ky} \frac{(T_{k,y} - E_{k,y})^2}{E_{k,y}}$$

where $E_{k,y}$ represents the expected frequency under the null hypothesis of independence between group membership and disease status, typically computed as:

$$E_{k,y} = \frac{T_{k,.} \cdot T_{.,y}}{T_{..}}$$

With: $T_{k,.}$ = total in group k; $T_{.,y}$ = total with disease status y; $T_{..}$ = N, the total sample size.

The p-value was obtained from the χ^2 distribution with (K - 1) degrees of freedom, enabling formal hypothesis testing.

This procedure was repeated for each disease condition across multiple sociodemographic factors. Statistically significant results (e.g., $p < 0.05$ after correction) were interpreted as evidence of health disparities potentially driven by group-specific exposures, genetics, or access to care.

For each disease and grouping variable, contingency tables were constructed to compare observed and expected frequencies under the null hypothesis of independence.

The resulting chi-squared statistics were evaluated using the appropriate degrees of freedom, and statistical significance was assessed with correction for multiple testing where applicable. Statistically significant associations were interpreted as evidence of population-level disparities that may reflect differential exposure, access to care, structural determinants, or genetic background. These tests provided a rigorous statistical foundation for the subsequent application of machine learning models.

Feature encoding and data representation

To prepare the data for predictive modeling, all sociodemographic, lifestyle, and clinical variables were encoded into a unified numerical feature matrix. Let $X \in R^{N \times P}$ denote the final feature matrix, where: N is the number of individuals in the dataset, and P is the total number of engineered features (both numeric and categorical). Categorical variables such as ethnicity, religion, and cooking oil type were transformed using one-hot encoding, while continuous variables including BMI and dietary intake measures were normalized to ensure comparable scales across features. These processed variables were then concatenated horizontally to produce the final feature matrix:

$$X = [X^{(num)} \mid X^{(cat)}]$$

Where: $X^{(num)} \in R^{N \times P_1}$ represents the block of normalized numeric features; $X^{(cat)} \in R^{N \times P_2}$ contains the one-hot encoded categorical variables; $P = P_1 + P_2$ is the total number of features after encoding. The resulting design matrix combined numeric and categorical information in a format suitable for both tree-based and nonlinear learning algorithms.

Supervised disease risk modeling

For each disease outcome, we trained a separate binary classifier using gradient-boosted decision trees implemented via XGBoost. This approach was chosen for its strong performance on heterogeneous tabular data, robustness to missing values, and built-in regularization mechanisms. Models were trained to estimate the conditional probability of disease presence given an individual's sociodemographic and lifestyle profile by minimizing binary cross-entropy loss.

Let $X \in R^{N \times P}$ be the design matrix of encoded features, and let $y \in \{0,1\}$ be the binary label vector indicating the presence or absence of a specific disease for N individuals. Each XGBoost model aims to approximate the conditional probability $y^i = P(y_i=1 | x_i)$ by minimizing the binary cross-entropy (log-loss):

$$L = -\frac{1}{N} \sum_{i=1}^N [y_i \log(\hat{y}_i) + (1-y_i) \log(1-\hat{y}_i)]$$

Where: $\hat{y}_i \in (0,1)$ is the predicted probability for individual i ; $y_i \in \{0,1\}$ is the true disease label; L is the average loss across all samples. Hyperparameters controlling model complexity and learning dynamics were optimized using stratified cross-validation to mitigate overfitting and account for class imbalance. Model performance was evaluated on held-out data using receiver operating characteristic area under the curve (ROC-AUC), along with precision, recall, and F1-score, allowing disease-specific assessment of predictive performance.

Model interpretability and subgroup-level explanation

To ensure transparency and facilitate equity-oriented interpretation, we applied SHAP (SHapley Additive exPlanations) to all trained models. SHAP values quantify the contribution of each feature to an individual prediction, enabling both local and global interpretability.

Global feature importance was assessed by averaging the absolute SHAP values across all individuals, yielding a ranked summary of the most influential predictors of disease risk at the population level. To examine subgroup-specific risk drivers, SHAP values were further aggregated within sociodemographic groups such as ethnic categories, income tiers, and religious affiliations. This stratified aggregation allowed direct comparison of how feature contributions differ across population subgroups, highlighting potential sources of health inequity.

Essentially, mathematically, to assess overall feature importance across the population, we computed global SHAP values by averaging the absolute contributions of each feature over all individuals:

$$\phi_j = \frac{1}{N} \sum_{i=1}^N |\phi_{ij}|$$

Where: ϕ_{ij} denotes the SHAP value of feature j for individual i ; N is the total number of individuals; ϕ_j quantifies the average marginal contribution of feature j to the model's predicted log-odds or probability output. These global values provide a ranked summary of which features were most influential in disease risk prediction across the entire dataset.

To identify subgroup-specific risk drivers and potential sources of disparity, we stratified the SHAP values by socio demographic group g (e.g., tribe, religion, income tier). For each feature j , we computed the average SHAP value within group g as:

$$\overline{\phi}_j^{(g)} = \frac{1}{|g|} \sum_{i \in g} \phi_{ij}$$

Where: $|g|$ is the number of individuals in group g ; $\overline{\phi}_j^{(g)}$ represents the average contribution of feature j within that subgroup. This group-level aggregation enabled population-level interpretation of risk factors, revealing whether certain social groups were disproportionately influenced by specific features.

Unsupervised population stratification

To uncover latent subpopulations with distinct health profiles, we applied unsupervised learning techniques combining nonlinear dimensionality reduction and clustering. High-dimensional feature representations were embedded into a two-dimensional latent space using Uniform Manifold Approximation and Projection (UMAP), which preserves both local neighborhood relationships and global data structure.

We employed UMAP to project the high-dimensional feature matrix $X \in R^{N \times P}$ into a low-dimensional space that preserves both local neighborhoods and global data structure. Formally, the embedding is given by:

$$Z = \text{UMAP}(X), Z \in R^{N \times 2}$$

Where: N is the number of individuals, P is the number of input features, Z is the resulting 2D latent representation for visualization and clustering.

UMAP was selected due to its ability to maintain the manifold geometry of the input space, making it particularly well-suited for identifying meaningful structure in high-dimensional sociodemographic and health-related data.

We then applied KMeans clustering to the UMAP embedding Z to partition individuals into K latent clusters. For each individual i , cluster assignment was defined as:

$$C_i = \underset{k}{\operatorname{argmin}} ||Z_i - \mu_k||^2$$

Where: $C_i \in \{1, \dots, K\}$ is the assigned cluster label for individual i ; $Z_i \in R^2$ is the UMAP embedding of individual i ; μ_k is the centroid of cluster k . The number of clusters K was determined using silhouette analysis or the elbow method, depending on the specific disease context.

Clustering was performed on the UMAP embeddings using KMeans, with the number of clusters determined via standard cluster validation criteria. The resulting clusters were visualized and annotated with predicted disease risk and sociodemographic attributes, revealing emergent subgroups characterized by distinct combinations of social context and health risk.

Interactive visualization and decision-support tools

To promote accessibility, transparency, and real-world applicability, we developed interactive web-based dashboards for exploring model outputs and population-level patterns. Using the Dash framework, we implemented an interface for interactive inspection of SHAP values, subgroup comparisons, and disease-specific risk factors. A complementary Streamlit application enables real-time disease risk prediction, interactive cluster exploration, and population-level summaries, supporting scenario analysis and targeted intervention planning.

Together, this framework provides a scalable, interpretable, and equity-centered approach to modeling chronic disease risk in diverse populations. By integrating statistical testing, machine learning, explainability, and interactive visualization, the pipeline enables both rigorous scientific inference and actionable public health insights tailored to Nigerian populations.

Simulation Study

Notation and variable definitions

We index individuals by $i \in \{1, \dots, N\}$ and diseases by $d \in \{1, \dots, D\}$. Let $\sigma(\cdot)$ denote the logistic (sigmoid) function, $\sigma(x) = \{1 + \exp(-x)\}^{-1}$.

Population size and group structure

- N : number of individuals in a simulated cohort (here $N \approx 45,000$).
- $T_i \in \{1, \dots, K_T\}$: ethnolinguistic group (“tribe”) membership for individual i .
- K_T : number of tribes (high-cardinality; e.g., 80-120 in the narrative; 100 by default in our implementation).
- $R_i \in \{1, \dots, K_R\}$: religion category for individual i .
- K_R : number of religion categories (5 by default in our implementation).
- $G_i \in \{1, \dots, K_G\}$: geographic region for individual i (3 by default in our implementation; optional).
- K_G : number of regions.

Latent socioeconomic score and social determinants

- S_i : latent socioeconomic score used to induce correlation/confounding among social determinants.
- μT_i : tribe-specific shift in the mean of S_i .
- μR_i : religion-specific shift in the mean of S_i .
- μG_i : region-specific shift in the mean of S_i .
- Distributional statement: $S_i \sim N(\mu T_i + \mu R_i + \mu G_i, 1)$,
- I_i : ordinal income tier for individual i (e.g., low/mid/high; coded as integers in simulation).
- E_i : ordinal educational attainment for individual i (coded as integers in simulation).
- $U_i \in \{0,1\}$: urbanicity indicator (1 = urban, 0 = rural), generated with probability increasing in S_i .

Lifestyle/behavioral variables

- PA_i : physical activity score (continuous).
- $Smoking_i \in \{0,1\}$: smoking indicator.

Anthropometrics

- BMI_i : body mass index for individual i , simulated as a continuous trait:
$$BMI_i = 22 + 1.5 \mathbb{1}(U_i=1) + 1.0 I_i - 0.6 PA_i + bT_i + bG_i + \eta_i,$$

- $1(\cdot)$: indicator function (1 if the condition holds, else 0).
- bT_i : tribe-level random effect contributing to BMI.
- bG_i : region-level random effect contributing to BMI.
- η_i : individual-level noise term for BMI.

Latent multimorbidity driver

- H_i : shared latent health factor that induces multimorbidity:

$$H_i = 0.6 BMI_i + 0.4 I_i - 0.3 PA_i + 0.2 Smoking_i + uT_i + uG_i + \xi_i,$$
- uT_i : tribe-level random effect contributing to shared disease liability.
- uG_i : region-level random effect contributing to shared disease liability.
- ξ_i : individual-level noise term for H_i .

Disease outcomes and risk models

- $Y_{i,d}^{true} \in \{0,1\}$: latent (true) disease status for disease d in individual i.
- Disease model:

$$\Pr(Y_{i,d}^{true}=1) = \sigma(\alpha_d + \lambda_d H_i + f_d(BMI_i) + \beta_d^T X_i + INT_{i,d}),$$
- α_d : disease-specific intercept controlling baseline prevalence.
- λ_d : disease-specific loading on shared risk factor H_i .
- f_d : disease-specific nonlinear effect of BMI (e.g., hinge/threshold effect).
- X_i : vector of observed covariates (lifestyle + sociodemographics) for individuals (e.g., income, education, urbanity, diet, smoking, alcohol, physical activity).
- β_d : disease-specific coefficient vector for X_i .
- $INT_{i,d}$: interaction term(s) encoding heterogeneous “risk architectures,” such as effect modification (e.g., diet $\times$ income) or subgroup-specific amplification (e.g., BMI $\times$ selected tribes/regions).

Diagnosis access, ICD observation model, and coding noise

- $A_i \in \{0,1\}$: diagnosis/access indicator (1 = disease can be clinically recognized/recorded; 0 = underdiagnosed), modeled as:

$$\Pr(A_i=1) = \sigma(\delta_0 + 0.8 I_i + 0.7 E_i + 0.6 U_i + \delta_{G_i}),$$
- δ_0 : intercept for diagnosis access.

- δ_{G_i} : region-specific offset for diagnosis access.
- $Y_{i,d}^{obs} \in \{0,1\}$: observed ICD-based disease flag derived from $Y_{i,d}^{true}$ through access and coding noise.
- $sens_d$: ICD sensitivity for disease d (probability of recording a true case when access is present).
- $spec_d$: ICD specificity for disease d (probability of not recording disease when truly absent, given access).

Missingness

- Missingness is introduced in selected covariates (e.g., BMI_i , I_i diet variables) via probabilities that depend on socioeconomic and geographic factors; missingness is therefore non-random with respect to observed determinants (e.g., lower SES, rural residence, specific regions).

Evaluation metrics referenced

- χ^2 tests: Pearson chi-squared tests of association between subgroup membership and observed disease status (with multiple-testing correction).
- ARI: adjusted Rand index comparing discovered clusters to ground-truth clusters (when simulated).
- NMI: normalized mutual information comparing discovered clusters to ground-truth clusters (when simulated).

Rationale and objectives

To rigorously validate the SOCIOMAP framework under controlled conditions, we conducted an extensive simulation study designed to mirror the demographic, sociocultural, and epidemiological complexity of the Nigerian population (Supplementary Simulation Table S1). The primary objective was to assess whether SOCIOMAP can reliably recover known ground-truth structure across the full analytical pipeline, including stratified epidemiology, statistical association testing, supervised risk prediction, explainability, and unsupervised subgroup discovery. Unlike idealized simulations commonly used in machine-learning validation, our design explicitly incorporates realistic sources of bias and uncertainty—such as high-cardinality social groups, multimorbidity, underdiagnosis driven by healthcare access, imperfect ICD coding, and non-random missingness—to reflect the challenges encountered in real-world public health data .

The simulation was designed to evaluate four core claims: (i) accurate detection of sociodemographic heterogeneity in disease prevalence while controlling false discoveries; (ii) recovery of nonlinear and interaction-driven disease risk through supervised modeling; (iii) fidelity of SHAP-based global and subgroup-specific explanations relative to known data-generating mechanisms; and (iv) identification of latent population subgroups with distinct risk architectures using unsupervised learning, while avoiding spurious clustering when no such structure exists.

Simulated cohort and sociodemographic structure

Each simulation replicate generated a synthetic cohort of approximately $N \approx 45,000$ individuals, matching the scale of the real SOCIOMAP dataset. Individuals were assigned to a high-cardinality set of ethnolinguistic groups (“tribes”), denoted $T_i \in \{1, \dots, K_T\}$, with K_T ranging from 80 to 120. Tribe sizes were drawn from a heavy-tailed distribution to emulate the presence of a few large groups alongside many small ones, deliberately reproducing the small-cell instability that complicates stratified inference in real multi-ethnic populations (Supplementary Simulation Table S1).

To induce realistic correlation and confounding among social determinants, additional sociodemographic variables—including religion, income tier, educational attainment, urban–rural residence, and geographic region—were generated from a shared latent socioeconomic score S_i , defined as

$$S_i \sim N(\mu T_i + \mu R_i + \mu G_i, 1),$$

where μT_i and μG_i capture tribe-, religion-, and region-specific shifts. Income and education were then generated as ordinal transformations of S_i , while urbanicity followed a Bernoulli model with probability increasing in S_i . This construction ensured that no single social variable acted in isolation, reflecting the intertwined nature of cultural, economic, and geographic factors in shaping health outcomes.

Lifestyle, anthropometric, and behavioral variables

Lifestyle and behavioral variables were generated conditionally on sociodemographic context. Dietary features such as cooking oil type and fruit and vegetable intake depended on income, education, region, and cultural group, while smoking status, alcohol use, and physical activity reflected known urban–rural and socioeconomic gradients. Body mass index (BMI) was simulated as a continuous trait,

$$BMI_i = 22 + 1.5 \mathbb{1}(U_i=1) + 1.0 I_i - 0.6 PA_i + bT_i + bG_i + \eta_i,$$

where bT_i and bG_i represent small tribe- and region-level random effects, and η_i is individual-level noise. This formulation allowed BMI to function both as a global risk factor and as a context-dependent contributor to disease risk.

Disease generation and multimorbidity structure

Multiple binary disease outcomes were simulated using logistic models combining a shared latent health factor with disease-specific components. The shared latent factor,

$$H_i = 0.6 BMI_i + 0.4 I_i - 0.3 PA_i + 0.2 Smoking_i + uT_i + uG_i + \xi_i,$$

captured common cardiometabolic risk and induced realistic multimorbidity across diseases.

For each disease d , the true disease status followed

$$\Pr(Y_{i,d}^{true}=1) = \sigma(\alpha_d + \lambda_d H_i + f_d(BMI_i) + \beta_d^T X_i + INT_{i,d}),$$

where $f_d(\cdot)$ represents nonlinear BMI effects, X_i includes lifestyle and sociodemographic covariates, and $INT_{i,d}$ encodes explicit interaction structure.

Interaction terms were used to define heterogeneous “risk architectures,” including socioeconomic effect modification (e.g., dietary effects differing by income tier) and group-specific amplification of biological risk (e.g., stronger BMI effects within selected ethnolinguistic or regional subsets). These ground-truth interactions were predefined, enabling direct evaluation of whether SOCIOMAP’s stratification and explainability components could recover them. Disease-specific coefficients and ICD-10 noise parameters are reported in Supplementary Simulation Tables S2 and S2b.

Diagnosis bias, ICD noise, and missingness

To reflect real-world clinical data limitations, true disease status was not directly observed. Instead, observed ICD-10 disease flags were generated through a two-stage process. First, individuals were assigned a probability of being diagnosed,

$$\Pr(A_i=1) = \sigma(\delta_0 + 0.8 I_i + 0.7 E_i + 0.6 U_i + \delta_{G_i}),$$

modeling systematic underdiagnosis in populations with reduced healthcare access. Second, diagnosed cases were subject to imperfect ICD coding, with disease-specific sensitivity and specificity parameters governing false negatives and false positives (Supplementary Simulation Table S2).

Non-random missingness was further introduced into BMI, income, and dietary variables, with missingness probabilities depending on socioeconomic and geographic factors. This design allowed explicit evaluation of robustness to missing data and reliance on native missing-value handling in tree-based models.

Simulation scenarios, evaluation, and reproducibility

SOCIOMAP was evaluated across a grid of simulation scenarios (Supplementary Simulation Table S1), including null settings with no subgroup heterogeneity, main-effect heterogeneity, strong interaction-driven heterogeneity, access-driven diagnosis bias, high-cardinality group stress tests, rare-disease regimes, multimorbidity-dominated settings, and distribution shifts between training and test cohorts (Supplementary Simulation Table S3). Because the ground truth was known, performance was assessed at each stage of the pipeline, including prevalence recovery, power and false discovery control of chi-squared tests, predictive discrimination and calibration, explainability fidelity, and recovery of latent subgroup structure using adjusted Rand index and normalized mutual information. Rare disease specifications are detailed in Supplementary Simulation Table S4.

Each scenario was replicated multiple times with fixed random seeds. For every replicate, the full SOCIOMAP pipeline was applied without modification, mirroring the real-data analysis, and all metrics and figures were automatically saved and summarized.

Across all scenarios, the simulation study demonstrates that SOCIOMAP reliably recovers true sociodemographic heterogeneity, interaction-driven risk structures, and latent population subgroups under realistic data-generating conditions, while maintaining appropriate control of false discoveries and avoiding spurious stratification under null settings. By explicitly modeling underdiagnosis, ICD misclassification, and structured missingness, the simulations provide strong evidence that the observed real-data findings are not artifacts of data quality limitations or modeling assumptions.

In unsupervised settings, SOCIOMAP accurately recovers true latent risk subgroups across a wide range of heterogeneous and adversarial scenarios, while exhibiting near-zero false structure under null conditions (Supplementary Simulation Fig. S1). Complementarily, supervised models achieve meaningful discrimination only when genuine sociodemographic or interaction-driven risk is present, remain robust under access bias and small-cell stress, and degrade gracefully under distribution shift (Supplementary Simulation Fig. S2). Mutual information-based evaluation further confirms that unsupervised clustering captures true latent population structure while maintaining strict control against spurious stratification (Supplementary Simulation Fig. S3).

Beyond marginal effects, SOCIOMAP robustly preserves disease–disease dependency structure across heterogeneity, access bias, rare disease imbalance, and distribution shift (Supplementary

Simulation Fig. S4). Prevalence estimation similarly remains accurate across most regimes, with predictable degradation only under severe access-driven underdiagnosis and extreme population fragmentation (Supplementary Simulation Fig. S5). Collectively, these results establish SOCIOMAP as a robust and interpretable framework for equity-centered precision public health analysis in complex, underrepresented populations.




Figure 5 | Type I error control under the null scenario (S1_null).

This figure evaluates false positive rates for stratified χ^2 association testing when no true sociodemographic heterogeneity is present. Under raw (uncorrected) testing, the false positive rate is substantially inflated relative to the nominal significance level ($\alpha = 0.05$, dashed red line), reflecting the multiple-testing burden induced by high-cardinality social stratification. In contrast, Benjamini–Hochberg false discovery rate (BH-FDR) correction substantially reduces spurious discoveries, demonstrating effective control of Type I error under the null. These results confirm that SOCIOMAP’s inferential safeguards appropriately prevent false subgroup detection while preserving interpretability in high-dimensional, stratified public health settings.




Figure 6 | Detection power under heterogeneous and interaction-driven scenarios.

This figure reports statistical power for stratified χ^2 testing after false discovery rate (BH-FDR) correction across simulation scenarios with increasing degrees of sociodemographic and epidemiological heterogeneity. Power is lowest when heterogeneity is driven solely by marginal main effects (S2_main_effects), reflecting the limited detectability of weak, diffuse signals across high-cardinality groups. Power increases under explicit interaction-driven risk architectures (S3_interactions) and further improves under small-cell stress with structured heterogeneity (S5_many_small_tribes). The highest power is achieved in the strong multimorbidity regime (S7_multimorbidity_strong), where shared latent risk amplifies group-specific effects. These results demonstrate that SOCIOMAP attains meaningful sensitivity to true sociodemographic heterogeneity while preserving strict Type I error control, even under adverse population structure and disease correlation.

References

1. Martin, A. R., Kanai, M., Kamatani, Y., Okada, Y., Neale, B. M. & Daly, M. J. Clinical use of current polygenic risk scores may exacerbate health disparities. *Nat. Genet.* 51, 584–591 (2019).
2. Knowles, J. W. & Ashley, E. A. Cardiovascular disease: The rise of the genetic risk score. *PLoS Med.* 15, e1002547 (2018).

3. The 1000 Genomes Project Consortium. A global reference for human genetic variation. *Nature* 526, 68–74 (2015).
4. World Health Organization. *Noncommunicable Diseases Country Profiles 2021: Nigeria* (WHO, 2021).
5. Marmot, M. Social determinants of health inequalities. *Lancet* 365, 1099–1104 (2005).
5. Braveman, P., Cubbin, C., Egerter, S., Williams, D. R. & Pamuk, E. Socioeconomic disparities in health in the United States: What the patterns tell us. *Am. J. Public Health* 100, S186–S196 (2010).
6. Popejoy, A. B. & Fullerton, S. M. Genomics is failing on diversity. *Nature* 538, 161–164 (2016).
7. Nature Editorial. How to bring more diversity into polygenic risk scores. *Nature* 606, 9 (2022).
8. Chen, T. & Guestrin, C. XGBoost: A scalable tree boosting system. In *Proc. 22nd ACM SIGKDD Int. Conf. Knowledge Discovery and Data Mining*, 785–794 (ACM, 2016).
9. Lundberg, S. M. & Lee, S.-I. A unified approach to interpreting model predictions. *Adv. Neural Inf. Process. Syst.* 30, 4765–4774 (2017).
10. McInnes, L., Healy, J. & Melville, J. UMAP: Uniform Manifold Approximation and Projection for dimension reduction. *J. Open Source Softw.* 3, 861 (2018).
11. Becht, E., McInnes, L., Healy, J., Dutertre, C.-A., Kwok, I. W. H., Ng, L. G. & Newell, E. W. Dimensionality reduction for visualizing single-cell data using UMAP. *Nat. Biotechnol.* 37, 38–44 (2019).
12. Lundberg, S. M., Erion, G., Chen, H. *et al.* From local explanations to global understanding with explainable AI for trees. *Nat. Mach. Intell.* 2, 56–67 (2020).
13. H3Africa Consortium. Enabling the genomic revolution in Africa. *Science* 344, 1346–1348 (2014).
14. Sirugo, G., Williams, S. M. & Tishkoff, S. A. The missing diversity in human genetic studies. *Cell* 177, 26–31 (2019).
15. Khoury, M. J., Iademarco, M. F. & Riley, W. T. Precision public health for the era of precision medicine. *Am. J. Prev. Med.* 50, 398–401 (2016).

16. Rajkomar, A., Hardt, M., Howell, M. D., Corrado, G. & Chin, M. H. Ensuring fairness in machine learning to advance health equity. *Ann. Intern. Med.* 169, 866–872 (2018).
17. Obermeyer, Z., Powers, B., Vogeli, C. & Mullainathan, S. Dissecting racial bias in an algorithm used to manage the health of populations. *Science* 366, 447–453 (2019).
18. Collins, F. S. & Varmus, H. A. A new initiative on precision medicine. *N. Engl. J. Med.* **372**, 793–795 (2015).
19. Manrai, A. K. *et al.* Genetic misdiagnoses and the potential for health disparities. *N. Engl. J. Med.* **375**, 655–665 (2016).
20. Popejoy, A. B. *et al.* Diversity in precision medicine research: a first look at U.S. initiatives. *Health Aff.* **37**, 1749–1757 (2018).
21. Goldstein, B. A., Navar, A. M., Pencina, M. J. & Ioannidis, J. P. A. Opportunities and challenges in developing risk prediction models with electronic health records data. *J. Am. Med. Assoc.* **375**, 198–208 (2017).
22. Hernán, M. A. & Robins, J. M. Using big data to emulate a target trial when a randomized trial is not available. *Am. J. Epidemiol.* **183**, 758–764 (2016).
23. VanderWeele, T. J. & Robinson, W. R. On the causal interpretation of race in regressions adjusting for confounding and mediating variables. *Epidemiology* **25**, 473–484 (2014).
24. Williamson, E. J., Aitken, Z., Lawrie, J., Dharmage, S. C. & Burgess, J. A. Introduction to causal diagrams for confounder selection. *Respirology* **19**, 303–311 (2014).
25. Richiardi, L., Bellocco, R. & Zugna, D. Mediation analysis in epidemiology: methods, interpretation and bias. *Int. J. Epidemiol.* **42**, 1511–1519 (2013).
26. Khera, A. V. *et al.* Genetic risk, adherence to a healthy lifestyle, and coronary disease. *N. Engl. J. Med.* **375**, 2349–2358 (2016).
27. Duncan, L. *et al.* Analysis of polygenic risk score usage and performance in diverse human populations. *Nat. Commun.* **10**, 3328 (2019).
28. Carlson, C. S. *et al.* Generalization and dilution of association results from European GWAS in non-European populations. *PLoS Genet.* **9**, e1003607 (2013).

29. Tiffin, P. A., Shah, S., Støvring, H. & Chan, K. Machine learning in clinical medicine: a practical guide. *BMC Med.* **18**, 44 (2020).
30. Obermeyer, Z. & Emanuel, E. J. Predicting the future—big data, machine learning, and clinical medicine. *N. Engl. J. Med.* **375**, 1216–1219 (2016).
31. Riley, W. T., Glasgow, R. E., Etheredge, L. & Abernethy, A. P. Rapid, responsive, relevant (R3) research: a call for a rapid learning health research enterprise. *Clin. Transl. Sci.* **6**, 114–119 (2013).
32. Beam, A. L. & Kohane, I. S. Big data and machine learning in health care. *JAMA* **319**, 1317–1318 (2018).
33. Chouldechova, A. & Roth, A. A snapshot of the frontiers of fairness in machine learning. *Commun. ACM* **63**, 82–89 (2020).
34. Barocas, S., Hardt, M. & Narayanan, A. *Fairness and Machine Learning*. (fairmlbook.org, 2019).
35. Kleinberg, J., Mullainathan, S. & Raghavan, M. Inherent trade-offs in the fair determination of risk scores. *Proc. ITCS* **2017**, 43:1–43:23.
36. Ribeiro, M. T., Singh, S. & Guestrin, C. “Why should I trust you?” Explaining the predictions of any classifier. *Proc. KDD* **2016**, 1135–1144.
37. Molnar, C. *Interpretable Machine Learning*. (Lulu.com, 2020).
38. Doshi-Velez, F. & Kim, B. Towards a rigorous science of interpretable machine learning. *arXiv* **1702.08608** (2017).
39. Stuart, E. A. Matching methods for causal inference: a review and a look forward. *Stat. Sci.* **25**, 1–21 (2010).
40. Benjamini, Y. & Hochberg, Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J. R. Stat. Soc. B* **57**, 289–300 (1995).
41. Efron, B. Large-scale simultaneous hypothesis testing. *J. Am. Stat. Assoc.* **99**, 96–104 (2004).

42. Gelman, A., Hill, J. & Vehtari, A. Regression and other stories. (Cambridge Univ. Press, 2020).
43. Spiegelhalter, D. J. Probabilistic prediction in patient management and clinical trials. *Stat. Med.* **5**, 421–433 (1986).
44. Steyerberg, E. W. *Clinical Prediction Models*. (Springer, 2019).
45. Subramanian, S. V., Jones, K., Kaddour, A. & Krieger, N. Revisiting Robinson: the perils of individualistic and ecologic fallacy. *Int. J. Epidemiol.* **38**, 342–360 (2009).
46. Diez Roux, A. V. Investigating neighborhood and area effects on health. *Am. J. Public Health* **91**, 1783–1789 (2001).
47. Glass, T. A. & McAtee, M. J. Behavioral science at the crossroads in public health. *Soc. Sci. Med.* **62**, 1650–1671 (2006).
48. Boulle, A. *et al.* Data centre profile: the provincial health data centre of the Western Cape Province, South Africa. *Int. J. Popul. Data Sci.* **4**, 1143 (2019).
49. Kahn, R. *et al.* Precision public health and structural racism in the United States. *Lancet Digit. Health* **4**, e697–e706 (2022).
50. Dowd, J. B. & Hamoudi, A. Is life expectancy really falling for groups of low socioeconomic status? *Proc. Natl Acad. Sci. USA* **111**, 193–198 (2014).

Data Description

The analysis is based on a nationally representative cohort of 44,927 individuals from Nigeria, comprising harmonized sociodemographic, lifestyle, anthropometric, and health-related variables. The dataset integrates population survey data with clinical information and was curated to enable stratified epidemiologic analysis and interpretable machine learning at scale. Sociodemographic variables include age, sex, ethnolinguistic affiliation (tribe), religion, educational attainment, household income tier, region, and urban–rural residence. Lifestyle and behavioral factors encompass dietary practices (including primary cooking oil type and fruit and vegetable intake), physical activity indicators, smoking status, and alcohol consumption. Anthropometric measures include body mass index (BMI), derived from recorded height and weight where available. Health outcomes were constructed from structured clinical records and free-text entries mapped to ICD-10–coded diagnoses. These diagnoses were subsequently harmonized and transformed into binary disease phenotypes representing chronic and non-communicable conditions, including hypertension, diabetes (type 1 and type 2), cardiovascular and hypertensive heart disease, asthma, epilepsy, liver disease, and related outcomes. Multimorbidity patterns were retained to preserve disease co-occurrence structure. All variables underwent a standardized preprocessing and quality-control

pipeline, including normalization of categorical encodings, auditing and handling of missingness, and clinically informed phenotype construction. The data contain no direct personal identifiers, and all analyses were conducted on de-identified records in accordance with applicable ethical, legal, and data governance requirements.

Data and Code Availability

The data analysed in this study were obtained under a restricted-use agreement and contain sensitive individual-level health and sociodemographic information. As a result, the raw data cannot be made publicly available due to ethical, legal, and privacy constraints. Access to the data may be considered on a case-by-case basis by the data custodians, subject to appropriate institutional approvals, data use agreements, and compliance with relevant ethical and data protection regulations. All code used for data preprocessing, analysis, statistical modelling, explainability, and visualization is publicly available at <https://github.com/1234-Ariel-code/sociomap/>. To support transparency, reproducibility, and public engagement without releasing protected individual-level records, we additionally provide a fully open, interactive demonstration of the SOCIOMAP framework based on a simulated Nigerian-like cohort. The synthetic data reproduce key structural properties of the real dataset, including high-cardinality sociodemographic groups, multimorbidity, access-driven underdiagnosis, ICD misclassification, structured missingness, and interaction-driven risk architectures.

This allows safe public exploration of the full analytical pipeline and validation of methodological behavior under realistic conditions. The simulation code and demo materials are available at <https://github.com/1234-Ariel-code/sociomap/tree/main/sociomap-demo>, and an interactive public deployment can be accessed at <https://sociomap-owjbfna3cmr7zuvzs2euq.streamlit.app/>. All analyses were conducted on de-identified data in accordance with applicable institutional policies and ethical and data protection standards.

Acknowledgement

We are grateful to the participants and communities whose data made this research possible. We also acknowledge the broader public health, open science, and machine learning communities for developing the tools, frameworks, and resources that informed our methodology.

Appendix & Supplementary

All supplementary figures and tables are provided in a separate Supplementary Information file. This includes simulation figures (Figs. S1-S7) and tables (Tables S1-S4), which detail the simulation data-generating processes, global configuration parameters, disease-specific ground-truth specifications, ICD noise and access-bias models, and the full grid of simulation scenarios used to validate the SOCIOMAP framework.