

Relations

Chapter 9

Chapter Summary

- Relations and Their Properties
- n -ary Relations and Their Applications (*not currently included in overheads*)
- Representing Relations
- Closures of Relations
- Equivalence Relations
- Partial Orderings

Relations and Their Properties


Section 9.1

Binary Relations (二元關係)

Definition: A *binary relation* R from a set A to a set B is a subset $R \subseteq A \times B$.

Example:

- Let $A = \{0, 1, 2\}$ and $B = \{a, b\}$
- $\{(0, a), (0, b), (1, a), (2, b)\}$ is a relation from A to B .
- We can represent relations from a set A to a set B graphically or using a table:



R	a	b
0	×	×
1	×	
2		×

Relations are more general than functions. A function is a relation where exactly one element of B is related to each element of A .

Binary Relation on a Set

Definition: A binary relation R *on a set A* is a subset of $A \times A$ or a relation from A to A .

Example:

- Suppose that $A = \{a, b, c\}$. Then $R = \{(a, a), (a, b), (a, c)\}$ is a relation on A .
- Let $A = \{1, 2, 3, 4\}$. The ordered pairs in the relation $R = \{(a, b) \mid a \text{ divides } b\}$ are
 $R = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}$.

Binary Relation on a Set (*cont.*)

Question: How many relations are there on a set with n elements?

Solution:

Because a relation on A is a subset of $A \times A$, we count the subsets of $A \times A$.

Since $A \times A$ has n^2 elements when A has n elements, and a set with m elements has 2^m subsets, there are $2^{|A|^2}$ subsets of $A \times A$. Therefore, there are $2^{|A|^2}$ relations on a set A .

Binary Relations on a Set (*cont.*)

Example: Consider these relations on the **set of integers**:

$$R_1 = \{(a,b) \mid a \leq b\},$$

$$R_4 = \{(a,b) \mid a = b\},$$

$$R_2 = \{(a,b) \mid a > b\},$$

$$R_5 = \{(a,b) \mid a = b + 1\},$$

$$R_3 = \{(a,b) \mid a = b \text{ or } a = -b\},$$

$$R_6 = \{(a,b) \mid a + b \leq 3\}.$$

Note that these relations are on an infinite set and each of these relations is an infinite set.

Which of these relations contain each of the pairs?

$(1,1)$, $(1, 2)$, $(2, 1)$, $(1, -1)$, and $(2, 2)$?

Solution:

$(1,1)$ is in R_1 , R_3 , R_4 , and R_6

$(1,2)$ is in R_1 and R_6

$(2,1)$ is in R_2 , R_5 , and R_6

$(1, -1)$ is in R_2 , R_3 , and R_6

$(2,2)$ is in R_1 , R_3 , and R_4

Reflexive Relations

Definition: R is *reflexive* iff $(a,a) \in R$ for every element $a \in A$.
Written symbolically, R is reflexive if and only if

$$\forall x[x \in U \longrightarrow (x,x) \in R]$$

Example: The following relations on the integers are reflexive:

$$R_1 = \{(a,b) \mid a \leq b\},$$

$$R_3 = \{(a,b) \mid a = b \text{ or } a = -b\},$$

$$R_4 = \{(a,b) \mid a = b\}.$$

If $A = \emptyset$ then the empty relation is reflexive vacuously. That is the *empty relation on an empty set is reflexive!*

The following relations are *not reflexive*:

$$R_2 = \{(a,b) \mid a > b\} \text{ (note that } 3 \not> 3),$$

$$R_5 = \{(a,b) \mid a = b + 1\} \text{ (note that } 3 \neq 3 + 1),$$

$$R_6 = \{(a,b) \mid a + b \leq 3\} \text{ (note that } 4 + 4 \not\leq 3).$$

Symmetric Relations

Definition: R is *symmetric* iff $(b,a) \in R$ whenever $(a,b) \in R$ for all $a,b \in A$. Written symbolically, R is symmetric if and only if

$$\forall x \forall y [(x,y) \in R \longrightarrow (y,x) \in R]$$

Example: The following relations on the integers are symmetric:

$$R_3 = \{(a,b) \mid a = b \text{ or } a = -b\},$$

$$R_4 = \{(a,b) \mid a = b\},$$

$$R_6 = \{(a,b) \mid a + b \leq 3\}.$$

The following are not symmetric:

$$R_1 = \{(a,b) \mid a \leq b\} \text{ (note that } 3 \leq 4, \text{ but } 4 \not\leq 3),$$

$$R_2 = \{(a,b) \mid a > b\} \text{ (note that } 4 > 3, \text{ but } 3 \not> 4),$$

$$R_5 = \{(a,b) \mid a = b + 1\} \text{ (note that } 4 = 3 + 1, \text{ but } 3 \neq 4 + 1).$$

Antisymmetric Relations

Definition: A relation R on a set A such that for all $a, b \in A$ if $(a, b) \in R$ and $(b, a) \in R$, then $a = b$ is called *antisymmetric*. Written symbolically, R is antisymmetric if and only if

$$\forall x \forall y [(x, y) \in R \wedge (y, x) \in R \rightarrow x = y]$$

Example: The following relations on the integers are antisymmetric:

$$R_1 = \{(a, b) \mid a \leq b\},$$

$$R_2 = \{(a, b) \mid a > b\},$$

$$R_4 = \{(a, b) \mid a = b\},$$

$$R_5 = \{(a, b) \mid a = b + 1\}.$$

For any integer, if $a \leq b$ and $b \leq a$, then $a = b$.

The following relations are not antisymmetric:

$$R_3 = \{(a, b) \mid a = b \text{ or } a = -b\}$$

(note that both $(1, -1)$ and $(-1, 1)$ belong to R_3),

$$R_6 = \{(a, b) \mid a + b \leq 3\} \text{ (note that both } (1, 2) \text{ and } (2, 1) \text{ belong to } R_6).$$

Transitive Relations

Definition: A relation R on a set A is called **transitive** if whenever $(a,b) \in R$ and $(b,c) \in R$, then $(a,c) \in R$, for all $a,b,c \in A$. Written symbolically, R is transitive if and only if

$$\forall x \forall y \forall z [(x,y) \in R \wedge (y,z) \in R \rightarrow (x,z) \in R]$$

- **Example:** The following relations on the integers are transitive:

$$R_1 = \{(a,b) \mid a \leq b\},$$

$$R_2 = \{(a,b) \mid a > b\},$$

$$R_3 = \{(a,b) \mid a = b \text{ or } a = -b\},$$

$$R_4 = \{(a,b) \mid a = b\}.$$

For every integer, $a \leq b$ and $b \leq c$, then $b \leq c$.

The following are not transitive:

$$R_5 = \{(a,b) \mid a = b + 1\} \text{ (note that both } (3,2) \text{ and } (4,3) \text{ belong to } R_5, \text{ but not } (3,3)),$$

$$R_6 = \{(a,b) \mid a + b \leq 3\} \text{ (note that both } (2,1) \text{ and } (1,2) \text{ belong to } R_6, \text{ but not } (2,2)).$$

Combining Relations

- Given two relations R_1 and R_2 , we can combine them using **basic set operations** to form new relations such as $R_1 \cup R_2$, $R_1 \cap R_2$, $R_1 - R_2$, and $R_2 - R_1$.
- Example:** Let $A = \{1,2,3\}$ and $B = \{1,2,3,4\}$. The relations $R_1 = \{(1,1),(2,2),(3,3)\}$ and $R_2 = \{(1,1),(1,2),(1,3),(1,4)\}$ can be combined using basic set operations to form new relations:

$$R_1 \cup R_2 = \{(1,1),(1,2),(1,3),(1,4),(2,2),(3,3)\}$$

$$R_1 \cap R_2 = \{(1,1)\}$$

$$R_1 - R_2 = \{(2,2),(3,3)\}$$

$$R_2 - R_1 = \{(1,2),(1,3),(1,4)\}$$

Composition


Definition: Suppose

- R_1 is a relation from a set A to a set B .
- R_2 is a relation from B to a set C .

Then the *composition* (or *composite*) of R_2 and R_1 , **denoted as $R_2 \circ R_1$** , is a relation from A to C where

- if (x,y) is a member of R_1 and (y,z) is a member of R_2 , then (x,z) is a member of $R_2 \circ R_1$.

Representing the Composition of a Relation



$$R_1 \circ R_2 = \{(b, x), (b, z)\}$$

Powers of a Relation

Definition: Let R be a binary relation on A . Then the powers R^n of the relation R can be defined recursively by:

- $R^1 = R$
- $R^{n+1} = R^n \circ R$

Theorem 1: The relation R on a set A is transitive iff $R^n \subseteq R$ for $n = 1, 2, 3, \dots$

Proof:

(1) $\leftarrow$ (if)

$R^n \subseteq R \Rightarrow R^2 \subseteq R \Rightarrow$ Based on the definition of $R \circ R$, $\forall (a,b) \in R$ and $(b,c) \in R \rightarrow (a,c) \in R^2 \subseteq R$, i.e., R is transitive.

(2) $\rightarrow$ (only if)

We use mathematical induction to prove: R is transitive $\rightarrow R^n \subseteq R$ for $n = 1, 2, 3, \dots$

Basis Step: $R^1 \subseteq R$

Inductive Step: (Inductive Hypothesis: $R^n \subseteq R, n \in \mathbb{N}$)

$\forall (a,b) \in R^{n+1}$, $(R^{n+1} = R^n \circ R)$, $\exists x \in A$ such that $(a,x) \in R$ and $(x,b) \in R^n$

Since $R^n \subseteq R$, $(x,b) \in R$

Since R is transitive: $(a,x) \in R$ and $(x,b) \in R \rightarrow (a,b) \in R$

Now we have $\forall (a,b) \in R^{n+1} \rightarrow (a,b) \in R$, that is $R^{n+1} \subseteq R$

It follows from the mathematical induction that R is transitive $\rightarrow R^n \subseteq R$ for $n = 1, 2, 3, \dots$

Representing Relations

Section 9.3

Representing Relations Using Matrices

- A relation between finite sets can be represented using a **zero-one matrix** (零壹矩陣).
- Suppose R is a relation from $A = \{a_1, a_2, \dots, a_m\}$ to $B = \{b_1, b_2, \dots, b_n\}$.
 - The elements of the two sets can be listed in any particular arbitrary order. When $A = B$, we use the same ordering.
- The relation R is represented by the matrix $M_R = [m_{ij}]$, where

$$m_{ij} = \begin{cases} 1 & \text{if } (a_i, b_j) \in R, \\ 0 & \text{if } (a_i, b_j) \notin R. \end{cases}$$

- The matrix representing R has a 1 as its (i,j) entry when a_i is related to b_j and a 0 if a_i is not related to b_j .

Examples of Representing Relations Using Matrices

Example 1: Suppose that $A = \{1,2,3\}$ and $B = \{1,2\}$. Let R be the relation from A to B containing (a,b) if $a \in A$, $b \in B$, and $a > b$. What is the matrix representing R (assuming $a_1=1$, $a_2=2$, $a_3=3$, and $b_1=1$, $b_2=2$)?

Solution: Because $R = \{(2,1), (3,1), (3,2)\}$, the matrix is

$$M_R = \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 1 & 1 \end{bmatrix}.$$

Examples of Representing Relations Using Matrices (*cont.*)

Example 2: Let $A = \{a_1, a_2, a_3\}$ and $B = \{b_1, b_2, b_3, b_4, b_5\}$. Which ordered pairs are in the relation R represented by the matrix

$$M_R = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 1 & 0 \\ 1 & 0 & 1 & 0 & 1 \end{bmatrix} ?$$

Solution: Because R consists of those ordered pairs (a_i, b_j) with $m_{ij} = 1$, it follows that:

$$R = \{(a_1, b_2), (a_2, b_1), (a_2, b_3), (a_2, b_4), (a_3, b_1), (a_3, b_3), (a_3, b_5)\}.$$

Matrices of Relations **on a Set**

- A **square matrix**
- If R is a **reflexive** relation, all the elements on the main diagonal of M_R are equal to 1.

$$\begin{bmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & & & \ddots \\ & & & & 1 & \\ & & & & & 1 \end{bmatrix}$$

- R is a **symmetric** relation, if and only if $m_{ij} = 1$ whenever $m_{ji} = 1$. R is an antisymmetric relation, if and only if $m_{ij} = 0$ or $m_{ji} = 0$ when $i \neq j$.

$$\begin{bmatrix} 1 & 1 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

(a) Symmetric

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

(b) Antisymmetric

Example of a Relation on a Set

Example 3: Suppose that the relation R on a set is represented by the matrix

$$M_R = \begin{bmatrix} 1 & 1 & 0 \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix}.$$

Is R reflexive, symmetric, and/or antisymmetric?


Solution: Because all the diagonal elements are equal to 1, R is reflexive. Because M_R is symmetric, R is symmetric and not antisymmetric because both $m_{1,2}$ and $m_{2,1}$ are 1.

Representing Relations Using Digraphs

Definition: A *directed graph* (有向圖), or *digraph*, consists of a set V of *vertices* (or *nodes*) together with a set E of ordered pairs of elements of V called *edges* (or *arcs*). The vertex a is called the *initial vertex* (起點) of the edge (a,b) , and the vertex b is called the *terminal vertex* (終點) of this edge.

- An edge of the form (a,a) is called a *loop* (迴圈).


Example 7: A drawing of the directed graph with vertices a, b, c , and d , and edges (a, b) , (a, d) , (b, b) , (b, d) , (c, a) , (c, b) , and (d, b) is shown here.



Example 8: What are the ordered pairs in the relation represented by this directed graph?

Solution: The ordered pairs in the relation are


$(1,3)$, $(1,4)$, $(2,1)$, $(2,2)$, $(2,3)$, $(3,1)$, $(3,3)$, $(4,1)$, and $(4, 3)$



Determining which Properties a Relation has from its Digraph


- *Reflexivity*: A **loop** must be present at **all vertices** in the graph.
- *Symmetry*: If (x,y) is an edge, then so is (y,x) .
- *Antisymmetry*: If (x,y) with $x \neq y$ is an edge, then (y,x) is **not** an edge.
- *Transitivity*: If (x,y) and (y,z) are edges, then so is (x,z) .

Determining which Properties a Relation has from its Digraph – Example 1




- *Reflexive?* No, not every vertex has a loop
- *Symmetric?* Yes (trivially), there is no edge from one vertex to another
- *Antisymmetric?* Yes (trivially), there is no edge from one vertex to another
- *Transitive?* Yes, (trivially) since there is no edge from one vertex to another

Determining which Properties a Relation has from its Digraph – Example 2




- *Reflexive?* No, there are no loops
- *Symmetric?* No, there is an edge from a to b , but not from b to a
- *Antisymmetric?* No, there is an edge from d to b and b to d
- *Transitive?* No, there are edges from a to c and from c to b , but there is no edge from a to d

Determining which Properties a Relation has from its Digraph – Example 3



- *Reflexive?* No, there are no loops
- *Symmetric?* No, for example, there is no edge from c to a
- *Antisymmetric?* Yes, whenever there is an edge from one vertex to another, there is not one going back
- *Transitive?* No, there is no edge from a to b

Determining which Properties a Relation has from its Digraph – Example 4



- *Reflexive?* No, there are no loops
- *Symmetric?* No, for example, there is no edge from d to a
- *Antisymmetric?* Yes, whenever there is an edge from one vertex to another, there is not one going back
- *Transitive?* Yes (trivially), there are no two edges where the first edge ends at the vertex where the second edge begins

Closures of Relations

Section 9.4

Closures of Relations

- For any property P , the “ P closure” of relation R on a set A is defined as the “smallest” superset of R that has property P .
- The *reflexive closure* of a relation R on A is obtained by adding (a,a) to R for each $a \in A$. i.e., it is $R \cup \Delta$, $\Delta = \{(a,a) \mid a \in A\}$
- The *symmetric closure* of R is obtained by adding (b,a) to R for each (a,b) in R . i.e., it is $R \cup R^{-1}$
- The *transitive closure* or *connectivity relation* of R is obtained by repeatedly adding (a,c) to R for each $(a,b), (b,c)$ in R .
 - i.e., it is

$$R^* = \bigcup_{n \in \mathbb{Z}^+} R^n$$

Paths in Digraphs/Binary Relations

- A *path of length n* from node a to b in the directed graph G (or the binary relation R) is a sequence $(a, x_1), (x_1, x_2), \dots, (x_{n-1}, b)$ of n ordered pairs in E_G (or R).
 - An empty sequence of edges is considered a path of length 0 from a to a .
 - If any path from a to b exists, then we say that a is *connected to b* . (“You can get there from here.”)
- A path of length $n \geq 1$ from a to a is called a *circuit* or a *cycle*.
- Note that there exists a path of length n from a to b in R if and only if $(a, b) \in R^n$.

Simple Transitive Closure Alg.

A procedure to compute R^* with 0-1 matrices.

procedure *transClosure*($\mathbf{M}_R$:rank- n 0-1 mat.)

A := **B** := $\mathbf{M}_R$; {note **A** represents R^i }

for $i := 2$ to n **begin**

A := $\mathbf{A} \odot \mathbf{M}_R$; **B** := **B** $\vee$ **A** {boolean product; join}

end


return **B** {Alg. takes $\Theta(n^4)$ time}

$$R^* = \bigcup_{n \in \mathbb{Z}^+} R^n$$

Roy-Warshall's Transitive closure Alg.


Main idea: a path exists between two vertices i, j , iff

- there is an edge from i to j ; or
- there is a path from i to j going through vertex 1; or
- there is a path from i to j going through vertex 1 and/or 2; or
- there is a path from i to j going through vertex 1, 2, and/or 3; or
- ...
- there is a path from i to j going through any of the other vertices




R_0

0	0	1	0
1	0	0	1
0	0	0	0
0	1	0	0




R_1

0	0	1	0
1	0	1	1
0	0	0	0
0	1	0	0




R_2

0	0	1	0
1	0	1	1
0	0	0	0
1	1	1	1



R_3

0	0	1	0
1	0	1	1
0	0	0	0
1	1	1	1



R_4

0	0	1	0
1	1	1	1
0	0	0	0
1	1	1	1

Roy-Warshall Algorithm

- Uses only $\Theta(n^3)$ operations!

Procedure *Warshall*($\mathbf{M}_R$: rank- n 0-1 matrix)

$\mathbf{W} := \mathbf{M}_R$

path from i to k and from k to j using just $1, \dots, k-1$


for $k := 1$ **to** n

for $i := 1$ **to** n

for $j := 1$ **to** n

$w_{ij} := \mathbf{w}_{ij} \vee (\mathbf{w}_{ik} \wedge \mathbf{w}_{kj})$

return $\mathbf{W}$ {this represents R^* }



$w_{ij} = 1$ means there is a path from i to j going only through nodes $\leq k$

Equivalence Relations

Section 9.5

Equivalence Relations

Definition 1: A relation on a set A is called an *equivalence relation* if it is reflexive, symmetric, and transitive.

Definition 2: Two elements a , and b that are related by an equivalence relation are called *equivalent*. The notation $a \sim b$ is often used to denote that a and b are equivalent elements with respect to a particular equivalence relation.

Strings

Example: Suppose that R is the relation on the set of strings of English letters such that aRb if and only if $l(a) = l(b)$, where $l(x)$ is the length of the string x . Is R an equivalence relation?

Solution: Show that all of the properties of an equivalence relation hold.

- *Reflexivity*: Because $l(a) = l(a)$, it follows that aRa for all strings a .
- *Symmetry*: Suppose that aRb . Since $l(a) = l(b)$, $l(b) = l(a)$ also holds and bRa .
- *Transitivity*: Suppose that aRb and bRc . Since $l(a) = l(b)$, and $l(b) = l(c)$, $l(a) = l(c)$ also holds and aRc .

Congruence Modulo m (模 m 同餘)

Example: Let m be an integer with $m > 1$. Show that the relation

$$R = \{(a, b) \mid a \equiv b \pmod{m}\}$$

is an equivalence relation on the set of integers.

Solution: Note that $a \equiv b \pmod{m}$ if and only if m divides $a - b$.

- *Reflexivity*: $a \equiv a \pmod{m}$ since $a - a = 0$ is divisible by m since $0 = 0 \cdot m$.
- *Symmetry*: Suppose that $a \equiv b \pmod{m}$. Then $a - b$ is divisible by m , and so $a - b = km$, where k is an integer. It follows that $b - a = (-k)m$, so $b \equiv a \pmod{m}$.
- *Transitivity*: Suppose that $a \equiv b \pmod{m}$ and $b \equiv c \pmod{m}$. Then m divides both $a - b$ and $b - c$. Hence, there are integers k and l with $a - b = km$ and $b - c = lm$. We obtain by adding the equations:

$$a - c = (a - b) + (b - c) = km + lm = (k + l)m.$$

Therefore, $a \equiv c \pmod{m}$.

Divides

Example: Show that the “divides” relation on the set of positive integers is **not** an equivalence relation.

Solution: The properties of reflexivity, and transitivity do hold, but the relation is not symmetric. Hence, “divides” is not an equivalence relation.

- *Reflexivity*: $a \mid a$ for all a .
- *Not Symmetric*: For example, $2 \mid 4$, but $4 \nmid 2$. Hence, the relation is not symmetric.
- *Transitivity*: Suppose that a divides b and b divides c . Then there are positive integers k and l such that $b = ak$ and $c = bl$. Hence, $c = a(kl)$, so a divides c . Therefore, the relation is transitive.

Equivalence Classes

Definition 3: Let R be an equivalence relation on a set A . The set of all elements that are related to an element a of A is called the *equivalence class* of a . The *equivalence class of a with respect to R* is denoted by $[a]_R$. When only one relation is under consideration, we can write $[a]$, without the subscript R , for this equivalence class.

Note that $[a]_R = \{s \mid (a,s) \in R\}$.

- If $b \in [a]_R$, then b is called a *representative of this equivalence class*. Any element of a class can be used as a representative of the class.
- The *equivalence classes of the relation congruence modulo m* are called the *congruence classes modulo m* . The congruence class of an integer a modulo m is denoted by $[a]_m$, so $[a]_m = \{\dots, a-2m, a-m, a, a+m, a+2m, \dots\}$. For example,

$$[0]_4 = \{\dots, -8, -4, 0, 4, 8, \dots\}$$

$$[1]_4 = \{\dots, -7, -3, 1, 5, 9, \dots\}$$

$$[2]_4 = \{\dots, -6, -2, 2, 6, 10, \dots\}$$

$$[3]_4 = \{\dots, -5, -1, 3, 7, 11, \dots\}$$

Equivalence Classes and Partitions

Theorem 1: let R be an **equivalence relation** on a set A .
These statements for elements a and b of A are equivalent:

- (i) aRb
- (ii) $[a] = [b]$
- (iii) $[a] \cap [b] \neq \emptyset$

Proof: We show that (i) implies (ii).

Assume that aRb . $\forall c \in [a]$, aRc .

Because aRb and R is symmetric, bRa .

Because R is transitive: bRa and $aRc \rightarrow bRc$. Hence, $c \in [b]$.

Therefore, $[a] \subseteq [b]$.

A similar argument (omitted here) shows that $[b] \subseteq [a]$.


Since $[a] \subseteq [b]$ and $[b] \subseteq [a]$, we have shown that $[a] = [b]$.

(see text for proof that (ii) implies (iii) and (iii) implies (i))

Partition of a Set

Definition: A *partition* (分割) of a set S is a collection of disjoint nonempty subsets of S that have S as their union. In other words, the collection of subsets A_i , where $i \in I$ (where I is an index set), forms a partition of S if and only if

- $A_i \neq \emptyset$ for $i \in I$,
- $A_i \cap A_j = \emptyset$ when $i \neq j$,
- and $\bigcup_{i \in I} A_i = S$.



A Partition of a Set

An Equivalence Relation Partitions a Set

- Let R be an equivalence relation on a set A . The union of all the equivalence classes of R is all of A , since an element a of A is in its own equivalence class $[a]_R$. In other words,

$$\bigcup_{a \in A} [a]_R = A.$$

- From Theorem 1, $[a]_R \cap [b]_R \neq \emptyset \rightarrow [a]_R = [b]_R$
it follows that $[a]_R \neq [b]_R \rightarrow [a]_R \cap [b]_R = \emptyset$, i.e.,
these equivalence classes are either equal or disjoint
- Therefore, **different equivalence classes form a partition of A** , because they split A into disjoint subsets.

An Equivalence Relation Partitions a Set (*continued*)

Theorem 2: Let R be an equivalence relation on a set S . Then the equivalence classes of R form a partition of S . Conversely, given a partition $\{A_i \mid i \in I\}$ of the set S , there is an equivalence relation R that has the sets $A_i, i \in I$, as its equivalence classes.

Proof: We have already shown the first part of the theorem

For the second part, assume that $\{A_i \mid i \in I\}$ is a partition of S .

Let R be the relation on S consisting of the pairs (x, y) where x and y belong to the same subset A_i in the partition. We must show that R satisfies the properties of an equivalence relation.

- **Reflexivity:** For every $a \in S$, $(a, a) \in R$, because a is in the same subset as itself.
- **Symmetry:** If $(a, b) \in R$, then b and a are in the same subset of the partition, so $(b, a) \in R$.
- **Transitivity:** If $(a, b) \in R$ and $(b, c) \in R$, then a and b are in the same subset of the partition, as are b and c . Since the subsets are disjoint and b belongs to both, the two subsets of the partition must be identical. Therefore, $(a, c) \in R$ since a and c belong to the same subset of the partition.

Partial Orderings

Section 9.6

Partial Orderings

Definition 1: A relation R on a set S is called a *partial ordering* (偏序), or *partial order*, if it is reflexive, antisymmetric, and transitive.

A set S together with a partial ordering R is called a *partially ordered set*, or *poset* (偏序集), and is denoted by (S, R) .

Members of S are called *elements* of the poset.

Partial Orderings (*continued*)

Example 1: Show that the “greater than or equal” relation ($\geq$) is a partial ordering on the set of integers.

- *Reflexivity*: $a \geq a$ for every integer a .
- *Antisymmetry*: If $a \geq b$ and $b \geq a$, then $a = b$.
- *Transitivity*: If $a \geq b$ and $b \geq c$, then $a \geq c$.

Partial Orderings (*continued*)

Example 2: Show that the divisibility relation ($|$) is a partial ordering on the set of **positive** integers.

- *Reflexivity*: $a | a$ for all integers a .
 - *Antisymmetry*: If a and b are positive integers with $a | b$ and $b | a$, then $a = b$.
 - *Transitivity*: Suppose that a divides b and b divides c . Then there are positive integers k and l such that $b = ak$ and $c = bl$. Hence, $c = a(kl)$, so a divides c . Therefore, the relation is transitive.
- $(\mathbf{Z}^+, |)$ is a poset.

Partial Orderings (*continued*)

Example 3: Show that the inclusion relation ($\subseteq$) is a partial ordering on **the power set of a set S** .

- *Reflexivity*: $A \subseteq A$ whenever A is a subset of S .
 - *Antisymmetry*: If A and B are two subsets of S with $A \subseteq B$ and $B \subseteq A$, then $A = B$.
 - *Transitivity*: If $A \subseteq B$ and $B \subseteq C$, then $A \subseteq C$.
- $(P(S), \subseteq)$ is a poset

Comparability (可比較性)

Definition 2: The elements a and b of a poset $(S, \preceq)$ are *comparable* if either $a \preceq b$ or $b \preceq a$. When a and b are elements of S so that neither $a \preceq b$ nor $b \preceq a$, then a and b are called *incomparable*.

The symbol $\preceq$ is used to denote the relation in any poset.

Definition 3: If $(S, \preceq)$ is a poset and every two elements of S are comparable, S is called a *totally ordered set* or *linearly ordered set*, and $\preceq$ is called a *total order* (全序) or a *linear order*. A totally ordered set is also called a *chain* (鏈).

Definition 4: $(S, \preceq)$ is a *well-ordered set* if it is a poset such that $\preceq$ is a *total ordering* and every nonempty subset of S has a least element.


Lexicographic Order

Definition: Given two posets $(A_1, \preceq_1)$ and $(A_2, \preceq_2)$, the *lexicographic ordering* on $A_1 \times A_2$ is defined by specifying that (a_1, a_2) is less than (b_1, b_2) , that is, $(a_1, a_2) < (b_1, b_2)$, either if $a_1 <_1 b_1$ or if $a_1 = b_1$ and $a_2 <_2 b_2$.

- The lexicographic order (with the equality being added) is a partial ordering on the Cartesian product of two posets
- **Example:** Consider strings of lowercase English letters. A lexicographic ordering can be defined using the ordering of the letters in the alphabet. This is the same ordering as that used in dictionaries.
 - *discreet* < *discrete*, because these strings differ in the seventh position and $e < t$.
 - *discreet* < *discreetness*, because the first eight letters agree, but the second string is longer.

Hasse Diagrams

Definition: A *Hasse diagram*(哈斯圖) is a visual representation of a partial ordering that remove edges that must be present because of the reflexive and transitive properties.



- A partial ordering is shown in (a) of the figure above.
- The loops due to the reflexive property are deleted in (b).
- The edges that must be present due to the transitive property are deleted in (c).
- The Hasse diagram for the partial ordering (a), is depicted in (c).


Procedure for Constructing a Hasse Diagram

- To represent a finite poset $(S, \preceq)$ using a Hasse diagram, start with the directed graph of the relation:
 - **Remove the loops** (a, a) present at every vertex due to the reflexive property.
 - Remove all edges (x, y) for which there is an element $z \in S$ such that $x \prec z$ and $z \prec y$. These are the edges that must be present due to the transitive property.
 - Arrange each edge so that its initial vertex is below the terminal vertex.
 - Remove all the arrows, because all edges point upwards toward their terminal vertex.

Example

Let $A = \{1, 2, 3, 9, 19\}$ and consider the “divides” relation on A :

For all $a, b \in A$, $a \mid b \Leftrightarrow b = ka$ for some integer k .




Example

Eliminate the loops at all the vertices.

Eliminate all arrows whose existence is implied by the transitive property.

Eliminate the direction indicators on the arrows.



Construct the Hasse diagram of $(P(\{a, b, c\}), \subseteq)$.

The elements of $P(\{a, b, c\})$ are


$\emptyset$

$\{a\}, \{b\}, \{c\}$

$\{a, b\}, \{a, c\}, \{b, c\}$


$\{a, b, c\}$

The digraph is



Maximal and Minimal Elements


- a is a *maximal* in the poset $(S, \leq)$ if there is no element $b \in S$ such that $a \leq b$.
- Similarly, an element of a poset is called *minimal* if it is not greater than any element of the poset. That is, a is *minimal* if there is no element $b \in S$ such that $b \leq a$.
- It is possible to have multiple minimals and maximals.



Greatest Element

Least Element

- a is the *greatest element* in the poset $(S, \leq)$ if $b \leq a$ for all $b \in S$.
- Similarly, an element of a poset is called the *least element* if it is less or equal than all other elements in the poset. That is, a is the *least element* if $a \leq b$ for all $b \in S$.



Upper bound, Lower bound


- Sometimes it is possible to find (an) element(s) that is greater than all the elements in a subset A of a poset $(S, \preceq)$.

If u is an element of S such that $a \preceq u$ for all elements $a \in A$, then u is called an *upper bound* of A .

- Likewise, there may be (an) element(s) less than all the elements in A . If l is an element of S such that $l \preceq a$ for all elements $a \in A$, then l is called a *lower bound* of A .


Least Upper Bound, Greatest Lower Bound

- The element x is called the *least upper bound* (lub) of the subset A if x is an upper bound that is less than every other upper bound of A .
- The element y is called the *greatest lower bound* (glb) of A if y is a lower bound of A and $z \leq y$ whenever z is a lower bound of A .




Lattices

- A partially ordered set in which *every pair of elements* has both a least upper bound and a greatest lower bound is called a *lattice*.



Lattices

- For the poset represented by the Hasse diagram
 - Consider element 1 and 3
 - Upper bound of $\{1\}$ is 1,2,4,5
 - Upper bound of $\{3\}$ is 3,2,4,5
 - Upper bound of $\{1,3\}$ is 2,4,5
 - There is no lub of $\{1,3\}$
 - There is no glb of $\{1,3\}$ either
 - The poset is not a lattice




Topological Sorting

- A **total ordering** $\preceq$ is said to be **compatible** with the **partial ordering** R if $a \preceq b$ whenever $a R b$.
- Constructing a compatible total ordering from a partial ordering is called **topological sorting**. (not unique)
- An Algorithm: To sort a poset (S, R)
 1. Selected a (any) minimal element and put it in the list.
Delete it from S
 2. Continue until all elements appear in the list (and S is void)

Example

- Consider the set $A = \{2, 3, 4, 6, 18, 24\}$ ordered by the “divides” relation. The Hasse diagram follows:



- The ordinary “less than or equal to” relation $\leq$ on this set is a topological sorting for it since for positive integers a and b , if $a \mid b$ then $a \leq b$.