Table S2 Receptor structures and docking boxes

A. Receptor identification

Role	Target	PDB	Auth chains	Cognate CCD	Cognate instance	Retained / removed components
Primary	EGFR	3POZ	A	03P	A:1023	Selected protein chain retained
Primary	HER2	3RCD	A	03P	A:9001	Selected protein chain retained
Primary	JAK1	6N7A	AB	KEV	A:1203	Chains A and B retained; PTR converted to TYR
Primary	JAK2	8BXH	A	C87	A:1201	PTR converted to TYR
Primary	TYK2	3LXP	A	IZA	A:1183	PTR converted to TYR
Primary	PIK3CA	4L23	A	X6K	A:1101	p85 removed
Primary	mTOR	4JT6	A	X6K	A:2601	Second mTOR copy and mLST8 removed
Primary	AChE	4EY7	AB	E20	A:604	AChE chains A and B retained
Primary	BChE	4BDS	A	THA	A:701	Selected protein chain retained
Primary	F2	4UDW	HIL	N6L	H:1249	Heavy chain, light chain and hirudin retained
Primary	F10	2JKH	AL	BI7	A:1244	Heavy and light chains retained
Primary	PPARG	9V8H	AB	BRL	A:501	PPARG and PG08-NL peptide retained
Primary	PPARA	6LXA	A	EPA	A:502	Selected protein chain retained
Primary	PPARD	5U3Q	B	7UJ	B:501	PPARD chain B retained
Alternative	AChE	4EY6	AB	GNT	A:604	AChE chains A and B retained
Alternative	BChE	1P0M	A	CHT	A:607	Selected protein chain retained
Alternative	F2	3SHC	HIL	B01	H:3	Heavy chain, light chain and hirudin retained
Alternative	F10	2Y5F	AL	XWG	A:1244	Heavy and light chains retained
Alternative	PPARA	6KAX	A	PLM	A:502	Selected protein chain retained
Alternative	PPARD	5U46	A	7T1	A:505	Selected protein chain retained

Coordinates and dimensions are in angstrom. Waters and cocrystal ligands are removed from the receptor. Box centers are bounding-box midpoints; padding is 5 Å with minimum edge 20 Å. Chain identifiers are author chain identifiers. Formal prepared files and preparation exceptions, including observed-fragment templates, are documented in the accompanying source files. Alternative structures cover only the six conditions used in the current analysis; no Q110 alternative-receptor result is included.

Table S2 Receptor structures and docking boxes

B. Docking-box coordinates

Role	Target	PDB	Center (x, y, z), Å	Size (x, y, z), Å
Primary	EGFR	3POZ	18.7460, 31.8320, 11.6260	20.7500, 20.0000, 21.2180
Primary	HER2	3RCD	12.4800, 2.9635, 28.0150	21.1620, 22.1630, 20.0000
Primary	JAK1	6N7A	5.9110, -24.3175, 21.4945	20.0000, 20.5970, 20.0000
Primary	JAK2	8BXH	-12.5897, 14.5939, 21.2626	24.0687, 20.0000, 20.0000
Primary	TYK2	3LXP	-4.4985, 25.5775, -31.0640	20.0000, 20.0000, 20.0000
Primary	PIK3CA	4L23	32.5530, 45.3340, 42.0935	20.0000, 20.0000, 20.0000
Primary	mTOR	4JT6	52.1230, -0.0840, -47.6645	20.4680, 20.0000, 20.0000
Primary	AChE	4EY7	-13.9875, -43.9055, 27.1085	23.3410, 20.0000, 20.3550
Primary	BChE	4BDS	133.0760, 116.1130, 41.3350	20.0000, 20.0000, 20.0000
Primary	F2	4UDW	129.7390, -14.4025, 164.9460	20.6000, 20.0000, 20.0000
Primary	F10	2JKH	-3.6430, 10.7695, 22.7710	20.0000, 20.0000, 21.1180
Primary	PPARG	9V8H	-5.4540, -5.3730, -23.5915	20.0000, 20.1400, 20.0000
Primary	PPARA	6LXA	11.9975, 5.7420, -7.4430	20.0000, 20.2000, 23.4320
Primary	PPARD	5U3Q	45.4160, -37.6795, 103.9765	20.0000, 20.0000, 22.4490
Alternative	AChE	4EY6	-10.4920, -43.5255, 29.4535	20, 20, 20
Alternative	BChE	1P0M	133.5840, 116.7965, 40.3950	20, 20, 20
Alternative	F2	3SHC	16.7620, -12.0230, 22.9065	20.5680, 20, 20
Alternative	F10	2Y5F	-4.6240, 10.6510, 22.4765	20, 20.2300, 21.4290
Alternative	PPARA	6KAX	12.0005, 6.0880, -7.6585	20, 20, 22.6890
Alternative	PPARD	5U46	45.4215, -1.7525, 135.5685	20, 20, 20.4870

Coordinates and dimensions are in angstrom. Waters and cocrystal ligands are removed from the receptor. Box centers are bounding-box midpoints; padding is 5 Å with minimum edge 20 Å. Chain identifiers are author chain identifiers. Formal prepared files and preparation exceptions, including observed-fragment templates, are documented in the accompanying source files. Alternative structures cover only the six conditions used in the current analysis; no Q110 alternative-receptor result is included.