

4.1.5

(page 124)

(a) If a and b are consecutive integers, show that $a^2 + b^2 + (ab)^2$ is a perfect square.

proof)

Let $b = a + 1$ and substituting into the original expression we arrive at:

$$a^2 + (a + 1)^2 + (a^2 + a)^2 = a^2 + a^2 + 2a + 1 + a^4 + 2a^3 + a^2$$

$$a^4 + 2a^3 + 2a^2 + a^2 + 2a + 1 = a^4 + a^2 + 1 + 2a^3 + 2a^2 + 2a$$

We recognize that $(x + y + z)^2 = x^2 + y^2 + z^2 + 2xy + 2yz + 2xz$ thus,

$$a^4 + a^2 + 1 + 2a^3 + 2a^2 + 2a = (a^2 + a + 1)^2$$

which is a perfect square.

(b) If $2a$ is the harmonic mean of b and c (i.e. $2a = 2/(1/b + 1/c)$), show that the sum of the squares of the three numbers a, b , and c is the square of a rational number.

proof)

We are to prove that

$$a^2 + b^2 + c^2 = \frac{p^2}{q^2}, p, q \in \mathbb{Z}, q \neq 0$$

. We substitute $2a = \frac{1}{(\frac{1}{b} + \frac{1}{c})}$ into the expression, and we get:

$$a^2 + b^2 + c^2 = \left(\frac{1}{(\frac{1}{b} + \frac{1}{c})}\right)^2 + b^2 + c^2$$

$$\left(\frac{bc}{bc(\frac{1}{b} + \frac{1}{c})}\right)^2 + b^2 + c^2 = \left(\frac{bc}{c + b}\right)^2 + b^2 + c^2$$

$$\frac{b^2c^2}{(c + b)^2} + b^2 + c^2 = \frac{b^2c^2}{(b + c)^2} + \frac{b^2(b + c)^2}{(b + c)^2} + \frac{c^2(b + c)^2}{(b + c)^2}$$

$$\frac{b^2(b + c)^2 + b^2c^2 + c^2(b + c)^2}{(b + c)^2} = \frac{b^2(b^2 + 2bc + c^2) + b^2c^2 + c^2(b^2 + 2bc + c^2)}{(b + c)^2}$$

$$\frac{b^4 + 2b^3c + b^2c^2 + b^2c^2 + b^2c^2 + 2bc^3 + c^4}{(b + c)^2} = \frac{b^4 + c^4 + b^2c^2 + 2b^3c + 2bc^3 + 2b^2c^2}{(b + c)^2}$$

We recognize that $(x + y + z)^2 = x^2 + y^2 + z^2 + 2xy + 2yz + 2xz$ thus,

$$\frac{b^4 + c^4 + b^2c^2 + 2b^3c + 2bc^3 + 2b^2c^2}{(b + c)^2} = \frac{(b^2 + c^2 + bc)^2}{(b + c)^2}$$

(c) If N differs from two successive squares between which it lies by x and y respectively, prove that $N - xy$ is a square.

proof)

Let's assume the two successive squares are a^2 and $(a + 1)^2$ and N is placed such that:

$$N - a^2 = x$$

$$(a + 1)^2 - N = y$$

and we need to show that $N - xy$ is a square.

$$N - xy = N - (N - a^2)((a + 1)^2 - N)$$

$$N - (N - a^2)(a^2 + 2a + 1 - N) = N - (a^2N + 2aN + N - N^2 - a^4 - 2a^3 - a^2 + a^2N)$$

$$N - (a^2N + 2aN + N - N^2 - a^4 - 2a^3 - a^2 + a^2N) = N - a^2N - 2aN - N + N^2 + a^4 + 2a^3 + a^2 - a^2N$$

$$N^2 + a^4 + a^2 - 2a^2N - 2aN + 2a^3$$

We recognize that $(x + y + z)^2 = x^2 + y^2 + z^2 + 2xy + 2yz + 2xz$ thus,

$$N^2 + a^4 + a^2 - 2a^2N - 2aN + 2a^3 = (N - a^2 - a)^2$$

which is a square.