



# Malaria

Elizabeth A Ashley, Aung Pyae Phyo, Charles J Woodrow

Lancet 2018; 391: 1608–21

Published Online

April 6, 2018

[http://dx.doi.org/10.1016/S0140-6736\(18\)30324-6](http://dx.doi.org/10.1016/S0140-6736(18)30324-6)

Myanmar–Oxford Clinical Research Unit, Yangon, Myanmar (E A Ashley MD); Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine, University of Oxford, Oxford, UK (E A Ashley, C J Woodrow MD), Shoklo Malaria Research Unit, Mae Sot, Thailand (A Pyae Phyo MD); and Mahidol–Oxford Tropical Medicine Research Unit (MORU), Faculty of Tropical Medicine, Mahidol University, Bangkok, Thailand (A Pyae Phyo, C J Woodrow)

Correspondence to: Dr Elizabeth A Ashley, Myanmar Oxford Clinical Research Unit, 32A1 Kokkine Swimming Club Lane, Yangon, Myanmar [liz@tropmedres.ac](mailto:liz@tropmedres.ac)

Following unsuccessful eradication attempts there was a resurgence of malaria towards the end of the 20th century. Renewed control efforts using a range of improved tools, such as long-lasting insecticide-treated bednets and artemisinin-based combination therapies, have more than halved the global burden of disease, but it remains high with 445 000 deaths and more than 200 million cases in 2016. Pitfalls in individual patient management are delayed diagnosis and overzealous fluid resuscitation in severe malaria. Even in the absence of drug resistance, parasite recurrence can occur, owing to high parasite densities, low host immunity, or suboptimal drug concentrations. Malaria elimination is firmly back as a mainstream policy but resistance to the artemisinin derivatives, their partner drugs, and insecticides present major challenges. Vaccine development continues on several fronts but none of the candidates developed to date have been shown to provide long-lasting benefits at a population level. Increased resources and unprecedented levels of regional cooperation and societal commitment will be needed if further substantial inroads into the malaria burden are to be made.

## Introduction

Malaria is a vector-borne parasitic tropical disease found in 91 countries worldwide.<sup>1</sup> Of more than 120 *Plasmodium* species infecting mammals, birds, and reptiles, only six are known to infect human beings regularly. *Plasmodium falciparum* produces high levels of blood-stage parasites that sequester in critical organs in all age groups and cause severe anaemia in African children, in whom the vast majority of malaria deaths occur. *Plasmodium vivax* usually produces milder disease, but can be severe, and recurrent episodes bring significant associated morbidity. *Plasmodium malariae*, and the morphologically indistinguishable sympatric species *Plasmodium ovale curtisi* and *Plasmodium ovale wallikeri* are understudied, but severity of illness is generally similar to uncomplicated vivax malaria. *Plasmodium knowlesi* is a primarily zoonotic infection encountered in southeast Asia that can cause severe malaria.

## Epidemiology

Malaria is a disease of tropical and subtropical regions, having been eradicated from temperate countries steadily over the last 100 years. It is transmitted by the bite of the female *Anopheles* mosquito. Disease incidence depends on environmental suitability for local vectors in terms of altitude, climate, vegetation, and implementation of


control measures, and hence is inextricably linked to poverty, natural disasters, and war. Less common transmission routes are from mother to child, or via blood transfusion, a rare occurrence in non-endemic countries thanks to blood donor screening procedures, but a significant risk in resource-poor settings.<sup>2,3</sup> Predictions as to the effect of climate change on global malaria distribution in the future vary, but have suggested the population at risk of malaria will increase, in particular in tropical highland areas.<sup>4</sup>

*Plasmodium falciparum* and *P vivax* are the predominant species worldwide with an estimated incidence of 207 million and 8·5 million cases respectively in 2016.<sup>1</sup> The great majority of falciparum malaria occurs in sub-Saharan Africa (approximately 190 million cases) where transmission remains intense in many locations, although there is considerable variation in incidence within and between countries.<sup>5,6</sup> Vivax malaria is much less common in this region because the human population is largely Duffy antigen negative (see predisposing factors below). In Asia and Oceania, malaria case numbers are generally lower and proportions caused by *P vivax* and *P falciparum* are similar, whereas in the Americas, vivax malaria cases exceed falciparum by more than two times.<sup>1</sup>

*P malariae* and *P ovale* have a global distribution but incidence is low with *P ovale* found mainly in Africa and southeast Asia. Macaques are the natural hosts of *P knowlesi*. In Malaysia, which has a high burden of knowlesi malaria, cases were initially misdiagnosed as *P malariae* because of morphological similarities when examined by light microscopy.<sup>7</sup> The true global burden of disease is unknown; however, although this parasite is capable of being transmitted by *Anopheles dirus*, an important vector of human malaria, it is predominantly a zoonosis. Human infections with other simian malarias such as *Plasmodium cynomolgi* and *Plasmodium simium* can occur. They are assumed to be infrequent events with the caveat that routine microscopic examination could fail to distinguish these from the more common species.<sup>8,9</sup>

## Search strategy and selection criteria

We searched PubMed, Embase, and the Cochrane Library for all clinical trials, meta-analyses, systematic reviews, and diagnostic test accuracy studies published between Jan 1, 2014, and July 31, 2017, in the English language, using the search term “malaria”. International malaria treatment guidelines and policy documents on the WHO website were also consulted. References cited in these publications were screened to identify other recent original journal articles and highly relevant older references—eg, definitive trial reports, or articles linked to a particular discovery.



**Figure 1: Human stages of the malaria lifecycle**

The inset illustrates the differing outcomes of treatment of falciparum malaria with quinine (QN) and artesunate (ART). In most cases the majority of parasites are young ring stages at clinical presentation (A). Quinine kills sequestered parasites but not circulating rings which continue to develop after treatment (B); in non-immune patients treated with quinine, sequestration is the dominant method of parasite clearance (C). Artesunate brings the advantage of rapid action against both circulating rings and sequestered parasites; killing of early stages produces pyknotic parasite forms (D), which the spleen removes by pitting, leading to potentially large numbers of once-infected red cells (E) with reduced lifespans and the phenomenon of late haemolysis. These cells contain the PfHRP2 antigen (indicated by green outline) explaining why, despite parasite clearance, this persists. Individuals with partial immunity to malaria clear both parasitised and once-infected red cells by antibody-mediated phagocytosis in the spleen relatively rapidly. P=Plasmodium. RBC=red blood cell.

## Biology

The human phases of the malaria life cycle are shown in figure 1. Sporozoites are inoculated by the bite of an infected female *Anopheles* mosquito. The parasite undergoes a pre-erythrocytic liver stage which typically lasts for 1–2 weeks before the onset of the blood stage, where serial cycles of asexual replication produce rising parasite numbers and hence human disease. A subpopulation of intraerythrocytic parasites switches to sexual development,<sup>10</sup> producing female and male gametocytes.<sup>11</sup> These distinctive transitional stages transmit malaria to the mosquito via a blood meal. Male gametocytes exflagellate in the mosquito midgut and male and female gametes fuse to form a zygote that transforms into a mobile ookinete and passes through the gut wall. The oocyst releases sporozoites which migrate to the mosquito salivary glands, completing the lifecycle.

In vivax and ovale infections, a proportion of sporozoites become dormant hypnozoites, causing relapses months or years after the initial infection.

## Pathogenesis

Symptoms of malaria develop once the erythrocytic cycle produces a parasitaemia above a certain threshold (roughly 100 parasites per  $\mu\text{L}$ ). Incubation periods are typically 10–14 days for *P falciparum* or *P knowlesi*, 2–3 weeks for *P vivax* and *P ovale*, and 18 days or longer for *P malariae*; however, there is variation—eg some strains of *P vivax* have a 3–6 month primary incubation period.<sup>12,13</sup> Classic accounts describe periodic fever spikes at intervals corresponding to the erythrocytic cycle length

of the infecting species (48 h for *P falciparum*, *vivax*, or *ovale* and 72 h for *P malariae*), resulting from synchronisation of developmental stages, but such patterns are now observed rarely.<sup>12</sup>

*P falciparum* is unique in that erythrocytes containing mature parasites sequester inside small and medium-sized vessels, avoiding parasite clearance in the spleen but causing host endothelial cell injury and microvascular obstruction. Cytoadherence is mediated by *P falciparum* erythrocyte membrane protein 1 (PfEMP1), a clonally variant set of proteins exported to the infected erythrocyte surface and encoded by the *var* gene family. Subtypes of PfEMP1 bind to different endothelial receptors; for example, those that bind intercellular adhesion molecule-1 and endothelial protein C receptor are associated with cerebral malaria.<sup>14,15</sup> Infected erythrocytes also bind to uninfected cells (rosetting), and uninfected cells become less deformable, exacerbating microvascular obstruction.

The clinical effect of sequestration and associated endothelial dysfunction depends on the organ(s) involved. In the brain, it contributes towards coma, in the lungs it predisposes to respiratory failure, and in pregnant women, sequestration in the intervillous space of the placenta leads to placental malaria with the consequences of maternal anaemia, low birthweight, preterm labour, and increased risk of abortion and stillbirth.<sup>16,17</sup> Placental cytoadherence is mediated by binding to chondroitin sulphate A (CSA), specifically via the PfEMP1 variant VAR2CSA, and the effects are most severe in primigravid women.<sup>16</sup>

Anaemia is a common feature of malaria and is typically multifactorial in origin with red cell loss as the leading

**Panel 1: Diagnostic criteria for severe malaria<sup>20</sup>****Clinical criteria**

- Prostration
- Confusion or agitation (with Glasgow Coma Scale [GCS] >11)
- Coma (GCS ≤11 or Blantyre Coma Scale <3 in children)
- Respiratory distress (acidotic breathing)
- Convulsions
- Shock: prolonged capillary refill time (>2 s), with or without systolic blood pressure <80 mm Hg in adults (<70 in children)
- Pulmonary oedema (should be confirmed radiologically)
- Abnormal bleeding
- Jaundice
- Anuria
- Repeated vomiting

**Laboratory criteria**

- Haemoglobin <7 g/dL in adults, <5 g/dL in children
- Haemoglobinuria
- Hypoglycaemia (blood glucose <2.2 mmol/L or <40 mg/dL)
- Acidosis (ie, base deficit >8 meq/L or plasma bicarbonate <15 mmol/L or venous plasma lactate >5 mmol/L)
- Acute kidney injury (creatinine >3 mg/dL or urea >20 mmol/L)
- Asexual parasitaemia >10% of infected red blood cells (Note: national guidelines can vary—eg, UK parasitaemia cutoff is 2%<sup>21</sup>)

cause in acute infections as the spleen filters both infected and damaged uninfected red blood cells.<sup>18</sup> A degree of intravascular haemolysis also occurs, which can be massive. There is also bone marrow suppression and dyserythropoiesis.

Investigations in 2015 found some evidence of endothelial cell activation in vivax malaria and suggested that peripheral parasite density could underestimate total biomass but research into pathogenesis is at a very early stage compared to *P. falciparum*.<sup>19</sup>

**Clinical presentation**

Malaria is separated conveniently into two disease presentations: uncomplicated and severe. Symptoms of uncomplicated malaria are very non-specific, and can include fever, chills, body-aches, headache, cough, and diarrhoea, making clinical diagnosis unreliable. In non-endemic areas, taking an accurate travel history in all patients with fever is the key to making the diagnosis. Thrombocytopenia can provide another clue. The differential diagnosis will vary depending on location. Once malaria is suspected, the most appropriate course of action is to expedite laboratory testing (see Laboratory diagnosis).
















Severe falciparum malaria has specific diagnostic criteria. For rapid clinical assessment, a short list of

danger signs is used, which includes prostration, fast deep breathing (reflecting underlying acidosis), and impaired consciousness. A comprehensive list is shown in panel 1.

The most common manifestations of severe malaria are cerebral malaria, acute lung injury, which can progress to acute respiratory distress syndrome (in up to 25% of cases), acute kidney injury, typically presenting as acute tubular necrosis, and acidosis.<sup>22</sup> Lactic acid predominates but other acids have been identified in adults with severe malaria, including hydroxyphenyllactic acid, and α-hydroxybutyric and β-hydroxybutyric acids.<sup>23</sup> Severe anaemia (without major organ dysfunction) is a common presentation in children. Other differences in disease presentation in children compared with adults are more frequent seizures (in 60–80%), hypoglycaemia, and concomitant sepsis, and less frequent pulmonary oedema and renal failure.<sup>20,24</sup> The case fatality rate of treated cerebral malaria is usually 10–20% and can reach 50% in pregnant women.<sup>12</sup> Coma can be profound with extensor posturing, positive pout reflex, and teeth-grinding. Neuroimaging typically shows some evidence of brain swelling but this is less prominent in adults than in children, in whom brain swelling is strongly associated with a fatal outcome.<sup>25–27</sup> Characteristic fundoscopic findings in cerebral malaria include retinal whitening, changes in blood vessel colour, and haemorrhages. Papilloedema is unusual.<sup>27</sup> Two patterns of acute kidney injury are described in malaria: one in patients with severe malaria with multiorgan failure and the second in patients who have been successfully treated and do not have evidence of other organ involvement.<sup>20</sup>

**Laboratory diagnosis**

Confirming the presence of parasites in all malaria cases ensures species-specific antimalarial treatment and points to other illnesses in negative cases. The gold standard for malaria diagnosis remains light microscopy of stained blood films, thick films providing sensitivity and thin films allowing speciation and quantitation (figure 2). However, rapid diagnostic tests (RDTs) now predominate as the first-line investigation<sup>28</sup> with a wide range of devices available. Given the distribution of species, in much of Africa a *P. falciparum*-only test based on the highly expressed histidine-rich protein 2 (PfHRP2) antigen is often used; this can remain positive for several weeks after parasite clearance because of persisting pitted (once-infected) red blood cells.<sup>29</sup> Elsewhere, RDTs often incorporate both an HRP2-detecting strip for sensitive falciparum detection and a pan-species strip that detects the lactate dehydrogenase enzyme of all human malarias, although this is relatively insensitive for *P. knowlesi* diagnosis. In Latin America, HRP2 gene deletions mean that HRP2-based tests are unreliable,<sup>30</sup> and evidence is emerging that the problem could extend to Africa.<sup>31</sup> Very high falciparum parasitaemias can also produce negative results due to the prozone effect.<sup>32</sup>

Human malaria				
	Rings	Trophozoites	Schizonts	
<i>P falciparum</i>				<ul style="list-style-type: none"><li>Parasitised red cells (pRBCs) not enlarged</li><li>RBCs containing mature trophozoites sequestered in deep vessels</li><li>Total parasite biomass = circulating parasites + sequestered parasites</li></ul>
<i>P vivax</i>				<ul style="list-style-type: none"><li>Parasites prefer young red cells</li><li>pRBCs enlarged</li><li>Trophozoites are amoeboid in shape</li><li>All stages present in peripheral blood</li></ul>
<i>P malariae</i>				<ul style="list-style-type: none"><li>Parasites prefer old red cells</li><li>pRBCs not enlarged</li><li>Trophozoites tend to have a band shape</li><li>All stages present in peripheral blood</li></ul>
<i>P ovale</i>				<ul style="list-style-type: none"><li>pRBCs slightly enlarged and have an oval shape, with tufted ends</li><li>All stages present in peripheral blood</li></ul>
<i>P knowlesi</i>				<ul style="list-style-type: none"><li>pRBCs not enlarged</li><li>Trophozoites, pigment spreads inside cytoplasm; like <i>P malariae</i>, band forms may be seen</li><li>Multiple invasion and high parasitaemia can be seen like <i>P falciparum</i></li><li>All stages present in peripheral blood</li></ul>

**Figure 2: Blood films showing microscopic appearance of the human malariae**

All parasite stages are visible in peripheral blood except *P falciparum* RBCs containing mature trophozoites where the vast majority are sequestered in deep vessels. Thin blood films were prepared from specimens taken from patients with clinical malaria, stained with modified Field's stain and examined by light microscopy under oil immersion at x1000 magnification. P=*Plasmodium*.

The threshold of detection for these standard methods is approximately 50 parasites per  $\mu\text{L}$  (microscopy) and 200 parasites per  $\mu\text{L}$  for PfHRP2-based *P falciparum* RDTs (several times higher for non-falciparum RDTs). Nucleic acid amplification-based tests provide much greater sensitivity (often below one parasite per  $\mu\text{L}$ ).<sup>33</sup>

### Case management

The treatment of malaria, particularly that of *P falciparum*, was revolutionised by the introduction of the artemisinin derivatives in the 1990s, a group of semisynthetic compounds produced from *qinghaosu* (artemisinin), a natural product of the sweet wormwood plant (*Artemisia annua*). Artemisinins are rapidly effective, safe, and well tolerated. Their discovery by China's Project 523 was acknowledged by the award of the 2015 Nobel Prize to Tu Youyou.

### Management of severe malaria

All patients diagnosed with severe malaria, including women in all trimesters of pregnancy, should receive parenteral artesunate without delay (panel 2).<sup>20</sup> A higher

mg/kg bodyweight dose of artesunate is given to children weighing less than 20 kg. Artesunate was shown to be vastly superior to quinine in large trials (35% [95% CI 18.5–47.6] mortality reduction in southeast Asian adults and 22% [95% CI 8.1–36.9] reduction in African children).<sup>34,35</sup> If quinine is prescribed, a loading dose must be given and the second dose administered 8 h after the start of the first infusion.<sup>36</sup> Intramuscular artemether is another option but was inferior to parenteral artesunate for preventing malaria deaths in Asian adults.<sup>37</sup> Parenteral quinidine is still recommended as an alternative treatment in the USA but is associated with substantial cardiotoxicity. Hypoglycaemia (blood glucose  $<2.2$  mmol/L) is a serious complication of malaria and is aggravated by quinine therapy, especially in pregnant women. Management of hypovolaemia and acidosis requires cautious fluid replacement with crystalloids to reduce the risk of pulmonary oedema.<sup>38</sup> The dangers of overly aggressive volume replacement have been shown in adults and children. In the multicentre FEAST study of more than 3000 critically ill African children with

**Panel 2: Drug treatment of malaria<sup>53</sup>****Severe malaria***Initial treatment*

- Intravenous artesunate (2–4 mg/kg per dose hours 0, 12, and 24; then every 24 h; in patients with bodyweight <20 kg unit dose is 3.0 mg/kg)

*Alternatives*

- Intravenous quinine infusion; loading dose of 20 mg/kg (given over 4 h) at admission then 10 mg/kg (given over 2 h) every 8 h
- Intramuscular artemether injection: 3.2 mg/kg initial dose, then 1.6 mg/kg every 24 h

*Once able to eat and drink*

- Oral treatment with an artemisin-based combination therapy (ACT) for 3 days (not mefloquine due to increased risk of post-malaria neurological syndrome)

**Uncomplicated malaria***Uncomplicated Plasmodium falciparum\* or Plasmodium knowlesi malaria*

- Artemether–lumefantrine 1.4–4 mg/kg of artemether and 10–16 mg/kg of lumefantrine twice daily for 3 days with food containing fat, or
- Dihydroartemisinin–piperaquine 4 mg/kg of dihydroartemisinin and 18 mg/kg of piperaquine once daily for 3 days (children with bodyweight <25 kg should receive at least 2.5 mg/kg per day of dihydroartemisinin and 20 mg/kg per day piperaquine), or
- Artesunate 4 mg/kg per day with mefloquine 8 mg/kg per day for 3 days, or
- Artesunate 4 mg/kg per day with amodiaquine 10 mg base/kg per day for 3 days, or
- Artesunate 4 mg/kg per day for 3 days with single dose sulfadoxine–pyrimethamine (25 mg/kg–1.25 mg/kg)

*Chloroquine-sensitive Plasmodium vivax†, Plasmodium ovale, Plasmodium malariae‡*

- Chloroquine 10 mg base/kg per day at hour 0 and hour 24 followed by 5 mg base/kg at hour 48

Target doses and full dosing information of different formulations and non-artemisinin-based treatment options are shown in the appendix. \*Choice of ACT to treat *P. falciparum* depends on local resistance patterns; mefloquine is contraindicated in patients with a history of epilepsy or neuropsychiatric disorders. †Chloroquine-resistant *P. vivax* is treated with one of the ACTs (except artesunate plus sulfadoxine–pyrimethamine). ‡Initial treatment of *P. vivax* or *P. ovale* should be followed by a course of primaquine to prevent relapse, if no contraindications (glucose-6-phosphate-dehydrogenase deficiency, pregnancy, age <6 months).

sepsis and impaired perfusion, more than half (57%) of whom had malaria, those given fluid boluses had a higher mortality at 48 h than those who were not (relative risk [RR] for bolus vs control, 1.45; 95% CI, 1.13–1.86).<sup>39</sup> Severe anaemia (haemoglobin <50 g/L) requires urgent correction but evidence regarding optimal transfusion practices is absent. A trial is ongoing to define this in children.<sup>24,40,41</sup> Seizures should be controlled with benzodiazepines; if recurrent,

loading with longer-acting anticonvulsants should be considered with close monitoring for respiratory depression. Routine seizure prophylaxis with phenobarbital given to Kenyan children with cerebral malaria in a placebo-controlled trial was associated with increased rates of respiratory depression and death.<sup>42</sup> Other supportive measures depend on the clinical manifestations and the level of care available. Early renal replacement therapy is recommended. If there is no access to endotracheal intubation, a nasogastric tube should be passed but enteral feeding delayed for 72 h to reduce the risk of aspiration pneumonia.<sup>43</sup>

Patients with cerebral malaria should have blood cultures taken, a lumbar puncture performed, and broad-spectrum antibiotics commenced, pending negative culture results and clinical improvement. Severe malaria is associated with concomitant bacterial sepsis, with non-typhoidal *Salmonella* bacteraemia described frequently in African children.<sup>44</sup> Co-infections with other tropical diseases are unusual in travellers but more common in hyperendemic countries.

Patients with severe malaria should have frequent monitoring of vital signs, level of consciousness, glucose, renal function, and haemoglobin. Monitoring parasite density (every 6–12 h) confirms parasite clearance (typically within 72 h) but clinical improvement often takes considerably longer.<sup>45,46</sup> Once the patient is conscious, able to eat and drink, and has received at least 24 h of parenteral therapy, oral follow-on treatment with an artemisinin-based combination treatment is advised.


**Adjunctive therapies for severe malaria**

Various adjunctive therapies have been evaluated without success in terms of improving the outcome from severe malaria, including steroids, anti-TNF, mannitol, *N*-acetylcysteine (antioxidant properties), exchange transfusion, and levamisole (inhibitor of sequestration).<sup>47–49</sup>

**Severe vivax and knowlesi malaria**

Severe vivax malaria tends to occur in patients with comorbidities in endemic countries and has not been observed frequently in returned travellers. Respiratory failure and acute kidney injury have been reported repeatedly in fatal cases.<sup>50,51</sup> One of the most common manifestations of severe vivax malaria in children is respiratory distress.<sup>52</sup> Coma is a rare occurrence in patients with vivax parasitaemia.<sup>54</sup> The main burden of serious morbidity and mortality from vivax malaria is secondary to severe anaemia, with young children particularly vulnerable. Rates of hospital admission and death from vivax malaria approach those for falciparum malaria in some parts of the world such as Papua in Indonesia.<sup>52,55</sup> Severe knowlesi malaria is associated with high parasite densities and similarly can present as acute kidney injury, shock, or respiratory failure.<sup>56</sup>





**Figure 3: Global distribution of drug-resistant *Plasmodium falciparum***

Countries are shown according to the level of resistance of local *P. falciparum*; the 13 countries approaching elimination (as defined in the World Malaria Report 2016) are also shown. The inset graphs show WHO estimates of global annual malaria case numbers and deaths from 2000 to 2015 (with 95% upper and lower uncertainty intervals).<sup>58</sup> ACT=artemisinin-based combination therapy. Created with mapchart.net ©.

### Management of uncomplicated malaria

The main considerations when prescribing anti-malarials are the infecting species and the risk of drug resistance. Comprehensive, evidence-based guidelines for the treatment of uncomplicated malaria are available on the WHO website.<sup>53</sup> Given the global spread of *P. falciparum* resistant to chloroquine and antifolates, artemisinin-based combination treatments (ACTs) are recommended for the treatment of falciparum malaria or falciparum mixed with non-falciparum species, except in the first trimester of pregnancy (see appendix). ACTs consist of a combination of an artemisinin derivative that rapidly reduces parasitaemia and a partner drug that removes residual parasites over a longer period. These properties make them the drugs of choice to treat knowlesi malaria as well.<sup>56</sup> The leading ACTs in use are artemether–lumefantrine, artesunate–amodiaquine, dihydroartemisinin–piperaquine, artesunate–mefloquine,

and artesunate plus sulfadoxine–pyrimethamine. The first four ACTs on this list exist as fixed-dose combinations, with paediatric formulations available. Artemether–lumefantrine should be given with milk or food containing fat to enhance lumefantrine absorption. The ACTs were highly efficacious against all *P. falciparum* until recently when numbers of treatment failures increased in parts of southeast Asia (figure 3).

Atovaquone–proguanil is an alternative non-artemisinin-based treatment that is useful for individual patients (eg, returned travellers without hyperparasitaemia, or in combination with artesunate plus primaquine for patients in endemic countries who have failed treatment with standard ACTs); however, it is not recommended for widespread implementation in endemic countries, because of the propensity for rapid emergence of atovaquone resistance.<sup>53</sup> Quinine remains efficacious, although it requires a long course of treatment, is poorly

See Online for appendix

tolerated, particularly by children, and needs to be combined with a second agent such as doxycycline or clindamycin.

Uncomplicated vivax, malariae, and ovale malaria are treated with chloroquine, unless chloroquine-resistant *Plasmodium vivax* is likely (Indonesia, Oceania) when an ACT is used (panel 2).<sup>57</sup> This should be followed by primaquine to eradicate dormant hypnozoites.

The decision to admit a patient with uncomplicated malaria to hospital will depend on the setting and local guidelines. It is common practice to admit non-immune returned travellers for an initial period of observation until clinical improvement and a fall in parasitaemia are observed.

### Management of uncomplicated malaria in pregnancy

Early detection of malaria in pregnancy is vital. Uncomplicated falciparum malaria in the first trimester is treated with a 7-day course of quinine and clindamycin. Safety data of first trimester ACT use have been reviewed and are reassuring, and treatment guidelines will be reviewed in the near future.<sup>59</sup> After week 12 of gestation, treatment is as for non-pregnant patients.<sup>60</sup> Vivax malaria in pregnancy is treated with chloroquine unless resistance is suspected (when quinine should be given), but radical cure with primaquine is contraindicated as the glucose-6-phosphate-dehydrogenase (G6PD) status of the foetus cannot be ascertained. The pharmacokinetic properties of several antimalarial drugs are different in pregnancy with a tendency towards lower drug exposure.<sup>61</sup> This makes treatment failure more likely, especially in non-immune women.

### Congenital malaria

The diagnosis of congenital malaria is easy to miss, especially if the mother is asymptomatic. The clinical presentation mimics neonatal sepsis. Parenteral treatment (artesunate or quinine) should be given for at least the first dose in congenital falciparum malaria. Follow-on treatment is with an ACT. Congenital vivax malaria can be treated with oral chloroquine unless the infant is very unwell, in which case parenteral drugs should be used, or if chloroquine resistance is likely, then either an ACT or quinine should be given.

### Adjunctive primaquine therapy

To reduce the risk of relapse from dormant hypnozoites in the liver, a course of the 8-aminoquinoline primaquine is added to the treatment of vivax or ovale malaria.<sup>62</sup> Primaquine causes dose-dependent haemolytic anaemia in patients with G6PD deficiency, hence testing for this enzymopathy is recommended; however, worldwide, access to testing is poor. The standard primaquine treatment regimen is long (14 days) and, as a result, difficult to adhere to. Doubling the daily dose to 1 mg base/kg bodyweight and shortening the course to 7 days was shown to result in an increased risk of

clinically significant haemolysis in G6PD heterozygotes (9/17 [53%] patients had >25% fractional haematocrit reduction compared with 2/16 [3%] G6PD heterozygotes taking the standard 0.5 mg/kg dose for 14 days;  $p=0.022$ ).<sup>63</sup> Failure of primaquine to suppress relapses has been linked to reduced metabolism of the drug in individuals with polymorphisms in the cytochrome P-450 isoenzyme 2D6.<sup>64</sup> The longer-acting 8-aminoquinoline tafenoquine is currently under review by stringent regulatory authorities to replace primaquine for radical cure as a single dose treatment but the risk of haemolysis remains.<sup>65</sup>

Primaquine is the only drug able to kill mature (stage V) gametocytes of *P. falciparum*. In endemic areas, prescription of a single low (0.25 mg/kg) dose of primaquine with an ACT is recommended to reduce the risk of onward transmission. This dose is considered safe in G6PD deficiency.

### Safety of the antimalarial drugs

Antimalarials can have serious side-effects and the frequency with which they occur varies between populations. Important examples include quinidine-induced cardiotoxicity, hypoglycaemia, and hypotension following quinine, which can also cause QT-prolongation, although much less frequently than following quinidine, hepatotoxicity and cutaneous hypersensitivity reactions to sulfadoxine-pyrimethamine (Stevens-Johnson syndrome in approximately 1/10 000 recipients), and neuropsychiatric reactions to mefloquine (approximately 1/200–1/2000 at treatment doses).<sup>53,66</sup>

### Treatment failure

Malaria treatments are not always curative.<sup>67–70</sup> Treatment failure usually presents as a recurrence of symptoms with detectable parasitaemia 2–6 weeks after an apparently successful treatment and is not always due to drug resistance. Alternative explanations include high parasite densities (particularly in non-immune individuals), poor drug bioavailability, non-adherence to therapy, and falsified or substandard antimalarials.<sup>71</sup> Relapse of vivax malaria is common after an episode of falciparum in southeast Asia (roughly 30% of cases).<sup>72</sup>

### Artemisinin-resistant falciparum malaria in southeast Asia

Like combination treatments in other areas of medicine, ACTs were introduced to prevent or delay development of resistance in a population over time.<sup>73</sup> After more than a decade of use in southeast Asia, artemisinin resistance was confirmed, manifested by a phenotype of delayed parasite clearance.<sup>74–76</sup> Use of artemisinin monotherapies was probably a contributing factor. The loss of the rapid parasitocidal effect of the artemisinin derivatives led predictably to worsening partner drug resistance (to mefloquine and piperazine) and reduced efficacy of the corresponding ACTs. Artemether-lumefantrine efficacy was poor in Cambodia in the early 2000s and has not

been reassessed since then. Molecular markers for parasite resistance to many of the commonly used drugs have been discovered.<sup>77–80</sup> High rates of ACT failure have now been reported from Cambodia, Thailand, and Vietnam.<sup>81–83</sup> A newer ACT, artesunate–pyronaridine, is likely to be approved for use in these countries in the near future. Triple artemisinin-based combinations (dihydroartemisinin–piperaquine with mefloquine and artemether–lumefantrine with amodiaquine) are being evaluated in an attempt to bridge the gap until new medicines become available.<sup>84</sup>

### New antimalarial drugs

Investment in antimalarial drug discovery and development, with the creation of product development partnerships in the early 2000s, have revitalised a near empty drug pipeline. Most of the drugs under development are blood schizontocides with a few exceptions.<sup>85–91</sup> There are efforts to develop alternative transmission-blocking drugs as an elimination tool.<sup>92</sup>

### Complications of malaria

Delayed haemolytic anaemia can follow artemisinin treatment of travellers with falciparum malaria.<sup>93</sup> The key event appears to be pitting, a splenic process whereby ring-stage parasites killed by artesunate are expelled from their host erythrocytes which return to the circulation, but with a reduced lifespan (figure 1).<sup>94</sup> This explains why the diagnostic antigen PfHRP2 persists for weeks after artemisinin treatment (it is exported to the erythrocyte periphery); indeed PfHRP2 concentration following parasite clearance predicts later haemolysis.<sup>29</sup> Treatment with quinine and other slow-acting drugs allows unhindered development of parasites until sequestration, so haemoglobin falls early. Delayed haemolysis can hence be considered a predictable event related to the early beneficial effect of artemisinins.<sup>94</sup> The problem appears to be somewhat less in African children,<sup>95</sup> presumably because of developing immunity, although cases have been documented.<sup>96</sup> It has also been suggested that post-artesunate delayed haemolysis could contribute to the high frequency of haemoglobinuria reported in eastern Uganda recently.<sup>97</sup>

Other haematological complications of malaria include hyper-reactive malarial splenomegaly (HMS), and rarely, splenic rupture. HMS is characterised by massive splenomegaly, raised IgM antimalarial antibodies, and anaemia. It responds to a prolonged course of weekly treatment with antimalarial drugs at prophylactic doses.<sup>98</sup>

Neurological complications following an episode of severe malaria range from a reversible post-malaria neurological syndrome to permanent deficits including visual, motor, or language disorders and epilepsy. In a study of African children with severe falciparum malaria, the prevalence of persistent neurological sequelae in the 4898 children who survived was 3.2%.<sup>35</sup> *P. malariae* can be complicated by anaemia and the nephrotic syndrome.<sup>99</sup>

### Panel 3: Factors predisposing to and protecting from severe malaria

#### Predisposing factors

##### Genetic factors

- Sickle cell disease
- Blood group B

##### Acquired factors

- Pregnancy and early post-partum period
- Malnutrition
- HIV
- Hyposplenism

#### Protective factors

##### Genetic factors

- Haemoglobin AS (sickle cell trait)
- Haemoglobin C
- Thalassaemia
- Glycophorins A and B
- Blood group O

### Malaria co-infection with HIV

HIV co-infection increases malaria severity and parasite densities tend to be higher. Co-trimoxazole prophylaxis is protective against malaria and concomitant antifolate antimalarial drugs should be avoided.<sup>100</sup> There are drug–drug interactions between common antimalarials and antiretroviral drugs—eg, efavirenz increases amodiaquine exposure and the risk of toxicity so these drugs should not be coadministered. Efavirenz decreases lumefantrine exposure but no formal recommendations have been made to adjust the dose.<sup>101,102</sup>

### Immunity

Repeated exposure over a long period to malaria leads to premunition—protection from disease but ongoing blood stage infection. In *P. falciparum* this involves sequential acquisition of antibodies to PfEMP1 subtypes. Asymptomatic falciparum or vivax parasitaemia are common in areas of high endemicity and use of more sensitive molecular detection methods for malaria has revealed that parasites can persist for years longer than thought possible previously.<sup>103</sup> Antibodies act on parasite blood stages to inhibit replication with complement fixation thought to play a part.<sup>104</sup> Antigen presentation takes place at different stages of the parasite lifecycle and T cells regulate both liver and blood stages of infection.<sup>105</sup> Immunity is lost steadily after an individual leaves an endemic area, or in a population with falling transmission.

### Predisposing and protective factors

A wide range of inherited and acquired factors influence an individual's chance of infection and severe illness with malaria (panel 3). Haldane's hypothesis suggesting that inherited human red cell conditions reflect evolutionary



#### Panel 4: Strategies to prevent malaria in different target groups

##### Chemoprophylaxis

###### Intermittent preventive treatment

- Pregnant women receive sulfadoxine–pyrimethamine (SP) at all antenatal care visits from second trimester (minimum dose interval of 1 month)
- Infants in moderate transmission areas receive SP with routine immunisations

###### Seasonal malaria chemoprevention

- Children aged 3–59 months in areas of seasonal transmission in the Sahel (intermittent SP plus amodiaquine at treatment doses, maximum four courses)

###### Fixed-term

- Travellers (atovaquone–proguanil, doxycycline, mefloquine, or primaquine)
- HIV-infected patients receiving co-trimoxazole as prophylaxis for opportunistic infections are protected against malaria

##### Vaccination

- RTS,S/AS01 in children (four doses for children aged 5–17 months) is the only registered malaria vaccine (still under evaluation)

##### Vector control or bite prevention

- Long-lasting insecticide-treated bednets: most effective in high-transmission areas where vectors rest indoors at night<sup>113</sup>
- Indoor residual spraying (with insecticides)
- Repellents (topical and spatial): convincing protective effect not shown

protection against malaria continues to be confirmed and refined.<sup>106</sup> Powerful studies combining clinical data from thousands of patients and controls with advanced sequencing methods have confirmed the protective role of structural variants of  $\beta$ -globin against severe malaria, the HbS variant (in heterozygous form causing sickle cell trait) reducing the risk of severe malaria by ten times, and the west African HbC variant to a lesser extent.<sup>107</sup> A consensus on the mechanism of protection has still to emerge, although suppression of parasite multiplication is probably involved given the protection against both cerebral malaria and severe anaemia.

The protective effect of X-linked G6PD deficiency, the commonest inherited human enzymopathy, against *P. falciparum* appears complex. The A allele is common in many parts of Africa, and male hemizygotes and female heterozygotes have reduced risk of cerebral malaria, but male hemizygotes and female homozygotes have increased risk of severe malarial anaemia.<sup>107</sup> In Asia, G6PD deficiency protects against vivax malaria.<sup>108,109</sup>

Genetic differences in red cell surface proteins also influence malaria. It has long been known that individuals negative for the Duffy antigen receptor for chemokines are

resistant to *P. vivax* and *P. knowlesi*, which preferentially invade reticulocytes via their Duffy binding protein. This protection has turned out not to be absolute; reports of vivax malaria in Duffy-negative individuals indicate that alternative ligands can mediate invasion.<sup>110,111</sup> Nevertheless, clinical vivax malaria remains rare in countries where the population is completely Duffy negative (most of sub-Saharan Africa).

Other families of red cell surface proteins also influence malaria infection, with individuals with blood group O relatively protected against severe falciparum malaria and those with group B at higher risk.<sup>107</sup> In 2017, a complex rearrangement in the polymorphic MNS blood group system (the DUP4 haplotype) producing hybrid glycophorin proteins was shown to be associated with protection against severe malaria in east Africa.<sup>112</sup>

#### Prevention of malaria

Malaria is prevented by chemoprophylaxis, vaccination, bite-avoidance and vector-control measures (panel 4).

##### Chemoprophylaxis

Target groups for chemoprophylaxis are pregnant women, young children, and travellers. Intermittent preventive therapy in pregnancy (IPTp) and infants has been slow to be scaled up in many areas and their impact is threatened by sulfadoxine–pyrimethamine resistance.<sup>114,115</sup> The use of alternative antimalarials such as dihydroartemisinin–piperaquine is under evaluation. Intermittent screening and treatment of pregnant women has not been demonstrated to be an effective alternative strategy to IPTp, attributed to lack of sensitivity of existing RDTs to detect malaria in pregnancy.<sup>115,116</sup> Seasonal malaria chemoprevention in children with sulfadoxine–pyrimethamine plus amodiaquine has been adopted widely and has reduced malaria in areas of highly seasonal transmission in the Sahel.<sup>117</sup>

To prevent malaria in travellers, the choice of chemoprophylaxis should consider the risks of malaria and drug resistance, which should be balanced with the risk of drug toxicity. Atovaquone–proguanil and doxycycline are prescribed as prophylaxis frequently. Weekly mefloquine at prophylactic doses is a convenient option but unpopular owing to concerns about neurotoxicity. Primaquine is a highly effective causal prophylactic agent with the added advantage of enhanced protection against vivax malaria (hypnozoites stages) but requires exclusion of G6PD deficiency.<sup>118</sup>

##### Vaccine development

Malaria subunit vaccines are designed to provide immunity against proteins exposed at critical stages of the lifecycle. Targeting sporozoite stages via one of the surface proteins that mediate homing to the liver and host cell traversal or invasion aims to reduce frequency of infection. The RTS,S/AS01 vaccine based on *P. falciparum* circumsporozoite protein is the most studied vaccine.

In a landmark study in African children, RTS,S/AS01 provided significant protection against falciparum malaria infection over a 3–4 year period; in older children, vaccine efficacy was 36·3% (95% CI 31·8–40·5) with a 20-month booster and 28·3% (95% CI 23·3–32·9) without.<sup>119</sup> However, efficacy was relatively lower (25·9% [95% CI 19·9–31·5] with booster and 18·3% [95% CI 11·7–24·4] without) in very young children (6–12 weeks old at first dose). Further, in contrast to earlier data, RTS,S/AS01 did not provide even efficacy across all strains.<sup>120</sup> Longer-term follow-up at one centre revealed a higher incidence of malaria in later years in vaccinated children with higher-than-average exposure to malaria.<sup>121</sup> An overall reduction in long-term mortality remains to be demonstrated. WHO is supporting pilot implementation of the four-dose regimen in children aged 5–17 months in three countries, allowing study of long-term outcomes, safety, and feasibility.

A contrasting approach to producing sporozoite-based immunity is the *P. falciparum* sporozoite (PfSPZ) vaccine, an intravenous injection of irradiation-attenuated sporozoites. PfSPZ has now entered clinical trials in Africa;<sup>122</sup> again the challenges are likely to centre on obtaining durable protection against all relevant strains.

The use of merozoite-stage proteins as vaccine targets aims to reduce asexual replication rate and hence protect against disease rather than produce sterile immunity, potentially allowing immunity to develop naturally while the vaccinee is protected from severe disease. Of the various extracellular merozoite proteins that collectively mediate erythrocyte invasion by *P. falciparum*,<sup>123</sup> there is considerable interest in targeting PfRh5 since its binding (to the Ok blood group antigen basigin) is critical for erythrocyte invasion.<sup>124</sup> Another asexual stage vaccine targets the product of the PfEMP1 VAR2CSA type aiming to prevent parasitised cell binding to CSA1 and hence protect against placental malaria.<sup>125</sup>

Transmission-blocking vaccines against sexual-stage antigens aim to generate antibodies that are ingested in the mosquito blood meal, potentially providing immunity at a population level. There is also increasing study of how vaccine responses to multiple targets might synergise to produce higher levels of overall protection,<sup>126</sup> although such an approach is likely to increase costs substantially. Vaccine development overall is complicated by the absence of reliable laboratory correlates of immunity.

### Vector control

More than 40 species of *Anopheles* are important malaria vectors. The mainstays of vector control are long-lasting insecticide (pyrethroid) treated bednets (LLINs) and indoor residual spraying with insecticides. LLINs have reduced morbidity and mortality from malaria and have the biggest impact in high transmission areas where vectors bite indoors at night, such as the highly successful *Anopheles gambiae* complex.<sup>113</sup> Their success is threatened by widespread pyrethroid resistance in anopheline vectors,

although the relationship between resistance and LLIN efficacy is not well characterised.<sup>127–129</sup> Alternative insecticides are needed urgently. In the meantime, addition of the synergist piperonyl butoxide to pyrethroid-treated bednets has been evaluated in some countries with mixed results. WHO have given an interim endorsement to this new class of products but have not recommended widespread deployment.<sup>130</sup> Bite-prevention strategies including topical repellents have not been shown to have an effect on malaria incidence.<sup>131,132</sup> Other approaches under consideration are mass treatment with ivermectin, which shortens mosquito survival, and transgenic mosquitoes.<sup>133–136</sup>

### From control to elimination

Progress towards malaria elimination is uneven. Indigenous cases in Europe, central Asia (north of Afghanistan), Sri Lanka, and several countries in Latin America are now extremely rare. However, in many sub-Saharan African countries, where transmission is highest, eliminating malaria has proved more difficult and there are signs that progress in this direction has stalled.<sup>1,6,137</sup> Areas with civil disruption have experienced substantial increases in malaria, exemplified by Venezuela. Pilot studies of mass drug administration (MDA) of ACT with single-dose primaquine to accelerate elimination of drug-resistant malaria in southeast Asia have taken place and early reports suggest it is effective and safe.<sup>138</sup>

### Controversies and uncertainties

There is more to be learned about the pathogenesis of severe vivax malaria, the relationship between pyrethroid resistance and LLIN efficacy, and the longer-term effects of MDA. Elimination of vivax malaria will require increased uptake of primaquine, which means addressing safety concerns in populations where G6PD deficiency is common. Elimination of human knowlesi malaria infections while there is a reservoir of parasites in macaques is another challenge. Whether artemisinin resistance spreads or pops up in different locations has been the subject of debate; recent evidence suggests both are true.<sup>139</sup> The effect of artemisinin resistance on artesunate efficacy in the treatment of severe falciparum malaria is unknown because severe malaria is now rare in southeast Asia. The loss of ring stage susceptibility to artemisinin raises the possibility that the treatment advantage over quinine has been eroded. Addition of parenteral quinine to artesunate has been proposed to provide a safety net for patients with presumed artemisinin resistance but there is no evidence to support this practice.

### Future perspectives

More than 130 years have passed since the protozoan cause of malaria was discovered. Over this period, there have been many scientific breakthroughs, with a subset

translating into interventions capable of reducing the burden of disease and death, notably the discovery and development of antimalarial drugs and insecticides.

The progress towards elimination in some countries shows that existing tools can be enough to eliminate malaria if the right conditions are in place: political commitment, access to health care, and adequate human and financial resources. There is evidence that access to high quality ACTs is still much too low (<25%) in some areas.<sup>140</sup> The spread of pyrethroid resistance among *Anopheles* vectors and increasing reports of ACT failures in southeast Asia signal that the window of opportunity to eliminate malaria with existing tools might be closing. Increased resources for disease control usually only come in times of crisis, but a concerted effort now could capitalise on recent gains and accelerate progress towards elimination.

#### Contributors

EAA performed the literature search. EAA and CJW wrote the first draft of text. APP drafted the tables and panels and figure 3. All authors reviewed and approved the final version of the manuscript.

#### Declaration of interests

We declare no competing interests.

#### Acknowledgments

We thank Kamolrat Silamut who created figure 2, and our colleagues and collaborators. The Myanmar–Oxford Clinical Research Unit, the Shoklo Malaria Research Unit, and the Mahidol–Oxford Tropical Medicine Research Unit (MORU) are part of the MORU Tropical Health Network, supported by the Wellcome Trust.

#### References

- WHO. World malaria report 2017. Geneva: World Health Organization, 2017.
- O'Brien SF, Delage G, Seed CR, et al. The epidemiology of imported malaria and transfusion policy in 5 nonendemic countries. *Transfus Med Rev* 2015; **29**: 162–71.
- Abdullah S, Karunamoorthi K. Malaria and blood transfusion: major issues of blood safety in malaria-endemic countries and strategies for mitigating the risk of *Plasmodium* parasites. *Parasitol Res* 2016; **115**: 35–47.
- Caminade C, Kovats S, Rocklov J, et al. Impact of climate change on global malaria distribution. *Proc Natl Acad Sci USA* 2014; **111**: 3286–91.
- Nkumama IN, O'Meara WP, Osier FH. Changes in malaria epidemiology in Africa and new challenges for elimination. *Trends Parasitol* 2017; **33**: 128–40.
- Snow RW, Sartorius B, Kyalo D, et al. The prevalence of *Plasmodium falciparum* in sub-Saharan Africa since 1900. *Nature* 2017; **550**: 515–18.
- Singh B, Kim Sung L, Matusop A, et al. A large focus of naturally acquired *Plasmodium knowlesi* infections in human beings. *Lancet* 2004; **363**: 1017–24.
- Ta TH, Hisam S, Lanza M, Jiram AI, Ismail N, Rubio JM. First case of a naturally acquired human infection with *Plasmodium cynomolgi*. *Malar J* 2014; **13**: 68.
- Brasil P, Zalis MG, de Pina-Costa A, et al. Outbreak of human malaria caused by *Plasmodium simium* in the Atlantic Forest in Rio de Janeiro: a molecular epidemiological investigation. *Lancet Glob Health* 2017; **5**: e1038–46.
- Josling GA, Llinas M. Sexual development in *Plasmodium* parasites: knowing when it's time to commit. *Nat Rev Microbiol* 2015; **13**: 573–87.
- Bousema T, Drakeley C. Epidemiology and infectivity of *Plasmodium falciparum* and *Plasmodium vivax* gametocytes in relation to malaria control and elimination. *Clin Microbiol Rev* 2011; **24**: 377–410.
- White NJ. Malaria. In: Farrar J, ed. *Manson's tropical diseases*, 23rd edn. London: Elsevier, 2013: 532–600.
- Coatney GR, Collins WE, Warren M, et al. CD-ROM. The primate malaras [original book published 1971]. Atlanta: Centers for Disease Control and Prevention; 2003.
- Lennartz F, Adams Y, Bengtsson A, et al. Structure-guided identification of a family of dual receptor-binding PfEMP1 that is associated with cerebral malaria. *Cell Host Microbe* 2017; **21**: 403–14.
- Turner L, Lavstsen T, Berger SS, et al. Severe malaria is associated with parasite binding to endothelial protein C receptor. *Nature* 2013; **498**: 502–05.
- Fried M, Duffy PE. Malaria during pregnancy. *Cold Spring Harb Perspect Med* 2017; **7**: a025551.
- Moore KA, Simpson JA, Wiladphaingern J, et al. Influence of the number and timing of malaria episodes during pregnancy on prematurity and small-for-gestational-age in an area of low transmission. *BMC Med* 2017; **15**: 117.
- Buffet PA, Safeukui I, Deplaine G, et al. The pathogenesis of *Plasmodium falciparum* malaria in humans: insights from splenic physiology. *Blood* 2011; **117**: 381–92.
- Barber BE, William T, Grigg MJ, et al. Parasite biomass-related inflammation, endothelial activation, microvascular dysfunction and disease severity in vivax malaria. *PLoS Pathog* 2015; **11**: e1004558.
- WHO. Severe malaria. *Trop Med Int Health* 2014; **19** (suppl 1): 7–131.
- Lalloo DG, Shingadia D, Bell DJ, et al. UK malaria treatment guidelines 2016. *J Infect* 2016; **72**: 635–49.
- Taylor WRJ, Hanson J, Turner GDH, White NJ, Dondorp AM. Respiratory manifestations of malaria. *Chest* 2012; **142**: 492–505.
- Herdman MT, Sriboonvorakul N, Leopold SJ, et al. The role of previously unmeasured organic acids in the pathogenesis of severe malaria. *Crit Care* 2015; **19**: 317.
- Maitland K. Management of severe paediatric malaria in resource-limited settings. *BMC Med* 2015; **13**: 42.
- Maude RJ, Barkhof F, Hassan MU, et al. Magnetic resonance imaging of the brain in adults with severe falciparum malaria. *Malar J* 2014; **13**: 177.
- Seydel KB, Kampondeni SD, Valim C, et al. Brain swelling and death in children with cerebral malaria. *N Engl J Med* 2015; **372**: 1126–37.
- Taylor TE, Molyneux ME. The pathogenesis of pediatric cerebral malaria: eye exams, autopsies, and neuroimaging. *Annals NY Acad Sci* 2015; **1342**: 44–52.
- Chiodini PL. Malaria diagnostics: now and the future. *Parasitology* 2014; **141**: 1873–79.
- Ndour PA, Larreche S, Moury O, et al. Measuring the *Plasmodium falciparum* HRP2 protein in blood from artesunate-treated malaria patients predicts post-artesunate delayed hemolysis. *Science Transl Med* 2017; **9**: eaaf9377.
- Gamboa D, Ho MF, Bendezu J, et al. A large proportion of *P. falciparum* isolates in the Amazon region of Peru lack pfhrp2 and pfhrp3: implications for malaria rapid diagnostic tests. *PLoS One* 2010; **5**: e8091.
- Berhane A, Russom M, Bahta I, Hagos F, Ghirmay M, Uqubay S. Rapid diagnostic tests failing to detect *Plasmodium falciparum* infections in Eritrea: an investigation of reported false negative RDT results. *Malar J* 2017; **16**: 105.
- Luchavez J, Baker J, Alcantara S, et al. Laboratory demonstration of a prozone-like effect in HRP2-detecting malaria rapid diagnostic tests: implications for clinical management. *Malar J* 2011; **10**: 286.
- Mathison BA, Pritt BS. Update on malaria diagnostics and test utilization. *J Clin Microbiol* 2017; **55**: 2009–17.
- Dondorp A, Nosten F, Stepniewska K, Day N, White N, South East Asian Quinine Artesunate Malaria Trial group. Artesunate versus quinine for treatment of severe falciparum malaria: a randomised trial. *Lancet* 2005; **366**: 717–25.
- Dondorp AM, Fanello CI, Hendriksen IC, et al. Artesunate versus quinine in the treatment of severe falciparum malaria in African children (AQUAMAT): an open-label, randomised trial. *Lancet* 2010; **376**: 1647–57.
- White NJ, Looareesuwan S, Warrell DA, et al. Quinine loading dose in cerebral malaria. *Am J Trop Med Hyg* 1983; **32**: 1–5.
- Esu E, Effa EE, Opie ON, Uwaoma A, Meremikwu MM. Artemether for severe malaria. *Cochrane Database Syst Rev* 2014; **9**: CD010678.
- Hanson JP, Lam SW, Mohanty S, et al. Fluid resuscitation of adults with severe falciparum malaria: effects on acid-base status, renal function, and extravascular lung water. *Crit Care Med* 2013; **41**: 972–81.

- 39 Maitland K, Kiguli S, Opoka RO, et al. Mortality after fluid bolus in African children with severe infection. *N Engl J Med* 2011; **364**: 2483–95.
- 40 Kiguli S, Maitland K, George EC, et al. Anaemia and blood transfusion in African children presenting to hospital with severe febrile illness. *BMC Med* 2015; **13**: 21.
- 41 ISRCTN Registry. Transfusion and treatment of severe anaemia in African children trial registration ISRCTN84086586. 2013. <https://doi.org/10.1186/ISRCTN84086586> (accessed Aug 20, 2017).
- 42 Crawley J, Waruiru C, Mithwani S, et al. Effect of phenobarbital on seizure frequency and mortality in childhood cerebral malaria: a randomised, controlled intervention study. *Lancet* 2000; **355**: 701–06.
- 43 Maude RJ, Hoque G, Hasan MU, et al. Timing of enteral feeding in cerebral malaria in resource-poor settings: a randomized trial. *PLoS One* 2011; **6**: e27273.
- 44 Takem EN, Roca A, Cunningham A. The association between malaria and non-typhoid *Salmonella* bacteraemia in children in sub-Saharan Africa: a literature review. *Malar J* 2014; **13**: 400.
- 45 Worldwide Antimalarial Resistance Network Parasite Clearance Study Group. Baseline data of parasite clearance in patients with falciparum malaria treated with an artemisinin derivative: an individual patient data meta-analysis. *Malar J* 2015; **14**: 359.
- 46 White NJ. Malaria parasite clearance. *Malar J* 2017; **16**: 88.
- 47 John CC, Kutamba E, Mugarura K, Opoka RO. Adjunctive therapy for cerebral malaria and other severe forms of *Plasmodium falciparum* malaria. *Expert Rev Anti-infect Ther* 2010; **8**: 997–1008.
- 48 Maude RJ, Silamut K, Plewes K, et al. Randomized controlled trial of levamisole hydrochloride as adjunctive therapy in severe falciparum malaria with high parasitemia. *J Infect Dis* 2014; **209**: 120–29.
- 49 Tan KR, Wiegand RE, Arguin PM. Exchange transfusion for severe malaria: evidence base and literature review. *Clin Infect Dis* 2013; **57**: 923–28.
- 50 Val F, Machado K, Barbosa L, et al. Respiratory complications of *Plasmodium vivax* malaria: systematic review and meta-analysis. *Am J Trop Med Hyg* 2017; **97**: 733–43.
- 51 Rahimi BA, Thakkestian A, White NJ, Sirivichayakul C, Dondorp AM, Chokejindachai W. Severe vivax malaria: a systematic review and meta-analysis of clinical studies since 1900. *Malar J* 2014; **13**: 481.
- 52 Anstey NM, Douglas NM, Poespoprodjo JR, Price RN. *Plasmodium vivax*: clinical spectrum, risk factors and pathogenesis. *Adv Parasitol* 2012; **80**: 151–201.
- 53 Lampah DA, Yeo TW, Hardianto SO, et al. Coma associated with microscopy-diagnosed *Plasmodium vivax*: a prospective study in Papua, Indonesia. *PLoS Negl Trop Dis* 2011; **5**: e1032.
- 54 Douglas NM, Pontororing GJ, Lampah DA, et al. Mortality attributable to *Plasmodium vivax* malaria: a clinical audit from Papua, Indonesia. *BMC Med* 2014; **12**: 217.
- 55 Barber BE, Grigg MJ, William T, Yeo TW, Anstey NM. The treatment of *Plasmodium knowlesi* malaria. *Trends Parasitol* 2017; **33**: 242–53.
- 56 World Health Organization. Guidelines for the treatment of malaria, 3rd edn. Geneva: WHO, 2015.
- 57 Price RN, von Seidlein L, Valecha N, Nosten F, Baird JK, White NJ. Global extent of chloroquine-resistant *Plasmodium vivax*: a systematic review and meta-analysis. *Lancet Infect Dis* 2014; **14**: 982–91.
- 58 WHO. World malaria report 2016. Geneva: World Health Organization, 2016.
- 59 Dellicour S, Seveane E, McGready R. First-trimester artemisinin derivatives and quinine treatments and the risk of adverse pregnancy outcomes in Africa and Asia: a meta-analysis of observational studies. *PLoS Med* 2017; **14**: e1002290.
- 60 Pekyi D, Ampromfi AA, Tinto H, et al. Four artemisinin-based treatments in African pregnant women with malaria. *N Engl J Med* 2016; **374**: 913–27.
- 61 Wilby KJ, Ensom MH. Pharmacokinetics of antimalarials in pregnancy: a systematic review. *Clin Pharmacokinet* 2011; **50**: 705–23.
- 62 Chu CS, White NJ. Management of relapsing *Plasmodium vivax* malaria. *Expert Rev Anti-infect Ther* 2016; **14**: 885–900.
- 63 Chu CS, Bancone G, Moore KA, et al. Haemolysis in G6PD heterozygous females treated with primaquine for *Plasmodium vivax* malaria: a nested cohort in a trial of radical curative regimens. *PLoS Med* 2017; **14**: e1002224.
- 64 Marcsisin SR, Reichard G, Pybus BS. Primaquine pharmacology in the context of CYP 2D6 pharmacogenomics: current state of the art. *Pharmacol Ther* 2016; **161**: 1–10.
- 65 Llanos-Cuentas A, Lacerda MV, Rueangweerayut R, et al. Tafenoquine plus chloroquine for the treatment and relapse prevention of *Plasmodium vivax* malaria (DETECTIVE): a multicentre, double-blind, randomised, phase 2b dose-selection study. *Lancet* 2014; **383**: 1049–58.
- 66 Taylor WR, White NJ. Antimalarial drug toxicity: a review. *Drug Saf* 2004; **27**: 25–61.
- 67 White NJ. Why is it that antimalarial drug treatments do not always work? *Ann Trop Med Parasitol* 1998; **92**: 449–58.
- 68 WorldWide Antimalarial Resistance Network Lumefantrine PK/PD Study Group. Artemether-lumefantrine treatment of uncomplicated *Plasmodium falciparum* malaria: a systematic review and meta-analysis of day 7 lumefantrine concentrations and therapeutic response using individual patient data. *BMC Med* 2015; **13**: 227.
- 69 WorldWide Antimalarial Resistance Network DP Study Group. The effect of dosing regimens on the antimalarial efficacy of dihydroartemisinin-piperaquine: a pooled analysis of individual patient data. *PLoS Med* 2013; **10**: e1001564.
- 70 Adjui MA, Allan R, Anvikar AR, et al. The effect of dosing strategies on the therapeutic efficacy of artesunate-amodiaquine for uncomplicated malaria: a meta-analysis of individual patient data. *BMC Med* 2015; **13**: 66.
- 71 Nayyar GM, Breman JG, Newton PN, Herrington J. Poor-quality antimalarial drugs in southeast Asia and sub-Saharan Africa. *Lancet Infect Dis* 2012; **12**: 488–96.
- 72 Douglas NM, Nosten F, Ashley EA, et al. *Plasmodium vivax* recurrence following falciparum and mixed species malaria: risk factors and effect of antimalarial kinetics. *Clin Infect Dis* 2011; **52**: 612–20.
- 73 White N. Antimalarial drug resistance and combination chemotherapy. *Philos Trans R Soc Lond B, Biol Sci* 1999; **354**: 739–49.
- 74 Dondorp AM, Nosten F, Yi P, et al. Artemisinin resistance in *Plasmodium falciparum* malaria. *N Engl J Med* 2009; **361**: 455–67.
- 75 Ashley EA, Dhorda M, Fairhurst RM, et al. Spread of artemisinin resistance in *Plasmodium falciparum* malaria. *N Engl J Med* 2014; **371**: 411–23.
- 76 Menard D, Khim N, Beghain J, et al. A worldwide map of *Plasmodium falciparum* K13-propeller polymorphisms. *N Engl J Med* 2016; **374**: 2453–64.
- 77 Amato R, Lim P, Miotto O, et al. Genetic markers associated with dihydroartemisinin-piperaquine failure in *Plasmodium falciparum* malaria in Cambodia: a genotype-phenotype association study. *Lancet Infect Dis* 2017; **17**: 164–73.
- 78 Witkowski B, Duru V, Khim N, et al. A surrogate marker of piperaquine-resistant *Plasmodium falciparum* malaria: a phenotype-genotype association study. *Lancet Infect Dis* 2017; **17**: 174–83.
- 79 Arie F, Witkowski B, Amaratunga C, et al. A molecular marker of artemisinin-resistant *Plasmodium falciparum* malaria. *Nature* 2014; **505**: 50–55.
- 80 Blasco B, Leroy D, Fidock DA. Antimalarial drug resistance: linking *Plasmodium falciparum* parasite biology to the clinic. *Nature Med* 2017; **23**: 917–28.
- 81 Phyo AP, Ashley EA, Anderson TJ, et al. Declining efficacy of artemisinin combination therapy against *P. falciparum* malaria on the Thai–Myanmar Border (2003–2013): the role of parasite genetic factors. *Clin Infect Dis* 2016; **63**: 784–91.
- 82 Thanh NV, Thuy-Nhien N, Tuyen NT, et al. Rapid decline in the susceptibility of *Plasmodium falciparum* to dihydroartemisinin-piperaquine in the south of Vietnam. *Malar J* 2017; **16**: 27.
- 83 Imwong M, Hien TT, Thuy-Nhien NT, Dondorp AM, White NJ. Spread of a single multidrug resistant malaria parasite lineage (PfPR10) to Vietnam. *Lancet Infect Dis* 2017; **17**: 1022–23.
- 84 WorldWide Antimalarial Resistance Network. Tracking resistance to artemisinin collaboration II. 2017. <http://www.wwarn.org/working-together/partner-projects/tracking-resistance-artemisinin-collaboration-ii> (accessed March 21, 2018).
- 85 White NJ, Pukrittayakamee S, Phyo AP, et al. Spiroindolone KAE609 for falciparum and vivax malaria. *N Engl J Med* 2014; **371**: 403–10.



- 86 McCarthy JS, Lotharius J, Ruckle T, et al. Safety, tolerability, pharmacokinetics, and activity of the novel long-acting antimalarial DSM265: a two-part first-in-human phase 1a/1b randomised study. *Lancet Infect Dis* 2017; **17**: 626–35.
- 87 White NJ, Duong TT, Uthaisin C, et al. Antimalarial activity of KAF156 in falciparum and vivax malaria. *N Engl J Med* 2016; **375**: 1152–60.
- 88 Phyto AP, Jittamala P, Nosten FH, et al. Antimalarial activity of artefenomel (OZ439), a novel synthetic antimalarial endoperoxide, in patients with *Plasmodium falciparum* and *Plasmodium vivax* malaria: an open-label phase 2 trial. *Lancet Infect Dis* 2016; **16**: 61–69.
- 89 Wells TN, Hooft van Huijsdijnen R, Van Voorhis WC. Malaria medicines: a glass half full? *Nature Rev Drug Discov* 2015; **14**: 424–42.
- 90 Phillips MA, Burrows JN, Manyando C, van Huijsdijnen RH, Van Voorhis WC, Wells TNC. Malaria. *Nature Rev Dis Primers* 2017; **3**: 17050.
- 91 Toure OA, Valecha N, Tshefu AK, et al. A phase 3, double-blind, randomized study of arterolane maleate-piperaquine phosphate vs artemether-lumefantrine for falciparum malaria in adolescent and adult patients in Asia and Africa. *Clin Infect Dis* 2016; **62**: 964–71.
- 92 Delves MJ, Ruecker A, Straschil U, et al. Male and female *Plasmodium falciparum* mature gametocytes show different responses to antimalarial drugs. *Antimicrob Agents Chemother* 2013; **57**: 3268–74.
- 93 Zoller T, Junghanss T, Kapaun A, et al. Intravenous artesunate for severe malaria in travelers, Europe. *Emerg Infect Dis* 2011; **17**: 771–77.
- 94 Jauréguiberry S, Ndour PA, Roussel C, et al. French Artesunate Working Group. Postartesunate delayed hemolysis is a predictable event related to the lifesaving effect of artemisinins. *Blood* 2014; **124**: 167–75.
- 95 Fanello C, Onyamboko M, Lee SJ, et al. Post-treatment haemolysis in African children with hyperparasitaemic falciparum malaria; a randomized comparison of artesunate and quinine. *BMC Infect Dis* 2017; **17**: 575.
- 96 Rolling T, Agbenyega T, Krishna S, Kreamsner PG, Cramer JP. Delayed haemolysis after artesunate treatment of severe malaria—review of the literature and perspective. *Trav Med Infect Dis* 2015; **13**: 143–49.
- 97 Olupot-Olupot P, Engoru C, Uyoga S, et al. High frequency of blackwater fever among children presenting to hospital with severe febrile illnesses in eastern Uganda. *Clin Infect Dis* 2017; **64**: 939–46.
- 98 Leoni S, Buonfrate D, Angheben A, Gobbi F, Bisoffi Z. The hyper-reactive malarial splenomegaly: a systematic review of the literature. *Malar J* 2015; **14**: 185.
- 99 Langford S, Douglas NM, Lampah DA, et al. *Plasmodium malariae* infection associated with a high burden of anemia: a hospital-based surveillance study. *PLoS Negl Trop Dis* 2015; **9**: e0004195.
- 100 Saadani Hassani A, Marston BJ. Impact of cotrimoxazole and insecticide-treated nets for malaria prevention on key outcomes among HIV-infected adults in low- and middle-income countries: a systematic review. *J Acquire Immune Defic Syndr* (1999) 2015; **68** (suppl 3): S306–17.
- 101 Hoglund RM, Byakika-Kibwika P, Lamorde M, et al. Artemether-lumefantrine co-administration with antiretrovirals: population pharmacokinetics and dosing implications. *Br J Clin Pharmacol* 2015; **79**: 636–49.
- 102 Kakuru A, Achan J, Muhindo MK, et al. Artemisinin-based combination therapies are efficacious and safe for treatment of uncomplicated malaria in HIV-infected Ugandan children. *Clin Infect Dis* 2014; **59**: 446–53.
- 103 Galatas B, Bassat Q, Mayor A. Malaria parasites in the asymptomatic: looking for the hay in the haystack. *Trends Parasitol* 2016; **32**: 296–308.
- 104 Boyle MJ, Reiling L, Feng G, et al. Human antibodies fix complement to inhibit *Plasmodium falciparum* invasion of erythrocytes and are associated with protection against malaria. *Immunity* 2015; **42**: 580–90.
- 105 Van Braeckel-Budimir N, Kurup SP, Harty JT. Regulatory issues in immunity to liver and blood-stage malaria. *Curr Opin Immunol* 2016; **42**: 91–97.
- 106 Taylor SM, Parobek CM, Fairhurst RM. Haemoglobinopathies and the clinical epidemiology of malaria: a systematic review and meta-analysis. *Lancet Infect Dis* 2012; **12**: 457–68.
- 107 Malaria Genomic Epidemiology Network. Reappraisal of known malaria resistance loci in a large multicenter study. *Nat Genet* 2014; **46**: 1197–204.
- 108 Leslie T, Briceno M, Mayan I, et al. The impact of phenotypic and genotypic G6PD deficiency on risk of *Plasmodium vivax* infection: a case-control study amongst Afghan refugees in Pakistan. *PLoS Med* 2010; **7**: e1000283.
- 109 Louicharoen C, Patin E, Paul R, et al. Positively selected G6PD-Mahidol mutation reduces *Plasmodium vivax* density in southeast Asians. *Science* 2009; **326**: 1546–49.
- 110 Batchelor JD, Malpede BM, Omattage NS, DeKoster GT, Henzler-Wildman KA, Tolia NH. Red blood cell invasion by *Plasmodium vivax*: structural basis for DBP engagement of DARC. *PLoS Pathog* 2014; **10**: e1003869.
- 111 Niangaly A, Karthigayan G, Amed O, et al. *Plasmodium vivax* infections over 3 years in Duffy blood group negative Malians in Bandiagara, Mali. *Am J Trop Med Hyg* 2017; **97**: 744–52.
- 112 Leffler EM, Band G, Busby GB, et al. Resistance to malaria through structural variation of red blood cell invasion receptors. *Science* 2017; **356**: 1140–152.
- 113 Nevill CG, Some ES, Mung'ala VO, et al. Insecticide-treated bednets reduce mortality and severe morbidity from malaria among children on the Kenyan coast. *Trop Med Int Health* 1996; **1**: 139–46.
- 114 Radeva-Petrova D, Kayentao K, ter Kuile FO, Sinclair D, Garner P. Drugs for preventing malaria in pregnant women in endemic areas: any drug regimen versus placebo or no treatment. *Cochrane Database Syst Rev* 2014; **10**: CD000169.
- 115 Desai M, Gutman J, L'Lanziva A, et al. Intermittent screening and treatment or intermittent preventive treatment with dihydroartemisinin-piperaquine versus intermittent preventive treatment with sulfadoxine-pyrimethamine for the control of malaria during pregnancy in western Kenya: an open-label, three-group, randomised controlled superiority trial. *Lancet* 2015; **386**: 2507–19.
- 116 Madanitsa M, Kalilani L, Mwapasa V, et al. Scheduled intermittent screening with rapid diagnostic tests and treatment with dihydroartemisinin-piperaquine versus intermittent preventive therapy with sulfadoxine-pyrimethamine for malaria in pregnancy in Malawi: an open-label randomised controlled trial. *PLoS Med* 2016; **13**: e1002124.
- 117 Cisse B, Ba EH, Sokhna C, et al. Effectiveness of seasonal malaria chemoprevention in children under ten years of age in Senegal: a stepped-wedge cluster-randomised trial. *PLoS Med* 2016; **13**: e1002175.
- 118 Kolifarhood G, Raeisi A, Ranjbar M, et al. Prophylactic efficacy of primaquine for preventing *Plasmodium falciparum* and *Plasmodium vivax* parasitaemia in travelers: a meta-analysis and systematic review. *Trav Med Infect Dis* 2017; **17**: 5–18.
- 119 RTS,S Clinical Trials Partnership. Efficacy and safety of RTS,S/AS01 malaria vaccine with or without a booster dose in infants and children in Africa: final results of a phase 3, individually randomised, controlled trial. *Lancet* 2015; **386**: 31–45.
- 120 Neafsey DE, Juraska M, Bedford T, et al. Genetic diversity and protective efficacy of the RTS,S/AS01 malaria vaccine. *N Engl J Med* 2015; **373**: 2025–37.
- 121 Olotu A, Fegan G, Wambua J, et al. Seven-year efficacy of RTS,S/AS01 malaria vaccine among young African children. *Engl J Med* 2016; **374**: 2519–29.
- 122 Sissoko MS, Healy SA, Katile A, et al. Safety and efficacy of PfSPZ vaccine against *Plasmodium falciparum* via direct venous inoculation in healthy malaria-exposed adults in Mali: a randomised, double-blind phase 1 trial. *Lancet Infect Dis* 2017; **17**: 498–509.
- 123 Cowman AF, Tonkin CJ, Tham WH, Duraisingh MT. The molecular basis of erythrocyte invasion by malaria parasites. *Cell Host Microbe* 2017; **22**: 232–45.
- 124 Crosnier C, Bustamante LY, Bartholdson SJ, et al. Basigin is a receptor essential for erythrocyte invasion by *Plasmodium falciparum*. *Nature* 2011; **480**: 534–37.
- 125 Fried M, Duffy PE. Designing a VAR2CSA-based vaccine to prevent placental malaria. *Vaccine* 2015; **33**: 7483–88.
- 126 Bustamante LY, Powell GT, Lin YC, et al. Synergistic malaria vaccine combinations identified by systematic antigen screening. *Proc Natl Acad Sci USA* 2017; **114**: 12045–50.
- 127 WHO. Global plan for insecticide resistance management in malaria vectors (GPIRM). Geneva: World Health Organization, 2013.
- 128 Ranson H, Lissenden N. Insecticide resistance in African anophelous mosquitoes: a worsening situation that needs urgent action to maintain malaria control. *Trends Parasitol* 2016; **32**: 187–96.



- 129 Strode C, Donegan S, Garner P, Enayati AA, Hemingway J. The impact of pyrethroid resistance on the efficacy of insecticide-treated bed nets against African anopheline mosquitoes: systematic review and meta-analysis. *PLoS Med* 2014; **11**: e1001619.
- 130 World Health Organization Global Malaria Programme. Conditions for deployment of mosquito nets treated with pyrethroid and piperonyl butoxide. September 2017 (updated December 2017). <http://www.who.int/malaria/publications/atoz/use-of-pbo-treated-lins/en/> (accessed March 21, 2018).
- 131 Wilson AL, Chen-Hussey V, Logan JG, Lindsay SW. Are topical insect repellents effective against malaria in endemic populations? A systematic review and meta-analysis. *Malar J* 2014; **13**: 446.
- 132 Sluydts V, Durnez L, Heng S, et al. Efficacy of topical mosquito repellent (picaridin) plus long-lasting insecticidal nets versus long-lasting insecticidal nets alone for control of malaria: a cluster randomised controlled trial. *Lancet Infect Dis* 2016; **16**: 1169–77.
- 133 Alout H, Foy BD. Ivermectin: a complimentary weapon against the spread of malaria? *Expert Rev Anti Infect Ther* 2017; **15**: 231–40.
- 134 McLean KJ, Jacobs-Lorena M. Genetic control of malaria mosquitoes. *Trends Parasitol* 2016; **32**: 174–76.
- 135 Marshall JM, Taylor CE. Malaria control with transgenic mosquitoes. *PLoS Med* 2009; **6**: e20.
- 136 Gantz VM, Jasinskiene N, Tatarenkova O, et al. Highly efficient Cas9-mediated gene drive for population modification of the malaria vector mosquito *Anopheles stephensi*. *Proc Natl Acad Sci USA* 2015; **112**: e6736–43.
- 137 Newby G, Bennett A, Larson E, et al. The path to eradication: a progress report on the malaria-eliminating countries. *Lancet* 2016; **387**: 1775–84.
- 138 Landier J, Kajeechiwa L, Thwin MM, et al. Safety and effectiveness of mass drug administration to accelerate elimination of artemisinin-resistant falciparum malaria: a pilot trial in four villages of eastern Myanmar. *Wellcome Open Res* 2017; **2**: 81.
- 139 Imwong M, Suwannasin K, Kunasol C, et al. The spread of artemisinin-resistant *Plasmodium falciparum* in the Greater Mekong subregion: a molecular epidemiology observational study. *Lancet Infect Dis* 2017; **17**: 491–97.
- 140 Bennett A, Bisanzio D, Yukich JO, et al. Population coverage of artemisinin-based combination treatment in children younger than 5 years with fever and *Plasmodium falciparum* infection in Africa, 2003–2015: a modelling study using data from national surveys. *Lancet Glob Health* 2017; **5**: e418–e27.

© 2018 Elsevier Ltd. All rights reserved.