

parameters:

- **Normal Distribution:**

⇒ Mean (μ) = center

⇒ Standard deviation (σ) = width

- **cauchy Distribution: / standard**

⇒ μ (Mu) = center

⇒ γ (gamma) = width

- **Bivariate distribution:**

⇒ μ_1 and μ_2 (Means of the two variables)

⇒ σ_1 and σ_2 (Standard deviations of the two variables)

⇒ ρ (rho, the correlation coefficient b/w the two variables).

- **t-Distributions:**

⇒ ν (nu) = Degrees of freedom

⇒ μ (mu) = center of the distribution

⇒ σ (sigma) = width of the distribution.

- **Lognormal Distribution:**

⇒ μ (mu) = Mean of the logarithm of the variable.

⇒ σ (sigma) = Standard deviation of the logarithm of the variable.

• Exponential distribution:

⇒ λ (lambda)

The Exponential distribution can also be expressed by:

⇒ β (beta) = which is the inverse of λ .

⇒ μ (mu) = Mean (which is equal to $1/\lambda$)

Note: λ (lambda) is the most commonly used parameter to describe the exponential distribution.

• Gamma distribution:

⇒ α (alpha) = Shape parameter.

⇒ β (beta) = Rate parameter.

Note:

the gamma distribution can also be described by:

⇒ α (alpha)

⇒ θ (theta) = which is inverse of β .

• Beta distribution:

⇒ α (alpha)

⇒ β (beta)

Note: Both α and β must be positive.

- **Weibull distribution:**

⇒ K (Kappa)

⇒ λ (lambda)

Note: Both K and λ must be positive values.

- **Chi-Square Distribution:**

⇒ K (degrees of freedom)

The chi-Square distribution has only one parameter, which is the number of degrees of freedom (K).

Note: K must be a positive integer.

- **F-Distribution:**

⇒ d_1 (df 1)

⇒ d_2 (df 2)

Note: The F-distribution has two parameters, both of which are positive integers.

- **Pareto Distribution:**

⇒ X_m (X minimum) = The smallest time value.

⇒ α (alpha)

The pareto distribution can also be described by

⇒ K = (related to X_m and α)

⇒ α = (alpha)

DATE: _____

DAY: _____

Order statistics

Order statistics don't have any parameters.

