10.3.1. Tabelle 3: Charakterisierung der Lebensqualitätsstudien unter Therapie mit Mistelextrakten

Wilfried Tröger

Lokalisation/Art Stadium	Lebens-qualitäts-erfassung	Intervention Verum	Kontrolle	Ergebnis SMD [95 % KI]	N Teilnehmer; N Frauen; Alter Verum	N Teilnehmer; N Frauen; Alter Kontrolle	multi-zentrisch	rando-misiert	offen	interven-ierend	Land	Referenz
Osteosarkom IB–IIIB	EORTC QLQ-C30	Iscador® P	Etoposid (oral)	0,68 [-0,23 bis 1,59]	9; 5; 28,0	11; 4; 39,0	nein	ja	ja	ja	Italien	Longhi 2014 (4)
Mamma Tx, T1–3	EORTC QLQ-C30	Chemotherapie, Helixor® A	Chemotherapie	0,59 [0,08 bis 1,11]	34; 34; 50,4	31; 31; 50,8	nein	ja	ja	ja	Serbien	Tröger 2014a (70)
Pankreas T3–4	EORTC QLQ-C30	Supportivtherapie, Iscador®	Supportivtherapie	2,34 [1,94 bis 2,73]	110; 45; 62,2	110; 47; 64,4	nein	ja	ja	ja	Serbien	Tröger 2014b (53)
Lunge (NSCLC), IIIa–IV	EORTC QLQ-C30	Chemotherapie, Iscador®	Chemotherapie	-0,17 [-0,70 bis 0,36]	33; 5; 63,0	39; 11; 62,0	nein	ja	ja	ja	Israel	Bar-Sela 2013 (5)
Gastrointestinal Ib, II	EORTC QLQ-C30	Chemotherapie, Abnobaviscum®	Chemotherapie	0,25 [-0,48 bis 0,98]	16; 3; 53,8	16; 3; 54,9	nein	ja	ja	ja	Südkorea	Kim 2012 (75)
Lunge (NSCLC), IIIb, IV	EORTC QLQ-C30	Chemotherapie, Iscador®	Chemotherapie	-0,50 [-1,07 bis 0,07]	26; 7; 64,3	24; 7; 63,0	nein	ja	ja	ja	Deutsch-land	Grah 2010 (76)
Mamma Tx, T1–3	EORTC QLQ-C30	Chemotherapie, Iscador® M	Chemotherapie	0,29 [-0,22 bis 0,79]	30; 30; 48,4	31; 31; 50,8	nein	ja	ja	ja	Serbien	Tröger 2009 (69)
Corpus uteri IA-C, IIIa–IVb	Selbstregulation	konventionelle Therapie, Iscador®	konventionelle Therapie	0,71 [0,43 bis 0,99]	105; 105; 58,5	105; 105; 59,1	ja	nein	ja	nein	Deutsch-land	Grossarth 2008 (9)
Corpus uteri IA-C, IIIa–IVb	Selbstregulation	konventionelle Therapie, Iscador®	konventionelle Therapie	0,80 [0,27 bis 1,33]	31; 31; 55,1	31; 31; 55,1	ja	ja	ja	ja	Deutsch-land	Grossarth 2008 (9)
Mamma TIa–c, TII	EORTC QLQ-C30	Chemotherapie, Iscador® M	Chemotherapie	-0,13 [-0,61 bis 0,36]	39; 39; 47,5	43; 43; 47,5	nein	nein	ja	ja	Deutsch-land	Loewe-Mesch 2008 (63)
Cervix uteri IB–IVa	Selbstregulation	konventionelle Therapie, Iscador®	konventionelle Therapie	0,68 [0,40 bis 0,97]	102; 102; 51,0	102; 102; 51,0	ja	nein	ja	nein	Deutsch-land	Grossarth 2007a (12)
Cervix uteri IB–IVa	Selbstregulation	konventionelle Therapie, Iscador®	konventionelle Therapie	0,79 [0,13 bis 1,46]	19; 19; 47,7	9; 19; 47,6	ja	ja	ja	ja	Deutsch-land	Grossarth 2007a (12)
Ovarial IA–C, IV	Selbstregulation	konventionelle Therapie, Iscador®	konventionelle Therapie	0,48 [0,15 bis 0,81]	75; 75; 43,0	75; 75; 45,1	ja	nein	ja	nein	Deutsch-land	Grossarth 2007b (10)
Ovarial IA–C, IV	Selbstregulation	konventionelle Therapie, Iscador®	konventionelle Therapie	1,23 [0,56 bis 1,89]	21; 21; 45,4	21; 21; 45,5	ja		ja	ja	Deutsch-land	Grossarth 2007b (10)
Melanom keine Angabe	Selbstregulation	konventionelle Therapie, Iscador®	konventionelle Therapie	0,80 [0,18 bis 1,41]	22; 8; 52,0	22; 8; 52,0	ja	ja	ja	ja	Deutsch-land	Grossarth 2007c (11)
Mamma T1a–T3	Selbstregulation	konventionelle Therapie, Iscador®	konventionelle Therapie	0,52 [0,06 bis 0,98]	97; 97; 52,2	97; 97; 52,2	ja	nein	ja	nein	Deutsch-land	Grossarth 2006a (13)
Mamma T1a–T3	Selbstregulation	konventionelle Therapie, Iscador®	konventionelle Therapie	0,30 [0,00 bis 0,61]	39; 39; 52,8	39; 39; 52,9	ja	ja	ja	ja	Deutschl and	Grossarth 2006a (13)
Mamma T2–4	Selbstregulation	konventionelle Therapie, Iscador®	konventionelle Therapie	0,66 [-0,04 bis 1,35]	17; 17; 44,5	17; 17; 44,6	ja	ja	ja	ja	Deutsch-land	Grossarth 2006b (14)
Mamma pTis–pT3	FACT-G, GLQ-8, Spitzer Uniscale	Chemotherapie, Lektinol®	Chemotherapie, Placebo	0,82 [0,60 bis 1,05]	176; 176; 46,4	176; 176; 45,9	ja	ja	ja	ja	BGR, RUS, UKR	Semiglazov 2006 (68)
Kopf/Hals Tumor, I–IV	EORTC QLQ-C30	Operation, Eurixor®	Operation	0,77 [0,28 bis 1,27]	200; 17; 55,0	199; 14; 55,0	ja	ja	ja	ja	Deutsch-land	Steuer-Vogt 2006 (77)
Gastrointestinal keine Angabe	Karnofsky-Index	Operation, Isorel® A	Operation	0,57 [0,33 bis 0,82]	40; kA; kA	30; kA; kA	nein	ja	ja	ja	Rumä-nien	Enesel 2005 (78)
verschiedene pTx, pT1–4	TCM, FLIC, Kar-nofsky Index	Chemotherapie, Helixor®	Chemotherapie, Lentinan (i.m.)	0,25 [-0,01 bis 0,52]	118; 91; 52,6	115; 91; 51,7	ja	ja	ja	ja	China	Piao 2004 (66)
Mamma pT1–3	GLQ-8, Spitzer Uniscale	Chemotherapie, Lektinol®	Chemotherapie, Placebo	0,50 [0,16 bis 0,85]	67; 67; 44,6	70; 70; 43,5	ja	ja	nein	ja	BGR, RUS, UKR	Semiglasov 2004 (67)
Mamma IV	Spitzer QoL	konventionelle Therapie danach Mistelextrakt	konventionelle Therapie, danach isot. Salzlösung	0,40 [-0,37 bis 1,16]	20; 20; kA	10; 10; kA	nein	ja	ja	ja	Italien	Borrelli 2001 (58)
verschiedene kA	ECOG	kA, Abnobaviscum®	kA, isot. Salzlösung	1,17 [0,29 bis 2,05]	29; 15; kA	29; 15; kA	nein	ja	nein	ja	Deutsch-land	Kaiser 2001 (79)
Kolorektal fortgeschritten	FACT-G	Chemotherapie, Eurixor®	Chemotherapie	2,17 [1,61 bis 2,73]	38; 16; 54,7	41; 18; 53,2	nein	ja	ja	ja	Deutschl and	Heiny 1998 (80)
Gliom III–IV	Spitzer QoL	konventionelle Therapie, Mistelextrakt	konventionelle Therapie	0,26 [-0,51 bis 1,03]	18; 6; 52,0	17; 7; 52,0	nein	ja	ja	ja	Deutsch-land	Lenartz 1996 (81)
Bronchial (NSCLC) I–IV	Karnofsky-Index	konventionelle Therapie, Iscador®	konventionelle Therapie, BVK Roche®	-0,09 [-0,39 bis 0,21]	114; 6; 65,6	113; 6; 67,5	ja	ja	ja	ja	Deutsch-land	Dold 1991 (20 S. 280)
Mamma fortgeschritten	Auszug aus FLIC	Chemotherapie, Eurixor®	Chemotherapie, isot. Salzlösung	0,83 [0,18 bis 1,48]	25; 25; kA	21; 21; kA	nein	ja	ja	ja	Deutsch-land	Heiny 1991 (60)
verschiedene keine Angabe	Karnofsky-Index	konventionelle Therapie, Helixor® A	konventionelle Therapie	0,44 [-0,16 bis 1,04]	35; 9; 58,3	33; 9; 60,2	nein	ja	ja	ja	Deutsch-land	Lange 1983 (48)

BGR = Bulgarien, BVK = B-Vitamin-Komplex, ECOG = Eastern Cooperative Oncology Group Score, EORTC QLQ-C30 = European Organisation for Research and Treatment of Cancer Core Quality of Life Questionnaire, FACT-G = Functional Assessment of Cancer Therapy – General, FLIC = Functional Living Index Cancer, GLQ-8 = Global Life Quality Scale, kA = keine Angabe, QoL = Quality of Life Index, RUS = Russland, SMD = standardisierte Mittelwertdifferenz, TCM = Traditional Chinese Medicine Score, UKR = Ukraine.

Referenzen: siehe Folgeseite

Referenzen

1. Ostermann T, Appelbaum S, Poier D, Boehm K, Raak C, Büssing A. A Systematic Review and Meta-Analysis on the Survival of Cancer Patients Treated with a Fermented Viscum album L. Extract (Iscador): An Update of Findings. Complement Med Res 2020; 27(4): 260–271. ▸ https://doi.org/10.1159/000505202.
2. Loef M, Walach H. Survival of Cancer Patients Treated with Non-Fermented Mistletoe Extract: A Systematic Review and Meta-Analysis. Integr Cancer Ther 2022; 21: 15347354221133561. ▸ https://doi.org/10.1177/15347354221133561.
3. Loef M, Walach H. Quality of life in cancer patients treated with mistletoe: a systematic review and meta-analysis. BMC Complement Med Ther 2020; 20(1): 227. ▸ https://doi.org/10.1186/s12906-020-03013-3.
4. Longhi A, Reif M, Mariani E, Ferrari S. A Randomized Study on Postrelapse Disease-Free Survival with Adjuvant Mistletoe versus Oral Etoposide in Osteosarcoma Patients. Evid Based Complement Alternat Med 2014; 2014: 210198. ▸ https://doi.org/10.1155/2014/210198.
5. Bar-Sela G, Wollner M, Hammer L, Agbarya A, Dudnik E, Haim N. Mistletoe as complementary treatment in patients with advanced non-small-cell lung cancer treated with carboplatin-based combinations: A randomised phase II study. Eur J Cancer 2013; 49(5): 1058–1064. ▸ https://doi.org/10.1016/j.ejca.2012.11.007.
6. Tröger W, Galun D, Reif M, Schumann A, Stanković N, Milićević M. Viscum album [L.] extract therapy in patients with locally advanced or metastatic pancreatic cancer: A randomised clinical trial on overall survival. Eur J Cancer 2013; 49(18): 3788–3797. ▸ https://doi.org/10.1016/j.ejca.2013.06.043.
7. Zänker KS, Matthes H, Bock PR, Hanisch J. A Specific Mistletoe Preparation (Iscador-Qu®) in Colorectal Cancer (CRC) Patients: More than Just Supportive Care? J Cancer Sci Ther 2012;4(9): 264–270. ▸ https://doi.org/10.4172/1948-5956.1000153.
8. Matthes H, Friedel WE, Bock PR, Zänker KS. Molecular mistletoe therapy: friend or foe in established anti-tumor protocols? A multicenter, controlled, retrospective pharmaco-epidemiological study in pancreas cancer. Curr Mol Med 2010; 10(4): 430–439. ▸ https://www.eurekaselect.com/article/16735.
9. Grossarth-Maticek R, Ziegler R. Randomized and non-randomized prospective controlled cohort studies in matched pair design for the long-term therapy of corpus uteri cancer patients with a mistletoe preparation (Iscador). Eur J Med Res 2008; 13(3): 107–120. ▸ https://pubmed.ncbi.nlm.nih.gov/18499556/.
10. Grossarth-Maticek R, Ziegler R. Prospective Controlled Cohort Studies on Long-term Therapy of Ovarian Cancer Patients with Mistletoe (Viscum album L.) Extracts Iscador. Arzneimittelforschung 2007; 57(10): 665–678. ▸ https://doi.org/10.1055/s-0031-1296666.
11. Grossarth-Maticek R, Ziegler R. Wirksamkeit und Unbedenklichkeit einer Langzeitbehandlung von Melanompatienten mit einem Mistelpräparat (Iscador). Schweiz Z Ganzheitsmed 2007; 19(6): 325–332. ▸ https://doi.org/10.1159/000285940.
12. Grossarth-Maticek R, Ziegler R. Prospective Controlled Cohort Studies on Long-Term Therapy of Cervical Cancer Patients with a Mistletoe Preparation (Iscador®). Forsch Komplementmed 2007; 14(3): 140–147. ▸ https://doi.org/10.1159/000102956.
13. Grossarth-Maticek R, Ziegler R. Prospective Controlled Cohort Studies on Long-Term Therapy of Breast Cancer Patients with a Mistletoe Preparation (Iscador®). Forsch Komplementmed 2006; 13(5): 285–292. ▸ https://doi.org/10.1159/000095378.
14. Grossarth-Maticek R, Ziegler R. Randomised and non-randomised prospective controlled cohort studies in matched-pair design for the long-term therapy of breast cancer patients with a mistletoe preparation (Iscador): a re-analysis. Eur J Med Res 2006; 11(11): 485–495. ▸ https://pubmed.ncbi.nlm.nih.gov/17182361/.
15. Augustin M, Bock PR, Hanisch J, Karasmann M, Schneider B. Safety and Efficacy of the Long-term Adjuvant Treatment of Primary Intermediate- to High-Risk Malignant Melanoma (UICC/AJCC Stage II and III) with a Standardized Fermented European Mistletoe (Viscum album L.) Extract. Arzneimittelforschung 2005; 55(1): 38–49. ▸ https://doi.org/10.1055/s-0031-1296823.
16. Bock PR, Friedel WE, Hanisch J, Karasmann M, Schneider B. Retrolective, Comparative, Epidemiological Cohort Study with Parallel Groups Design for Evaluation of Efficacy and Safety of Drugs with „Well-Established Use“: Experience with the Long-Term Treatment Using the European Mistletoe Extract (Viscum album L.) in Addition to Conventional Oncological Therapy in Primary, Non-Metastatic Breast Carcinoma. Forsch Komplementarmed Klass Naturheilkd 2004; 11(Suppl. 1): 23–29. ▸ https://doi.org/10.1159/000080572.
17. Grossarth-Maticek R, Kiene H, Baumgartner S, Ziegler R. Synergieeffekte von Selbstregulation und Misteltherapie (Iscador) auf die Überlebenszeit bei Krebspatienten. Ergebnisse einer epidemiologischen Langzeitstudie, Teil II. Schweiz Z Ganzheitsmed 2004; 16(2): 81–89. ▸ https://www.researchgate.net/publication/290326317_Synergy_effects_of_self-regulation_and_mistletoe_therapy_Iscador_on_survival_of_cancer_patients.
18. Kleeberg UR, Suciu S, Bröcker EB, Ruiter DJ, Chartier C, Liénard D, Marsden J, Schadendorf D, Eggermont AMM. Final results of the EORTC 18871/DKG 80-1 randomised phase III trial. rIFN-alpha2b versus rIFN-gamma versus Iscador M versus observation after surgery in melanoma patients with either high-risk primary (thickness >3 mm) or regional lymph node metastasis. Eur J Cancer 2004; 40(3): 390–402. ▸ https://doi.org/10.1016/j.ejca.2003.07.004.
19. Grossarth-Maticek R, Kiene H, Baumgartner SM, Ziegler R. Use of Iscador, an extract of European mistletoe (Viscum album L.), in cancer treatment: prospective nonrandomized and randomized matched-pair studies nested within a cohort study. Altern Ther Health Med 2001; 7(3): 57–78. ▸ https://pubmed.ncbi.nlm.nih.gov/11347286/.
20. Dold U, Edler L, Mäurer HC, Müller-Wening D, Sakellariou B, Trendelenburg F, Wagner G. Krebszusatztherapie beim fortgeschrittenen nicht-kleinzelligen Bronchialkarzinom. Multizentrische kontrollierte Studie zur Prüfung der Wirksamkeit von Iscador und Polyerga. Stuttgart, New York: Georg Thieme Verlag 1991.
21. Salzer G, Danmayr E, Wutzlhofer F, Frey S. Adjuvante Iscador-Behandlung operierter nicht kleinzelliger Bronchuskarzinome. Ergebnisse einer randomisierten Studie. Dtsch Z Onkol 1991; 23(4): 93–98.
22. Schuppli R. Die adjuvante Behandlung des malignen Melanoms mit Iscador c. Hg. In: Jungi WF, Senn H-J (Hrsg.). Krebs und Alternativmedizin II. 1. Aufl. Berlin, Heidelberg: Springer Verlag 1990.
23. Salzer G, Havelec L. Adjuvante Iscador-Behandlung nach operiertem Magenkarzinom. Ergebnisse einer randomisierten Studie. Krebsgeschehen 1983; 15(4): 106–110.
24. Hoffmann J, Hajto T. Iscadorbehandlung beim metastasierenden Mammakarzinom. Krebsgeschehen 1982; 14(3): 71–75.
25. Leroi R, Hajto T. Die Iscadortherapie beim Ovarialkarzinom. Krebsgeschehen 1982; 14(2): 38–44.
26. Salzer G, Havelec L. Rezidivprophylaxe bei operierten Bronchuskarzinompatienten mit dem Mistelpräparat Iscador®. Ergebnisse eines klinischen Versuchs aus den Jahren 1969–1971. Onkologie 1978; 1(6): 264–267. ▸ https://doi.org/10.1159/000213966.
27. Hoffmann J. Die Iscador-Behandlung bei Lebermetastasen. Krebsgeschehen 1979; 6: 172–175.
28. Leroi R. Die Iscadorbehandlung bei inoperablen kolo-rektalen Tumoren. Krebsgeschehen 1979; 11(6): 163–165.
29. Leroi R. Nachbehandlung des operierten Mammakarzinoms mit Viscum album. Helv Chir Acta 1977; 44(3): 403–414. ▸ https://pubmed.ncbi.nlm.nih.gov/617173/.
30. Leroi R. Malignomtherapie mit neuen Iscador-Präparaten. Krebsgeschehen 1975; 7(5): 124–126.
31. Günczler M, Salzer G. Iscadortherapie in der Nachbehandlung operierter Carcinome. Osterr Azteztg 1969; 24(20): 2290–2292.
32. Fellmer KE. A clinical trial of Iscador. Br Homoeopath J 1968; 57(1): 43–47. ▸ https://doi.org/10.1016/S0007-0785(68)80049-3.
33. Günczler M, Osika C, Salzer G. Ergebnisse von Resektion und Nachbehandlung beim Magenkarzinom. Wien Klin Wochenschr 1968; 80(6): 105–106. ▸ https://pubmed.ncbi.nlm.nih.gov/5661513/.
34. Majewski A, Bentele W. Über Zusatzbehandlung beim weiblichen Genitalkarzinom. Zentralbl Gynäkol 1963; 85(20): 696–700.
35. Thronicke A, Matthes B, von Trott P, Schad F, Grah C. Overall Survival of Non-metastasized NSCLC Patients Treated With Add-On Viscum album L: A Multicenter Real-World Study. Integr Cancer Ther 2020; 19: 1534735420940384. ▸ https://doi.org/10.1177/1534735420940384.
36. Schad F, Thronicke A, Steele ML, Merkle A, Matthes B, Grah C, Matthes H. Overall survival of stage IV non-small cell lung cancer patients treated with Viscum album L. in addition to chemotherapy, a real-world observational multicenter analysis. PLoS One 2018; 13(8): e0203058. ▸ https://doi.org/10.1371/journal.pone.0203058. Erratum in: PLoS One 2022; 17(8): e0273387. ▸ https://doi.org/10.1371/journal.pone.0273387.
37. Axtner J, Steele M, Kröz M, Spahn G, Matthes H, Schad F. Health services research of integrative oncology in palliative care of patients with advanced pancreatic cancer. BMC Cancer 2016; 16: 579. ▸ https://doi.org/10.1186/s12885-016-2594-5.
38. Tröger W, Ždrale Z, Stanković N. Fünf-Jahres-Nachbeobachtung von Patientinnen mit Brustkrebs nach einer randomisierten Studie mit Viscum album (L.) Extrakt. Dtsch Z Onkol 2016; 48(03): 105–110. ▸ https://doi.org/10.1055/s-0042-113505.
39. Beuth J, Schneider B, Schierholz JM. Impact of Complementary Treatment of Breast Cancer Patients with Standardized Mistletoe Extract during Aftercare: A Controlled Multicenter Comparative Epidemiological Cohort Study. Anticancer Res 2008; 28(1B): 523–527. ▸ https://ar.iiarjournals.org/content/28/1B/523.
40. Schumacher K, Schneider B, Reich G, Stiefel T, Stoll G, Bock PR, Hanisch J, Beuth J. Influence of postoperative complementary treatment with lectin-standardized mistletoe extract on breast cancer patients. A controlled epidemiological multicentric retrolective cohort study. Anticancer Res 2003; 23(6D): 5081–5087. ▸ https://pubmed.ncbi.nlm.nih.gov/14981970/.
41. Cazacu M, Oniu T, Lungoci C, Mihailov A, Cipak A, Klinger R, Weiss T, Zarković N. The Influence of Isorel on the Advanced Colorectal Cancer. Cancer Biother Radiopharm 2003; 18(1): 27–34. ▸ https://doi.org/10.1089/108497803321269304.
42. Goebell PJ, Otto T, Suhr J, Rübben H. Evaluation of an Unconventional Treatment Modality with Mistletoe Lectin to Prevent Recurrence of Superficial Bladder Cancer: A Randomized Phase II Trial. J Urol 2002; 168(1): 72–75. ▸ https://doi.org/10.1016/S0022-5347(05)64834-3.
43. Steuer-Vogt MK, Bonkowsky V, Ambrosch P, Scholz M, Neiß A, Strutz J, Hennig M, Lenarz T, Arnold W. The effect of an adjuvant mistletoe treatment programme in resected head and neck cancer patients: a randomised controlled clinical trial. Eur J Cancer 2001; 37(1): 23–31. ▸ https://doi.org/10.1016/s0959-8049(00)00360-9.
44. Lenartz D, Dott U, Menzel J, Schierholz JM, Beuth J. Survival of glioma patients after complementary treatment with galactoside-specific lectin from mistletoe. Anticancer Res 2000; 20(3B): 2073–2076. ▸ https://pubmed.ncbi.nlm.nih.gov/10928154/.

45. Heiny BM, Albrecht V. Komplementäre Therapie mit Mistellektin-1-normiertem Extrakt; Lebensqualitätsstabilisierung beim fortgeschrittenen kolorektalen Karzinom. Med Welt 1997; 48(9): 419–423.
46. Douwes FR, Kalden M, Frank G, Holzhauer P. Behandlung des fortgeschrittenen kolorektalen Karzinoms. Dtsch Z Onkol 1988; 20(3): 63–67.
47. Gutsch J, Berger H, Scholz G, Denck H. Prospektive Studie beim radikal operierten Mammakarzinom mit Polychemotherapie, Helixor und unbehandelter Kontrolle. Dtsch Z Onkol 1988; 4: 94–101.
48. Lange O, Scholz G, Gutsch J. Modulation der subjektiven und objektiven Toxizität einer aggressiven Chemo/ Radiotherapie mit Helixor. Unveröffentlichter Studienbericht 1983.
49. Douwes FR, Wolfrum DI, Migeod F. Ergebnisse einer prospektiv randomisierten Studie: Chemotherapie versus Chemotherapie plus "Biological response modifier" bei metastasierendem kolorektalem Karzinom. Krebsgeschehen 1986; 18(6): 155–164.
50. Boie D, Gutsch J. Helixor bei Kolon- und Rektumkarzinom. In: Denk H, Karrer K (Hrsg.). Schriftenreihe Krebsgeschehen. Band 23. 1. Aufl. Heidelberg: Verlag für Medizin Dr. Ewald Fischer 1980.
51. Prediger F. Über Ergebnisse der Behandlung inoperabler weiblicher Genitalkarzinome mit Plenosol in einem Zeitraum von 5 Jahren. Medizinische 1956; 42: 1513–1515.
52. U.S. Department of Health and Human Services, FDA, Oncology Center of Excellence, CDER, CBER (Hrsg.). Clinical Trial Endpoints for the Approval of Cancer Drugs and Biologics. Guidance for Industry. U.S. Department of Health and Human Services, FDA, Oncology Center of Excellence, CDERm CBER 2018. ▸ https://www.fda.gov/media/71195/download (Abrufdatum: 04.08.2022).
53. Tröger W, Galun D, Reif M, Schumann A, Stanković N, Milićević M. Quality of Life of Patients With Advanced Pancreatic Cancer During Treatment With Mistletoe: A randomized controlled trial. Dtsch Arztebl Int 2014; 111(29–30): 493–502. ▸ https://doi.org/10.3238/arztebl.2014.0493.
54. Galun D, Tröger W, Milićević M. Cancer Surgery and Supportive Mistletoe Therapy: From Scepticism to Randomised Clinical Trials. In: Zänker KS, Kaveri SV (Hrsg.). Mistletoe: From Mythology to Evidence-Based Medicine. Volume 4: Translational Research in Biomedicine. Basel: Karger 2015. ▸ https://doi.org/10.1159/000375426.
55. Schmidt NEW. Die postoperative komplementäre Therapie des primären Mammakarzinom mit lektinnormiertem Mistelextrakt – eine epidemiologische, kontrollierte retrolektive Kohortenstudie. Dissertation, Humanmedizin. Medizinische Fakultät der Charité, Universitätsmedizin Berlin 2007. ▸ https://doi.org/10.17169/refubium-10460.
56. Oei SL, Thronicke A, Kröz M, Herbstreit C, Schad F. Impact of Viscum album Treatments on the Thermocoherence of Non-Metastasized Breast Cancer Patients. In: Scheer R, Alban S, Becker H, Beer A-M, Blaschek W, Klein R, Kreis W, Leneweit G, Matthes H, Spahn G, Stange R (Hrsg.). Die Mistel in der Tumortherapie 5. Aktueller Stand der Forschung und klinische Anwendung. 1. Aufl. Essen: KVC Verlag 2020.
57. Bock PR, Friedel WE, Hanisch J, Karasmann M, Schneider B. Wirksamkeit und Sicherheit der komplementären Langzeitbehandlung mit einem standardisierten Extrakt aus Europäischer Mistel (Viscum album L.) zusätzlich zur konventionellen adjuvanten onkologischen Therapie bei primärem, nicht metastasiertem Mammakarzinom. Ergebnisse einer multizentrischen, komparativen, epidemiologischen Kohortenstudie in Deutschland und der Schweiz. Arzneimittelforschung 2004; 54(8): 456–466. https://doi.org/10.1055/s-0031-1296999.
58. Borrelli E. Evaluation of the quality of life in breast cancer patients undergoing lectin standardized mistletoe extract therapy. Minerva Medica 2001, 92(Suppl. 1): 105–107.
59. Carlsson M, Arman M, Backman M, Flatters U, Hatschek T, Hamrin E. A Five-year Follow-up of Quality of Life in Women with Breast Cancer in Anthroposophic and Conventional Care. Evid Based Complement Alternat Med 2006; 3(4): 523–531. ▸ https://doi.org/10.1093/ecam/nel042.
60. Heiny BM. Additive Therapie mit standardisiertem Mistelextrakt reduziert die Leukopenie und verbessert die Lebensqualität von Patientinnen mit fortgeschrittenem Mammakarzinom unter palliativer Chemotherapie (VEC-Schema). Krebsmedizin 1991; 12: 3–15.
61. Kienle GS, Glockmann A, Schink M, Kiene H. Viscum album L. extracts in breast and gynaecological cancers: a systematic review of clinical and preclinical research. J Exp Clin Cancer Res 2009; 28: 79. ▸ https://doi.org/10.1186/1756-9966-28-79.
62. Konitzer M, Jäger B, Schüller I, Doering T, Fischer GC. High-dosage mistletoe-treatment (Vysorel) of mamma-carcinoma. A 10 years retrospective of 200 patients from a general medicine practice. Phytomedicine 2000; 7(Suppl. 2): 36.
63. Loewe-Mesch A, Kuehn JJ, Borho K, Abel U, Bauer C, Gerhard I, Schneeweiss A, Sohn C, Strowitzki T, von Hagens C. Adjuvante simultane Mistel-/Chemotherapie bei Mammakarzinom – Einfluss auf Immunparameter, Lebensqualität und Verträglichkeit. Forsch Komplementmed 2008; 15(1): 22–30. ▸ https://doi.org/10.1159/000112860.
64. Orange M, Fonseca M, Lace A, von Laue H-B, Geider S. Durable tumour responses following primary high dose induction with mistletoe extracts: Two case reports. Eur J Integr Med 2010; 2(2): 63–69. ▸ https://doi.org/10.1016/j.eujim.2010.04.001.
65. Pelzer F, Tröger W. Complementary Treatment with Mistletoe Extracts During Chemotherapy: Safety, Neutropenia, Fever, and Quality of Life Assessed in a Randomized Study. J Altern Complement Med 2018; 24(9–10): 954–961. ▸ https://doi.org/10.1089/acm.2018.0159.
66. Piao BK, Wang YX, Xie GR, Mansmann U, Matthes H, Beuth J, Lin HS. Impact of Complementary Mistletoe Extract Treatment on Quality of Life in Breast, Ovarian and Non-small Cell Lung Cancer Patients. A Prospective Randomized Controlled Clinical Trial. Anticancer Res 2004; 24(1): 303–309. ▸ https://ar.iiarjournals.org/content/anticanres/24/1/303.
67. Semiglasov VF, Stepula VV, Dudov A, Lehmacher W, Mengs U. The Standardised Mistletoe Extract PS76A2 Improves QoL in Patients with Breast Cancer Receiving Adjuvant CMF Chemotherapy: A Randomised, Placebo-controlled, Double-blind, Multicentre Clinical Trial. Anticancer Res 2004; 24(2C): 1293–1302. ▸ https://ar.iiarjournals.org/content/24/2C/1293.
68. Semiglazov VF, Stepula VV, Dudov A, Schnitker J, Mengs U. Quality of Life is Improved in Breast Cancer Patients by Standardised Mistletoe Extract PS76A2 during Chemotherapy and Follow-up: A Randomised, Placebo-controlled, Double-blind, Multicentre Clinical Trial. Anticancer Res 2006; 26(2B): 1519–1529. ▸ https://ar.iiarjournals.org/content/26/2B/1519.
69. Tröger W, Jezdić S, Ždrale Z, Tišma N, Hamre HJ, Matijašević M. Quality of Life and Neutropenia in Patients with Early Stage Breast Cancer: A Randomized Pilot Study Comparing Additional Treatment with Mistletoe Extract to Chemotherapy Alone. Breast Cancer (Auckl) 2009; 3: 35–45. ▸ https://doi.org/10.4137/bcbcr.s2905.
70. Tröger W, Matijašević M, Ždrale Z, Tišma N, Jezdić S. Additional therapy with mistletoe extracts in breast cancer patients receiving chemotherapy – a prospective randomized open label pilotstudy. In: Scheer R, Alban S, Becker H, Holzgrabe U, Kemper FH, Kreis W, Matthes H, Schilcher H (Hrsg.). Die Mistel in der Tumortherapie 2. Aktueller Stand der Forschung und klinische Anwendung. 1. Aufl. Essen: KVC Verlag 2009.
71. Tröger W, Ždrale Z, Tišma N, Matijašević M. Additional Therapy with a Mistletoe Product during Adjuvant Chemotherapy of Breast Cancer Patients Improves Quality of Life: An Open Randomized Clinical Pilot Trial. Evid Based Complement Alternat Med 2014; 2014: 430518. ▸ https://doi.org/10.1155/2014/430518.
72. Werthmann PG, Kuck A, Kienle GS. Long-term survival of a patient with relapsing triple-negative breast cancer stage IIB under surgical treatment and Viscum album extracts: A case report. In: Scheer R, Alban S, Becker H, Beer A-M, Blaschek W, Klein R, Kreis W, Leneweit G, Matthes H, Spahn G, Stange R (Hrsg.). Die Mistel in der Tumortherapie 5. Aktueller Stand der Forschung und klinische Anwendung. Essen: KVC Verlag 2020.
73. Cochrane Deutschland. Evidenzbasierte Medizin. ▸ https://www.cochrane.de/ueber-uns/evidenzbasierte-medizin (Abrufdatum: 27.05.2024).
74. Loef M, Paepke D, Walach H. Quality of Life in Breast Cancer Patients Treated With Mistletoe Extracts: A Systematic Review and Meta-Analysis. Integr Cancer Ther 2023; 22: 15347354231198074. ▸ https://doi.org/10.1177/15347354231198074.
75. Kim K-C, Yook J-H, Eisenbraun J, Kim B-S, Huber R. Quality of life, immunomodulation and safety of adjuvant mistletoe treatment in patients with gastric carcinoma – a randomized, controlled pilot study. BMC Complement Altern Med 2012; 12: 172. ▸ https://doi.org/10.1186/1472-6882-12-172.
76. Grah C. Misteltherapie bei nicht kleinzelligem Bronchialkarzinom. Randomisierte, offene Phase-II-Studie zur Untersuchung der Verträglichkeit, Sicherheit und Wirksamkeit von Viscum-album-Extrakt in der palliativen, additiven Behandlung des fortgeschrittenen nichtkleinzelligen Bronchialkarzinoms. Dissertation, Humanmedizin. Medizinische Fakultät der Charité, Universitätsmedizin Berlin 2010.
77. Steuer-Vogt MK, Bonkowsky V, Scholz M, Fauser C, Licht K, Ambrosch P. Einfluss eines ML-1-normierten Mistelextraktes auf die Lebensqualität bei Patienten mit Kopf-Hals-Karzinomen. HNO 2006; 54(4): 277–286. ▸ https://doi.org/10.1007/s00106-005-1318-y.
78. Enesel MB, Acalovschi I, Grosu V, Sbarcea A, Rusu C, Dobre A, Weiss T, Zarković N. Perioperative application of the Viscum album extract Isorel in digestive tract cancer patients. Anticancer Res 2005; 25(6C): 4583–4590. ▸ https://pubmed.ncbi.nlm.nih.gov/16334146/.
79. Kaiser G, Büschel G, Horneber M, Smetak M, Birkmann J, Braun W, Fischer S, von Laue H-B, Scheer R, Gallmeier WM. Studiendesign und erste Ergebnisse einer prospektiven, placebokontrollierten, doppelblinden, randomisierten Studie mit ABNOBAviscum Mali 4. In: Scheer R, Bauer R, Becker H, Berg PA, Fintelmann V (Hrsg.). Die Mistel in der Tumortherapie. Grundlagenforschung und Klinik. 1. Aufl. Essen: KVC Verlag 2001.
80. Heiny BM, Albrecht V, Beuth J. Lebensqualitätsstabilisierung durch Mistellektin-1 normierten Extrakt beim fortgeschrittenen kolorektalen Karzinom. Onkologe 1998; 4(Suppl. 1): 35–39. ▸ https://link.springer.com/article/10.1007/s007610050311.
81. Lenartz D, Stoffel B, Menzel J, Beuth J. Immunoprotective activity of the galactoside-specific lectin from mistletoe after tumor destructive therapy in glioma patients. Anticancer Res 1996; 16(6B): 3799–3802. ▸ https://pubmed.ncbi.nlm.nih.gov/9042260/.
82. Anonymus. Ein neues Krebsheilmittel. Cancrodoron. In: Neues Wiener Journal 1924; 11. Januar: 5.
83. Pelzer F, Loef M, Martin DD, Baumgartner S. Cancer-related fatigue in patients treated with mistletoe extracts: a systematic review and meta-analysis. Support Care Cancer 2022; 30(8): 6405–6418. ▸ https://doi.org/10.1007/s00520-022-06921-x.
84. Mustian KM, Alfano CM, Heckler C, Kleckner AS, Kleckner IR, Leach CR, Mohr D, Palesh OG, Peppone LJ, Piper BF, Scarpato J, Smith T, Sprod LK, Miller SM. Comparison of Pharmaceutical, Psychological, and Exercise Treatments for Cancer-Related Fatigue: A Meta-analysis. JAMA Oncol 2017; 3(7): 961–968. ▸ https://doi.org/10.1001/jamaoncol.2016.6914.
85. Grossarth-Maticek R, Kiene H, Baumgartner S, Ziegler R. Verlängerung der Überlebenszeit von Krebspatienten unter Misteltherapie (Iscador). Ergebnisse einer

epidemiologischen Langzeitstudie. Schweiz Z Ganzheitsmed 2001; 13(4): 217–225.
86. Kienle GS, Mussler M, Fuchs D, Kiene H. The Subjective Dimension of Integrative Cancer Care: A Qualitative Study Exploring the Perspectives, Themes, and Observations of Experienced Doctors from the Area of Anthroposophic Medicine. Explore 2018; 14(5): 342–351. ▸ https://doi.org/10.1016/j.explore.2018.03.007.
87. Friedel WE, Matthes H, Bock PR, Zänker KS. Systematic evaluation of the clinical effects of supportive mistletoe treatment within chemo- and/or radiotherapy protocols and long-term mistletoe application in non metastatic colorectal carcinoma: multicenter, controlled, observational cohort study. J Soc Integr Oncol 2009; 7(4): 137–145. ▸ https://doi.org/10.2310/7200.2009.0014.
88. Mansky PJ, Wallerstedt DB, Sannes TS, Stagl J, Johnson LL, Blackman MR, Grem JL, Swain SM, Monahan BP. NCCAM/NCI Phase 1 Study of Mistletoe Extract and Gemcitabine in Patients with Advanced Solid Tumors. Evid Based Complement Alternat Med 2013; 2013: 964592. ▸ https://doi.org/10.1155/2013/964592. Erratum in: Evid Based Complement Alternat Med 2014; 2014: 606348. ▸ https://doi.org/10.1155/2014/606348.
89. Siegle I, Fritz P, McClellan M, Gutzeit S, Mürdter TE. Combined cytotoxic action of Viscum album agglutinin-1 and anticancer agents against human A549 lung cancer cells. Anticancer Res 2001; 21(4A): 2687–2691. ▸ https://pubmed.ncbi.nlm.nih.gov/11724340/.
90. Weissenstein U, Kunz M, Urech K, Baumgartner S. Interaction of standardized mistletoe (Viscum album) extracts with chemotherapeutic drugs regarding cytostatic and cytotoxic effects in vitro. BMC Complement Altern Med 2014; 14: 6. ▸ https://doi.org/10.1186/1472-6882-14-6.
91. Engdal S, Nilsen OG. In vitro inhibition of CYP3A4 by herbal remedies frequently used by cancer patients. Phytother Res 2009; 23(7): 906–912. ▸ https://doi.org/10.1002/ptr.2750.
92. Doehmer J, Eisenbraun J. Assessment of Extracts from Mistletoe (Viscum album) for Herb-Drug Interaction by Inhibition and Induction of Cytochrome P450 Activities. Phytother Res 2012; 26(1): 11–17. ▸ https://doi.org/10.1002/ptr.3473.
93. Jurin M, Zarković N, Borovic S, Kissel D. Immunmodulation by the Viscum album L. preparation Isorel and its antitumorous effects. In: Scheer R, Becker H, Berg PA (Hrsg.). Grundlagen der Misteltherapie – Aktueller Stand der Forschung und klinische Anwendung. Stuttgart: Hippokrates Verlag 1996.
94. Jurin M, Zarković N, Hrzenjak M, Ilić Z. Antitumorous and immunomodulatory effects of the Viscum album L. preparation Isorel. Oncology 1993; 50(6): 393–398. ▸ https://doi.org/10.1159/000227217.
95. Kienle GS, Mussler M, Fuchs D, Kiene H. Intravenous Mistletoe Treatment in Integrative Cancer Care: A Qualitative Study Exploring the Procedures, Concepts, and Observations of Expert Doctors. Evid Based Complement Alternat Med 2016; 2016: 4628287. ▸ https://doi.org/10.1155/2016/4628287.
96. Matthes H, Buchwald D, Schad F, Jeschke E. Intratumorale Applikation von Viscum album L (Mistelgesamtextrakt; Helixor M®) in der Therapie des inoperablen Pankreaskarzinom. Z Gastroenterol 2007; 45: P015. ▸ https://doi.org/10.1055/s-2007-988162.
97. Rostock M, Huber R, Greiner T, Fritz P, Scheer R, Schueler J, Fiebig HH. Anticancer Activity of a Lectin-rich Mistletoe Extract Injected Intratumorally into Human Pancreatic Cancer Xenografts. Anticancer Res 2005; 25(3B): 1969–1975. ▸ https://ar.iiarjournals.org/content/25/3B/1969.long.
98. Schad F, Axtner J, Buchwald D, Happe A, Popp S, Kröz M, Matthes H. Intratumoral Mistletoe (Viscum album L) Therapy in Patients With Unresectable Pancreas Carcinoma: A Retrospective Analysis. Integr Cancer Ther 2014; 13(4): 332–340. ▸ https://doi.org/10.1177/1534735413513637.
99. Steele ML, Axtner J, Happe A, Kröz M, Matthes H, Schad F. Use and Safety of Intratumoral Application of European Mistletoe (Viscum album L) Preparations in Oncology. Integr Cancer Ther 2015; 14(2): 140–148. ▸ https://doi.org/10.1177/1534735414563977.
100. Chang Y, Cho D, Cho K, Cho M. Viscum pleurodesis is as effective as talc pleurodesis and tends to have less adverse effect. Support Care Cancer 2020; 28(11): 5463–5467. ▸ https://doi.org/10.1007/s00520-020-05405-0.
101. Stumpf C, Büssing A, Schietzel M. Instillation von Mistelextrakten bei malignen Pleuraergüssen. Zytologische und immunologische Ergebnisse einer prospektiven Studie. Forsch Komplementarmed 1996; 3(1): 11.
102. Eisenbraun J. Intravesikaler Mistelextract bei Patienten mit nicht-muskelinvasivem Harnblasenkarzinom: Eine Phase III-Wirksamkeitsstudie – Aktueller Stand. In: Scheer R, Alban S, Becker H, Beer A-M, Blaschek W, Kreis W, Matthes H, Schilcher H, Spahn G, Stange R (Hrsg.). Die Mistel in der Tumortherapie 4. Aktueller Stand der Forschung und klinische Anwendung. 1. Aufl. Essen: KVC Verlag 2016.
103. Heusser P, Berger Braun S, Bertschy M, Burkhard R, Ziegler R, Helwig S, van Wegberg B, Cerny T. Palliative In-Patient Cancer Treatment in an Anthroposophic Hospital: II. Quality of Life during and after Stationary Treatment, and Subjective Treatment Benefits. Forsch Komplementmed 2006; 13(3): 156–166. ▸ https://doi.org/10.1159/000092448.
104. Heusser P, Berger Braun S, Ziegler R, Bertschy M, Helwig S, van Wegberg B, Cerny T. Palliative In-Patient Cancer Treatment in an Anthroposophic Hospital: I. Treatment Patterns and Compliance with Anthroposophic Medicine. Forsch Komplementmed 2006; 13(2): 94–100. ▸ https://doi.org/10.1159/000091694.
105. Rexer H, Geschäftsstelle der AUO. Studie zur Therapie beim nicht-muskelinvasiven Harnblasenkarzinom: Eine Phase-III-Wirksamkeitsstudie zur intravesikalen Instillation von Mistelextrakt bei oberflächlichem Blasenkarzinom (TIM) – AB 40/11 der AUO. Urologe 2016; 55(7): 963–965. ▸ https://doi.org/10.1007/s00120-016-0135-0.
106. Wagner R. Pankreaskopf- Karzinom mit Lebermetastasen inoperabel, Laparotomie und Leberprobeexzision. Krankengeschichte Nr. 3. Mitteilungen aus der Behandlung maligner Tumoren mit Viscum album (Verein für Krebsforschung, Arlesheim) 1987; 19(1): 33–35.
107. Mahfouz MM, Ghaleb HA, Zawawy A, Scheffler A. Significant tumor reduction, improvement of pain and quality of life and normalization of sleeping patterns of cancer patients treated with a high dose of mistletoe. Ann Oncol 1998; 9: 129.
108. Mabed M, El-Helw L, Shamaa S. Phase II study of viscum fraxini-2 in patients with advanced hepatocellular carcinoma. Br J Cancer 2004; 90(1): 65–69. ▸ https://doi.org/10.1038/sj.bjc.6601463.
109. Kirsch A. Successful treatment of metastatic malignant melanoma with Viscum album extract (Iscador M). J Altern Complement Med 2007; 13(4): 443–445. ▸ https://doi.org/10.1089/acm.2007.6175.
110. Seifert G, Laengler A, Tautz C, Seeger K, Henze G. Response to subcutaneous therapy with mistletoe in recurrent multisystem Langerhans cell histiocytosis. Pediatr Blood Cancer 2007; 48(5): 591–592. ▸ https://doi.org/10.1002/pbc.20649.
111. Eldorry AK, Shaker MK, Barakat EM, Abdallah H, Khalifa MO, Elbreedy AM. The effect of a mistletoe preparation with defined lectin content in advanced hepatocellular carcinoma cases. Hepatology 2008; 48(4): 497A–498A.
112. Kunz C, Heiligtag H-R, Hintze A, Urech K. Topische Behandlung des Basalzellkarzinoms mit Viscum album, lipophiler Extrakt 10 %, Unguentum – Eine Fallserie-Studie. In: Scheer R, Alban S, Becker H, Blaschek W, Kemper FH, Kreis W, Matthes H, Schilcher H, Stange R (Hrsg.). Die Mistel in der Tumortherapie 3. Aktueller Stand der Forschung und klinische Anwendung. 1. Aufl. Essen: KVC Verlag 2013.
113. Werthmann PG, Sträter G, Friesland H, Kienle GS. Durable response of cutaneous squamous cell carcinoma following high-dose peri-lesional injections of Viscum album extracts – A case report. Phytomedicine 2013; 20(3–4): 324–327. ▸ https://doi.org/10.1016/j.phymed.2012.11.001.
114. von Schoen-Angerer T, Goyert A, Vagedes J, Kiene H, Merckens H, Kienle GS. Disappearance of an advanced adenomatous colon polyp after intratumoural injection with Viscum album (European mistletoe) extract: a case report. J Gastrointestin Liver Dis 2014; 23(4): 449–452. ▸ https://doi.org/10.15403/jgld.2014.1121.234.acpy.
115. Christen-Clottu O, Klocke P, Burger D, Straub R, Gerber V. Treatment of clinically diagnosed equine sarcoid with a mistletoe extract (Viscum album austriacus). J Vet Intern Med 2010; 24(6): 1483–1489. ▸ https://doi.org/10.1111/j.1939-1676.2010.0597.x.
116. Knottenbelt DC, Patterson-Kane JC, Snalune KL. Clinical Equine Oncology. 1. Aufl. London: Elsevier 2015.
117. Beermann A, Clottu O, Reif M, Biegel U, Unger L, Koch C. A randomized placebo-controlled double-blinded study comparing oral and subcutaneous administration of mistletoe extract for the treatment of equine sarcoid disease. JVet Intern Med 2024; 38(3): 1815–1824. ▸ https://doi.org/10.1111/jvim.17052.
118. Königsberger K, Urech K, Reif M, Baumgartner S, Martin D, Tröger W. Viscum album Lipophilic Extract in Actinic Keratosis, Cutaneous Squamous Cell Carcinoma and Basal Cell Carcinoma: A Retrospective Case Series. Complement Med Res 2024; Online ahead of print. ▸ https://doi.org/10.1159/000537979.
119. Girke M, Debus M, Kröz M. Ascites bei Non-Hodgkin-Lymphom (V.a. splenales Lymphom): Remission nach viermaliger intraperitonealer Viscum-album-Instillation. Der Merkurstab 2012; 65(3): 257–258. ▸ https://doi.org/10.14271/DMS-19967-DE.
120. Gutsch J, Rieger S, Schlodder D. Observational study on treatment of lymphocytic Non Hodgkin's Lymphoma (CLL) with Viscum album products Helixor P or A: clinical course and safety. Phytomedicine 2011; 18(Suppl. 1): S10. ▸ https://doi.org/10.1016/j.phymed.2011.09.025.
121. Delebinski CI, Jäger S, Kemnitz-Hassanin K, Henze G, Lode HN, Seifert G. Therapeutic efficacy of natural compounds from Viscum album L. in acute lymphoblastic leukaemia. Phytomedicine 2011; 18(Suppl. 1): S8. ▸ https://doi.org/10.1016/j.phymed.2011.09.066.
122. Kuehn JJ. Viscum album L. Pini in der Behandlung des Non-Hodgkin-Lymphoms – Wirksamkeit und Risikoeinschätzung einer subkutanen Misteltherapie im Rahmen einer retrospektiven Fallkontrollstudie. In: Scheer R, Alban S, Becker H, Holzgrabe U, Kemper FH, Kreis W, Matthes H, Schilcher H (Hrsg.). Die Mistel in der Tumortherapie 2. Aktueller Stand der Forschung und klinische Anwendung. 1. Aufl. Essen: KVC Verlag 2009.
123. Kuehn JJ. Misteltherapie bei malignen Lymphomen – Neue Erkenntnisse und Erfahrungen im Rahmen einer prospektiven Kasuistikserie bei Patienten mit follikulärem Non-Hodgkin-Lymphom. In: Scheer R, Bauer R, Becker H, Fintelmann V, Kemper FH, Schilcher H (Hrsg.). Fortschritte in der Misteltherapie. Aktueller Stand der Forschung und klinische Anwendung. 1. Aufl. Essen: KVC Verlag 2005.
124. Kuehn JJ. Langfristig guter Verlauf unter Misteltherapie bei einem Patienten mit einem zentroblastisch-zentrozytischen Non-Hodgkin-Lymphom. Dtsch Med Wochenschr 1999; 124(47): 1414–1418. ▸ https://doi.org/10.1055/s-2007-1024555.
125. Kovacs E. Untersuchung des Serumspiegels von IL-6 und von C-reaktivem Protein im Serum von Patienten mit Non-Hodgkin-Lymphomen (NHL). In: Krebstagung Arlesheim, Arlesheim, 19.–21. September 1999. Verein für Krebsforschung 1999.
126. Goyert A. Niedrig dosierte Misteltherapie bei niedrig malignem Non-Hodgkin-Lymphom (Erfahrungsbericht). Erfahrungsheilkunde 1997; Sonderausgabe August: 514–515.
127. Emme B. Zur Misteltherapie der Non-Hodgkin-Lymphome. Der Merkurstab 1994, 47(6): 629. ▸ https://www.anthromedics.org/DMS-16636-DE.
128. Werthmann PG, Helling D, Heusser P, Kienle GS. Tumour response following high-dose intratumoral application of Viscum album on a patient with adenoid cystic carcinoma. BMJ Case Rep 2014; bcr2013203180. ▸ https://doi.org/10.1136/bcr-2013-203180.

129. Werthmann PG, Hintze A, Kienle GS. Complete remission and long-term survival of a patient with melanoma metastases treated with high-dose fever-inducing Viscum album extract: A case report. Medicine (Baltimore) 2017; 96(46): e8731. ▸ https://doi.org/10.1097/md.0000000000008731.
130. Gutsch J, Werthmann PG, Rosenwald A, Kienle GS. Complete Remission and Long-term Survival of a Patient with a Diffuse Large B-cell Lymphoma Under Viscum album Extracts After Resistance to R-CHOP: A Case Report. Anticancer Res 2018; 38(9): 5363–5369. ▸ https://doi.org/10.21873/anticanres.12865.
131. Reynel M, Villegas Y, Kiene H, Werthmann PG, Kienle GS. Intralesional and subcutaneous application of Viscum album L. (European mistletoe) extract in cervical carcinoma in situ: A CARE compliant case report. Medicine (Baltimore) 2018; 97(48): e13420. ▸ https://doi.org/10.1097/md.0000000000013420. Erratum in: Medicine (Baltimore) 2018; 97(52): e13950. ▸ https://doi.org/10.1097/MD.0000000000013950.
132. Werthmann PG, Huber R, Kienle GS. Durable clinical remission of a skull metastasis under intralesional Viscum album extract therapy: Case report. Head Neck 2018; 40(7): E77–E81. ▸ https://doi.org/10.1002/hed.25320.
133. Werthmann PG, Inter P, Welsch T, Sturm AK, Grutzmann R, Debus M, Sterner MG, Kienle GS. Long-term tumor-free survival in a metastatic pancreatic carcinoma patient with FOLFIRINOX/Mitomycin, high-dose, fever inducing Viscum album extracts and subsequent R0 resection: A case report. Medicine (Baltimore) 2018; 97(49): e13243. ▸ https://doi.org/10.1097/MD.0000000000013243.
134. Werthmann PG, Kempenich R, Kienle GS. Long-Term Tumor-Free Survival in a Patient with Stage IV Epithelial Ovarian Cancer Undergoing High-Dose Chemotherapy and Viscum album Extract Treatment: A Case Report. Perm J 2018; 23: 18–025. ▸ https://doi.org/10.7812/TPP/18-025. Erratum in: Perm J 2019; 23: 18–025. ▸ https://doi.org/ 10.7812/TPP/18-025-err.
135. Werthmann PG, Kindermann L, Kienle GS. A 21-year course of Merkel cell carcinoma with adjuvant Viscum album extract treatment: A case report. Complement Ther Med 2018; 38: 58–60. ▸ https://doi.org/10.1016/j.ctim.2018.04.001.
136. Reynel M, Villegas Y, Kiene H, Werthmann PG, Kienle GS. Bilateral Asynchronous Renal Cell Carcinoma With Lung Metastases: A Case Report of a Patient Treated Solely With High-dose Intravenous and Subcutaneous Viscum album Extract for a Second Renal Lesion. Anticancer Res 2019; 39(10): 5597–5604. ▸ https://doi.org/10.21873/anticanres.13754.
137. Werthmann PG, Kempenich R, Lang-Avérous G, Kienle GS. Long-term survival of a patient with advanced pancreatic cancer under adjunct treatment with Viscum album extracts: A case report. World J Gastroenterol 2019; 25(12): 1524–1530. ▸ https://doi.org/10.3748/wjg.v25.i12.1524.
138. Werthmann PG, Kindermann L, Kienle GS. Chemoimmunotherapy in Advanced Renal Cell Carcinoma: A Case Report of a Long-Term Survivor Adjunctly Treated with Viscum album Extracts. Complement Med Res 2019; 26(4): 276–279. ▸ https://doi.org/10.1159/000496866.
139. Reynel M, Villegas Y, Werthmann PG, Kiene H, Kienle GS. Long-term survival of a patient with an inoperable thymic neuroendocrine tumor stage IIIa under sole treatment with Viscum album extract: A CARE compliant clinical case report. Medicine (Baltimore) 2020; 99(5): e18990. ▸ https://doi.org/10.1097/MD.0000000000018990.
140. Orange M, Poidimani N, Crosignani A, Werthmann PG, Bertotto C. Complete, Durable Remission of Advanced Hepatocellular Carcinoma under Treatment with Viscum album Extracts: A Case Report. Complement Med Red 2022; 29(6): 484–491. ▸ https://doi.org/10.1159/000525409.
141. Reynel M, Villegas Y, Werthmann PG, Kiene H, Kienle GS. Long-Term Survival of a Patient with Recurrent Dedifferentiated High-Grade Liposarcoma of the Retroperitoneum Under Adjuvant Treatment with Viscum album L. Extract: A Case Report. Integr Cancer Ther 2021; 20: 1534735421995258. ▸ https://doi.org/10.1177/1534735421995258.
142. Gagnier JJ, Riley D, Altman DG, Moher D, Sox H, Kienle G. The CARE Guidelines: Consensus-Based Clinical Case Reporting Guideline Development. Dtsch Arztebl Int 2013; 110(37): 603–608. ▸ https://doi.org/10.3238/arztebl.2013.0603.
143. Werthmann PG, Kiene H, Kienle G. How to Capture Clinical Experience – An Example From Case Reporting in Viscum album Extract Therapy. Glob Adv Health Med 2020; 9: OA14.02. ▸ https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7290253/.