



Russ Salakhutdinov



9 October at 21:59 · 🌐

I am teaching a Deep Learning graduate course this Fall at CMU with over 300 MSc and PhD students enrolled.

Today, after our midterm, I received the following anonymous feedback: "Did I take the wrong exam? Does this exam cover too little machine learning stuff and focus too much on mathematics?"

I guess there is a common belief that Deep Learning is all about installing TensorFlow or PyTorch and training a gigantic convnet on multiple GPUs 😊



Like



Comment



Share



Csaba Szepesvari, Anima Anandkumar and 1.3k others

Outline

Goals

- ▶ Review the supervised learning setting
- ▶ Describe the linear regression framework
- ▶ Apply the linear model to make predictions
- ▶ Derive the least squares estimate

Outline

Goals

- ▶ Review the supervised learning setting
- ▶ Describe the linear regression framework
- ▶ Apply the linear model to make predictions
- ▶ Derive the least squares estimate

Supervised Learning Setting

- ▶ Data consists of **input** and **output** pairs
- ▶ Inputs (also covariates, independent variables, predictors, features)
- ▶ Output (also variates, dependent variables, targets, labels)

Why study linear regression?

- ▶ **Least squares** is at least 200 years old going back to Legendre and Gauss
- ▶ Francis Galton (1886): “Regression to the mean”

Why study linear regression?

- ▶ **Least squares** is at least 200 years old going back to Legendre and Gauss
- ▶ Francis Galton (1886): “Regression to the mean”
- ▶ Often real processes can be **approximated** by linear models
- ▶ More complex models require understanding linear regression

Why study linear regression?

- ▶ **Least squares** is at least 200 years old going back to Legendre and Gauss
- ▶ Francis Galton (1886): “Regression to the mean”
- ▶ Often real processes can be **approximated** by linear models
- ▶ More complex models require understanding linear regression
- ▶ Closed form analytic solutions can be obtained
- ▶ Many **key notions** of machine learning can be introduced

A toy example : Commute Times

Want to predict commute time into city centre

What variables would be useful?

- ▶ Distance to city centre
- ▶ Day of the week



Data

dist (km)	day	commute time (min)
2.7	fri	25
4.1	mon	33
1.0	sun	15
5.2	tue	45
2.8	sat	22



Linear Models

Suppose the input is a vector $\mathbf{x} \in \mathbb{R}^D$ and the output is $y \in \mathbb{R}$.

Linear Models

Suppose the input is a vector $\mathbf{x} \in \mathbb{R}^D$ and the output is $y \in \mathbb{R}$.

We have data $\langle (\mathbf{x}_i, y_i) \rangle_{i=1}^N$

Notation: data dimension D , size of dataset N , column vectors

Linear Models

Suppose the input is a vector $\mathbf{x} \in \mathbb{R}^D$ and the output is $y \in \mathbb{R}$.

We have data $\langle (\mathbf{x}_i, y_i) \rangle_{i=1}^N$

Notation: data dimension D , size of dataset N , column vectors

Linear Model

$$y = w_0 + x_1 w_1 + \cdots + x_D w_D + \epsilon$$

Bias/intercept

Noise/uncertainty

Linear Models : Commute Time

Linear Model

$$y = w_0 + x_1 w_1 + \cdots + x_D w_D + \epsilon$$

Bias/intercept

Noise/uncertainty

Linear Models : Commute Time

Linear Model

$$y = w_0 + x_1 w_1 + \cdots + x_D w_D + \epsilon$$

Bias/intercept

Noise/uncertainty

Input encoding: mon-sun has to be converted to a number

- ▶ monday: 0, tuesday: 1, ..., sunday: 6
- ▶ 0 if weekend, 1 if weekday

Linear Models : Commute Time

Linear Model

$$y = w_0 + x_1 w_1 + \cdots + x_D w_D + \epsilon$$

Bias/intercept

Noise/uncertainty

Input encoding: mon-sun has to be converted to a number

- ▶ monday: 0, tuesday: 1, ..., sunday: 6
- ▶ 0 if weekend, 1 if weekday

Using 0-6 is a **bad** encoding.
Use seven 0-1 features instead
called **one-hot** encoding

Linear Models : Commute Time

Linear Model

$$y = w_0 + x_1 w_1 + \cdots + x_D w_D + \epsilon$$

Bias/intercept

Noise/uncertainty

Input encoding: mon-sun has to be converted to a number

- ▶ monday: 0, tuesday: 1, ..., sunday: 6
- ▶ 0 if weekend, 1 if weekday

Using 0-6 is a **bad** encoding.
Use seven 0-1 features instead
called **one-hot** encoding

Say $x_1 \in \mathbb{R}$ (distance) and $x_2 \in \{0, 1\}$ (weekend/weekday)

Linear model for commute time

$$y = w_0 + w_1 x_1 + w_2 x_2 + \epsilon$$

Linear Model : Adding a feature for bias term

dist	day	commute time
x_1	x_2	y
2.7	fri	25
4.1	mon	33
1.0	sun	15
5.2	tue	45
2.8	sat	22

Linear Model : Adding a feature for bias term

dist	day	commute time		one	dist	day	commute time
x_1	x_2	y		x_0	x_1	x_2	y
2.7	fri	25	\Leftrightarrow	1	2.7	fri	25
4.1	mon	33		1	4.1	mon	33
1.0	sun	15		1	1.0	sun	15
5.2	tue	45		1	5.2	tue	45
2.8	sat	22		1	2.8	sat	22

Linear Model : Adding a feature for bias term

dist	day	commute time
x_1	x_2	y
2.7	fri	25
4.1	mon	33
1.0	sun	15
5.2	tue	45
2.8	sat	22



one	dist	day	commute time
x_0	x_1	x_2	y
1	2.7	fri	25
1	4.1	mon	33
1	1.0	sun	15
1	5.2	tue	45
1	2.8	sat	22

Model

$$y = w_0 + w_1x_1 + w_2x_2 + \epsilon$$

Model

$$\begin{aligned} y &= w_0x_0 + w_1x_1 + w_2x_2 + \epsilon \\ &= \mathbf{w} \cdot \mathbf{x} + \epsilon \end{aligned}$$

Learning Linear Models

Data: $\langle (\mathbf{x}_i, y_i) \rangle_{i=1}^N$, where $\mathbf{x}_i \in \mathbb{R}^D$ and $y_i \in \mathbb{R}$

Model parameter \mathbf{w} , where $\mathbf{w} \in \mathbb{R}^D$

Learning Linear Models

Data: $\langle (\mathbf{x}_i, y_i) \rangle_{i=1}^N$, where $\mathbf{x}_i \in \mathbb{R}^D$ and $y_i \in \mathbb{R}$

Model parameter \mathbf{w} , where $\mathbf{w} \in \mathbb{R}^D$


Training phase: (learning/estimation \mathbf{w} from data)

Learning Linear Models

Data: $\langle (\mathbf{x}_i, y_i) \rangle_{i=1}^N$, where $\mathbf{x}_i \in \mathbb{R}^D$ and $y_i \in \mathbb{R}$

Model parameter \mathbf{w} , where $\mathbf{w} \in \mathbb{R}^D$

Training phase: (learning/estimation \mathbf{w} from data)




Learning Linear Models

Data: $\langle (\mathbf{x}_i, y_i) \rangle_{i=1}^N$, where $\mathbf{x}_i \in \mathbb{R}^D$ and $y_i \in \mathbb{R}$

Model parameter \mathbf{w} , where $\mathbf{w} \in \mathbb{R}^D$

Training phase: (learning/estimation \mathbf{w} from data)




Testing/Deployment phase: (predict $\hat{y}_{\text{new}} = \mathbf{x}_{\text{new}} \cdot \mathbf{w}$)

Learning Linear Models

Data: $\langle (\mathbf{x}_i, y_i) \rangle_{i=1}^N$, where $\mathbf{x}_i \in \mathbb{R}^D$ and $y_i \in \mathbb{R}$

Model parameter \mathbf{w} , where $\mathbf{w} \in \mathbb{R}^D$

Training phase: (learning/estimation \mathbf{w} from data)



Testing/Deployment phase: (predict $\hat{y}_{\text{new}} = \mathbf{x}_{\text{new}} \cdot \mathbf{w}$)

- ▶ How different is \hat{y}_{new} from y_{new} (actual observation)?
- ▶ We should keep some data aside for testing before deploying a model

$$\langle (x_i, y_i) \rangle_{i=1}^N, \quad \text{where } x_i \in \mathbb{R} \text{ and } y_i \in \mathbb{R}$$

$$\widehat{y}(x) = w_0 + x \cdot w_1, \quad (\text{no noise term in } \widehat{y})$$

$$\mathcal{L}(\mathbf{w}) = \mathcal{L}(w_0, w_1) = \frac{1}{2N} \sum_{i=1}^N (\widehat{y}_i - y_i)^2 = \frac{1}{2N} \sum_{i=1}^N (w_0 + x_i \cdot w_1 - y_i)^2$$

$\langle (x_i, y_i) \rangle_{i=1}^N$, where $x_i \in \mathbb{R}$ and $y_i \in \mathbb{R}$

$\hat{y}(x) = w_0 + x \cdot w_1$, (no noise term in \hat{y})

$$\mathcal{L}(\mathbf{w}) = \mathcal{L}(w_0, w_1) = \frac{1}{2N} \sum_{i=1}^N (\hat{y}_i - y_i)^2 = \frac{1}{2N} \sum_{i=1}^N (w_0 + x_i \cdot w_1 - y_i)^2$$


Loss function
Cost function
Objective Function
Energy Function
Notation - \mathcal{L}, J, E, R

$\langle (x_i, y_i) \rangle_{i=1}^N$, where $x_i \in \mathbb{R}$ and $y_i \in \mathbb{R}$

$\hat{y}(x) = w_0 + x \cdot w_1$, (no noise term in \hat{y})

$$\mathcal{L}(\mathbf{w}) = \mathcal{L}(w_0, w_1) = \frac{1}{2N} \sum_{i=1}^N (\hat{y}_i - y_i)^2 = \frac{1}{2N} \sum_{i=1}^N (w_0 + x_i \cdot w_1 - y_i)^2$$

Predict commute time using only distance



Loss function
Cost function
Objective Function
Energy Function
Notation - \mathcal{L}, J, E, R

This objective is known
as the residual sum
of squares or (RSS)

The estimate (w_0, w_1)
is known as the least
squares estimate

$$\langle (x_i, y_i) \rangle_{i=1}^N, \quad \text{where } x_i \in \mathbb{R} \text{ and } y_i \in \mathbb{R}$$

$$\widehat{y}(x) = w_0 + x \cdot w_1, \quad (\text{no noise term in } \widehat{y})$$

$$\mathcal{L}(\mathbf{w}) = \mathcal{L}(w_0, w_1) = \frac{1}{2N} \sum_{i=1}^N (\widehat{y}_i - y_i)^2 = \frac{1}{2N} \sum_{i=1}^N (w_0 + x_i \cdot w_1 - y_i)^2$$

$$\langle (x_i, y_i) \rangle_{i=1}^N, \quad \text{where } x_i \in \mathbb{R} \text{ and } y_i \in \mathbb{R}$$

$$\widehat{y}(x) = w_0 + x \cdot w_1, \quad (\text{no noise term in } \widehat{y})$$

$$\mathcal{L}(\mathbf{w}) = \mathcal{L}(w_0, w_1) = \frac{1}{2N} \sum_{i=1}^N (\widehat{y}_i - y_i)^2 = \frac{1}{2N} \sum_{i=1}^N (w_0 + x_i \cdot w_1 - y_i)^2$$

$$\frac{\partial \mathcal{L}}{\partial w_0} = \frac{1}{N} \sum_{i=1}^N (w_0 + w_1 \cdot x_i - y_i)$$

$$\langle (x_i, y_i) \rangle_{i=1}^N, \quad \text{where } x_i \in \mathbb{R} \text{ and } y_i \in \mathbb{R}$$

$$\widehat{y}(x) = w_0 + x \cdot w_1, \quad (\text{no noise term in } \widehat{y})$$

$$\mathcal{L}(\mathbf{w}) = \mathcal{L}(w_0, w_1) = \frac{1}{2N} \sum_{i=1}^N (\widehat{y}_i - y_i)^2 = \frac{1}{2N} \sum_{i=1}^N (w_0 + x_i \cdot w_1 - y_i)^2$$

$$\frac{\partial \mathcal{L}}{\partial w_0} = \frac{1}{N} \sum_{i=1}^N (w_0 + w_1 \cdot x_i - y_i)$$

$$\frac{\partial \mathcal{L}}{\partial w_1} = \frac{1}{N} \sum_{i=1}^N (w_0 + w_1 \cdot x_i - y_i) x_i$$

$$\langle (x_i, y_i) \rangle_{i=1}^N, \quad \text{where } x_i \in \mathbb{R} \text{ and } y_i \in \mathbb{R}$$

$$\widehat{y}(x) = w_0 + x \cdot w_1, \quad (\text{no noise term in } \widehat{y})$$

$$\mathcal{L}(\mathbf{w}) = \mathcal{L}(w_0, w_1) = \frac{1}{2N} \sum_{i=1}^N (\widehat{y}_i - y_i)^2 = \frac{1}{2N} \sum_{i=1}^N (w_0 + x_i \cdot w_1 - y_i)^2$$

$$\frac{\partial \mathcal{L}}{\partial w_0} = \frac{1}{N} \sum_{i=1}^N (w_0 + w_1 \cdot x_i - y_i)$$

$$\frac{\partial \mathcal{L}}{\partial w_1} = \frac{1}{N} \sum_{i=1}^N (w_0 + w_1 \cdot x_i - y_i) x_i$$

We obtain the solution for (w_0, w_1) by setting the partial derivatives to 0 and solving the resulting system.
(Normal Equations)

$$\langle (x_i, y_i) \rangle_{i=1}^N, \quad \text{where } x_i \in \mathbb{R} \text{ and } y_i \in \mathbb{R}$$

$$\widehat{y}(x) = w_0 + x \cdot w_1, \quad (\text{no noise term in } \widehat{y})$$

$$\mathcal{L}(\mathbf{w}) = \mathcal{L}(w_0, w_1) = \frac{1}{2N} \sum_{i=1}^N (\widehat{y}_i - y_i)^2 = \frac{1}{2N} \sum_{i=1}^N (w_0 + x_i \cdot w_1 - y_i)^2$$

$$\frac{\partial \mathcal{L}}{\partial w_0} = \frac{1}{N} \sum_{i=1}^N (w_0 + w_1 \cdot x_i - y_i)$$

$$\frac{\partial \mathcal{L}}{\partial w_1} = \frac{1}{N} \sum_{i=1}^N (w_0 + w_1 \cdot x_i - y_i) x_i$$

We obtain the solution for (w_0, w_1) by setting the partial derivatives to 0 and solving the resulting system.
(Normal Equations)

$$w_0 + w_1 \cdot \frac{\sum_i x_i}{N} = \frac{\sum_i y_i}{N} \quad (1)$$

$$w_0 \cdot \frac{\sum_i x_i}{N} + w_1 \cdot \frac{\sum_i x_i^2}{N} = \frac{\sum_i x_i y_i}{N} \quad (2)$$

$$\langle (x_i, y_i) \rangle_{i=1}^N, \quad \text{where } x_i \in \mathbb{R} \text{ and } y_i \in \mathbb{R}$$

$$\hat{y}(x) = w_0 + x \cdot w_1, \quad (\text{no noise term in } \hat{y})$$

$$\mathcal{L}(\mathbf{w}) = \mathcal{L}(w_0, w_1) = \frac{1}{2N} \sum_{i=1}^N (\hat{y}_i - y_i)^2 = \frac{1}{2N} \sum_{i=1}^N (w_0 + x_i \cdot w_1 - y_i)^2$$

$$\frac{\partial \mathcal{L}}{\partial w_0} = \frac{1}{N} \sum_{i=1}^N (w_0 + w_1 \cdot x_i - y_i)$$

$$\frac{\partial \mathcal{L}}{\partial w_1} = \frac{1}{N} \sum_{i=1}^N (w_0 + w_1 \cdot x_i - y_i) x_i$$

We obtain the solution for (w_0, w_1) by setting the partial derivatives to 0 and solving the resulting system.
(Normal Equations)

$$w_0 + w_1 \cdot \frac{\sum_i x_i}{N} = \frac{\sum_i y_i}{N} \quad (1)$$

$$w_0 \cdot \frac{\sum_i x_i}{N} + w_1 \cdot \frac{\sum_i x_i^2}{N} = \frac{\sum_i x_i y_i}{N} \quad (2)$$

$$\bar{x} = \frac{\sum_i x_i}{N}$$

$$\bar{y} = \frac{\sum_i y_i}{N}$$

$$\widehat{\text{var}}(x) = \frac{\sum_i x_i^2}{N} - \bar{x}^2$$

$$\widehat{\text{cov}}(x, y) = \frac{\sum_i x_i y_i}{N} - \bar{x} \cdot \bar{y}$$

$$\langle (x_i, y_i) \rangle_{i=1}^N, \quad \text{where } x_i \in \mathbb{R} \text{ and } y_i \in \mathbb{R}$$

$$\hat{y}(x) = w_0 + x \cdot w_1, \quad (\text{no noise term in } \hat{y})$$

$$\mathcal{L}(\mathbf{w}) = \mathcal{L}(w_0, w_1) = \frac{1}{2N} \sum_{i=1}^N (\hat{y}_i - y_i)^2 = \frac{1}{2N} \sum_{i=1}^N (w_0 + x_i \cdot w_1 - y_i)^2$$

$$\frac{\partial \mathcal{L}}{\partial w_0} = \frac{1}{N} \sum_{i=1}^N (w_0 + w_1 \cdot x_i - y_i)$$

$$\frac{\partial \mathcal{L}}{\partial w_1} = \frac{1}{N} \sum_{i=1}^N (w_0 + w_1 \cdot x_i - y_i) x_i$$

We obtain the solution for (w_0, w_1) by setting the partial derivatives to 0 and solving the resulting system.
(Normal Equations)

$$w_0 + w_1 \cdot \frac{\sum_i x_i}{N} = \frac{\sum_i y_i}{N} \quad (1)$$

$$w_0 \cdot \frac{\sum_i x_i}{N} + w_1 \cdot \frac{\sum_i x_i^2}{N} = \frac{\sum_i x_i y_i}{N} \quad (2)$$

$$\bar{x} = \frac{\sum_i x_i}{N}$$

$$\bar{y} = \frac{\sum_i y_i}{N}$$

$$\widehat{\text{var}}(x) = \frac{\sum_i x_i^2}{N} - \bar{x}^2$$

$$\widehat{\text{cov}}(x, y) = \frac{\sum_i x_i y_i}{N} - \bar{x} \cdot \bar{y}$$

$$w_1 = \frac{\widehat{\text{cov}}(x, y)}{\widehat{\text{var}}(x)}$$

$$w_0 = \bar{y} - w_1 \cdot \bar{x}$$

Linear Regression : General Case

Recall that the linear model is

$$\hat{y}_i = \sum_{j=0}^D x_{ij} w_j$$

where we assume that $x_{i0} = 1$ for all \mathbf{x}_i , so that the bias term w_0 does not need to be treated separately.

Linear Regression : General Case

Recall that the linear model is

$$\hat{y}_i = \sum_{j=0}^D x_{ij} w_j$$

where we assume that $x_{i0} = 1$ for all \mathbf{x}_i , so that the bias term w_0 does not need to be treated separately.

Expressing everything in matrix notation

$$\hat{\mathbf{y}} = \mathbf{X}\mathbf{w}$$

Here we have $\hat{\mathbf{y}} \in \mathbb{R}^{N \times 1}$, $\mathbf{X} \in \mathbb{R}^{N \times (D+1)}$ and $\mathbf{w} \in \mathbb{R}^{(D+1) \times 1}$

$$\begin{matrix} \hat{\mathbf{y}}_{N \times 1} \\ \begin{bmatrix} \hat{y}_1 \\ \hat{y}_2 \\ \vdots \\ \hat{y}_N \end{bmatrix} \end{matrix} = \begin{matrix} \mathbf{X}_{N \times (D+1)} \\ \begin{bmatrix} \mathbf{x}_1^T \\ \mathbf{x}_2^T \\ \vdots \\ \mathbf{x}_N^T \end{bmatrix} \end{matrix} \begin{matrix} \mathbf{w}_{(D+1) \times 1} \\ \begin{bmatrix} w_0 \\ \vdots \\ w_D \end{bmatrix} \end{matrix} = \begin{matrix} \mathbf{X}_{N \times (D+1)} \\ \begin{bmatrix} x_{10} & \cdots & x_{1D} \\ x_{20} & \cdots & x_{2D} \\ \vdots & \ddots & \vdots \\ x_{N0} & \cdots & x_{ND} \end{bmatrix} \end{matrix} \begin{matrix} \mathbf{w}_{(D+1) \times 1} \\ \begin{bmatrix} w_0 \\ \vdots \\ w_D \end{bmatrix} \end{matrix}$$

Back to toy example

one	dist (km)	weekday?	commute time (min)
1	2.7	1 (fri)	25
1	4.1	1 (mon)	33
1	1.0	0 (sun)	15
1	5.2	1 (tue)	45
1	2.8	0 (sat)	22

We have $N = 5$, $D + 1 = 3$ and so we get

$$\mathbf{y} = \begin{bmatrix} 25 \\ 33 \\ 15 \\ 45 \\ 22 \end{bmatrix}, \quad \mathbf{X} = \begin{bmatrix} 1 & 2.7 & 1 \\ 1 & 4.1 & 1 \\ 1 & 1.0 & 0 \\ 1 & 5.2 & 1 \\ 1 & 2.8 & 0 \end{bmatrix}, \quad \mathbf{w} = \begin{bmatrix} w_0 \\ w_1 \\ w_2 \end{bmatrix}$$

Back to toy example

one	dist (km)	weekday?	commute time (min)
1	2.7	1 (fri)	25
1	4.1	1 (mon)	33
1	1.0	0 (sun)	15
1	5.2	1 (tue)	45
1	2.8	0 (sat)	22

We have $N = 5$, $D + 1 = 3$ and so we get

$$\mathbf{y} = \begin{bmatrix} 25 \\ 33 \\ 15 \\ 45 \\ 22 \end{bmatrix}, \quad \mathbf{X} = \begin{bmatrix} 1 & 2.7 & 1 \\ 1 & 4.1 & 1 \\ 1 & 1.0 & 0 \\ 1 & 5.2 & 1 \\ 1 & 2.8 & 0 \end{bmatrix}, \quad \mathbf{w} = \begin{bmatrix} w_0 \\ w_1 \\ w_2 \end{bmatrix}$$

Suppose we get $\mathbf{w} = [6.09, 6.53, 2.11]^T$. Then our predictions would be


$$\hat{\mathbf{y}} = \begin{bmatrix} 25.83 \\ 34.97 \\ 12.62 \\ 42.16 \\ 24.37 \end{bmatrix}$$

Least Squares Estimate : Minimise the Squared Error

$$\mathcal{L}(\mathbf{w}) = \frac{1}{2N} \sum_{i=1}^N (\mathbf{x}_i^\top \mathbf{w} - y_i)^2 = \frac{1}{2N} (\mathbf{X}\mathbf{w} - \mathbf{y})^\top (\mathbf{X}\mathbf{w} - \mathbf{y})$$

Least Squares Estimate : Minimise the Squared Error

$$\mathcal{L}(\mathbf{w}) = \frac{1}{2N} \sum_{i=1}^N (\mathbf{x}_i^\top \mathbf{w} - y_i)^2 = \frac{1}{2N} (\mathbf{X}\mathbf{w} - \mathbf{y})^\top (\mathbf{X}\mathbf{w} - \mathbf{y})$$



Finding Optimal Solutions using Calculus

$$\mathcal{L}(\mathbf{w}) = \frac{1}{2N} \sum_{i=1}^N (\mathbf{x}_i^T \mathbf{w} - y_i)^2 = \frac{1}{2N} (\mathbf{X}\mathbf{w} - \mathbf{y})^T (\mathbf{X}\mathbf{w} - \mathbf{y})$$

Finding Optimal Solutions using Calculus

$$\begin{aligned}\mathcal{L}(\mathbf{w}) &= \frac{1}{2N} \sum_{i=1}^N (\mathbf{x}_i^\top \mathbf{w} - y_i)^2 = \frac{1}{2N} (\mathbf{X}\mathbf{w} - \mathbf{y})^\top (\mathbf{X}\mathbf{w} - \mathbf{y}) \\ &= \frac{1}{2N} \left(\mathbf{w}^\top (\mathbf{X}^\top \mathbf{X}) \mathbf{w} - \mathbf{w}^\top \mathbf{X}^\top \mathbf{y} - \mathbf{y}^\top \mathbf{X} \mathbf{w} + \mathbf{y}^\top \mathbf{y} \right)\end{aligned}$$

Finding Optimal Solutions using Calculus

$$\begin{aligned}\mathcal{L}(\mathbf{w}) &= \frac{1}{2N} \sum_{i=1}^N (\mathbf{x}_i^\top \mathbf{w} - y_i)^2 = \frac{1}{2N} (\mathbf{X}\mathbf{w} - \mathbf{y})^\top (\mathbf{X}\mathbf{w} - \mathbf{y}) \\ &= \frac{1}{2N} \left(\mathbf{w}^\top (\mathbf{X}^\top \mathbf{X}) \mathbf{w} - \mathbf{w}^\top \mathbf{X}^\top \mathbf{y} - \mathbf{y}^\top \mathbf{X} \mathbf{w} + \mathbf{y}^\top \mathbf{y} \right) \\ &= \frac{1}{2N} \left(\mathbf{w}^\top (\mathbf{X}^\top \mathbf{X}) \mathbf{w} - 2 \cdot \mathbf{y}^\top \mathbf{X} \mathbf{w} + \mathbf{y}^\top \mathbf{y} \right) \\ &= \dots\end{aligned}$$

Finding Optimal Solutions using Calculus

$$\begin{aligned}\mathcal{L}(\mathbf{w}) &= \frac{1}{2N} \sum_{i=1}^N (\mathbf{x}_i^\top \mathbf{w} - y_i)^2 = \frac{1}{2N} (\mathbf{X}\mathbf{w} - \mathbf{y})^\top (\mathbf{X}\mathbf{w} - \mathbf{y}) \\ &= \frac{1}{2N} \left(\mathbf{w}^\top (\mathbf{X}^\top \mathbf{X}) \mathbf{w} - \mathbf{w}^\top \mathbf{X}^\top \mathbf{y} - \mathbf{y}^\top \mathbf{X} \mathbf{w} + \mathbf{y}^\top \mathbf{y} \right) \\ &= \frac{1}{2N} \left(\mathbf{w}^\top (\mathbf{X}^\top \mathbf{X}) \mathbf{w} - 2 \cdot \mathbf{y}^\top \mathbf{X} \mathbf{w} + \mathbf{y}^\top \mathbf{y} \right) \\ &= \dots\end{aligned}$$

Then, write out all partial derivatives to form the gradient $\nabla_{\mathbf{w}} \mathcal{L}$

Finding Optimal Solutions using Calculus

$$\begin{aligned}\mathcal{L}(\mathbf{w}) &= \frac{1}{2N} \sum_{i=1}^N (\mathbf{x}_i^\top \mathbf{w} - y_i)^2 = \frac{1}{2N} (\mathbf{X}\mathbf{w} - \mathbf{y})^\top (\mathbf{X}\mathbf{w} - \mathbf{y}) \\ &= \frac{1}{2N} \left(\mathbf{w}^\top (\mathbf{X}^\top \mathbf{X}) \mathbf{w} - \mathbf{w}^\top \mathbf{X}^\top \mathbf{y} - \mathbf{y}^\top \mathbf{X} \mathbf{w} + \mathbf{y}^\top \mathbf{y} \right) \\ &= \frac{1}{2N} \left(\mathbf{w}^\top (\mathbf{X}^\top \mathbf{X}) \mathbf{w} - 2 \cdot \mathbf{y}^\top \mathbf{X} \mathbf{w} + \mathbf{y}^\top \mathbf{y} \right) \\ &= \dots\end{aligned}$$

Then, write out all partial derivatives to form the gradient $\nabla_{\mathbf{w}} \mathcal{L}$

$$\frac{\partial \mathcal{L}}{\partial w_0} = \dots$$

$$\frac{\partial \mathcal{L}}{\partial w_1} = \dots$$

$$\vdots$$

$$\frac{\partial \mathcal{L}}{\partial w_D} = \dots$$

Finding Optimal Solutions using Calculus

$$\begin{aligned}\mathcal{L}(\mathbf{w}) &= \frac{1}{2N} \sum_{i=1}^N (\mathbf{x}_i^\top \mathbf{w} - y_i)^2 = \frac{1}{2N} (\mathbf{X}\mathbf{w} - \mathbf{y})^\top (\mathbf{X}\mathbf{w} - \mathbf{y}) \\ &= \frac{1}{2N} \left(\mathbf{w}^\top (\mathbf{X}^\top \mathbf{X}) \mathbf{w} - \mathbf{w}^\top \mathbf{X}^\top \mathbf{y} - \mathbf{y}^\top \mathbf{X} \mathbf{w} + \mathbf{y}^\top \mathbf{y} \right) \\ &= \frac{1}{2N} \left(\mathbf{w}^\top (\mathbf{X}^\top \mathbf{X}) \mathbf{w} - 2 \cdot \mathbf{y}^\top \mathbf{X} \mathbf{w} + \mathbf{y}^\top \mathbf{y} \right) \\ &= \dots\end{aligned}$$

Then, write out all partial derivatives to form the gradient $\nabla_{\mathbf{w}} \mathcal{L}$

$$\frac{\partial \mathcal{L}}{\partial w_0} = \dots$$

$$\frac{\partial \mathcal{L}}{\partial w_1} = \dots$$

$$\vdots$$

$$\frac{\partial \mathcal{L}}{\partial w_D} = \dots$$

Instead, we will use matrix calculus shortcuts to differentiate using matrix notation directly

Differentiating Matrix Expressions

Rules (Tricks)

(i) Linear Form Expressions: $\nabla_{\mathbf{w}} \left(\mathbf{c}^T \mathbf{w} \right) = \mathbf{c}$

Differentiating Matrix Expressions

Rules (Tricks)

(i) Linear Form Expressions: $\nabla_{\mathbf{w}} \left(\mathbf{c}^T \mathbf{w} \right) = \mathbf{c}$

$$\mathbf{c}^T \mathbf{w} = \sum_{j=0}^D c_j w_j$$

$$\frac{\partial (\mathbf{c}^T \mathbf{w})}{\partial w_j} = c_j,$$

$$\text{and so } \nabla_{\mathbf{w}} \left(\mathbf{c}^T \mathbf{w} \right) = \mathbf{c} \quad (3)$$

Differentiating Matrix Expressions

Rules (Tricks)

(i) Linear Form Expressions: $\nabla_{\mathbf{w}} \left(\mathbf{c}^T \mathbf{w} \right) = \mathbf{c}$

$$\mathbf{c}^T \mathbf{w} = \sum_{j=0}^D c_j w_j$$

$$\frac{\partial (\mathbf{c}^T \mathbf{w})}{\partial w_j} = c_j, \quad \text{and so} \quad \nabla_{\mathbf{w}} \left(\mathbf{c}^T \mathbf{w} \right) = \mathbf{c} \quad (3)$$

(ii) Quadratic Form Expressions:

$$\nabla_{\mathbf{w}} \left(\mathbf{w}^T \mathbf{A} \mathbf{w} \right) = \mathbf{A} \mathbf{w} + \mathbf{A}^T \mathbf{w} \quad (= 2\mathbf{A} \mathbf{w} \text{ for symmetric } \mathbf{A})$$

Differentiating Matrix Expressions

Rules (Tricks)

(i) Linear Form Expressions: $\nabla_{\mathbf{w}} (\mathbf{c}^T \mathbf{w}) = \mathbf{c}$

$$\begin{aligned}\mathbf{c}^T \mathbf{w} &= \sum_{j=0}^D c_j w_j \\ \frac{\partial (\mathbf{c}^T \mathbf{w})}{\partial w_j} &= c_j, \quad \text{and so} \quad \nabla_{\mathbf{w}} (\mathbf{c}^T \mathbf{w}) = \mathbf{c}\end{aligned}\quad (3)$$

(ii) Quadratic Form Expressions:

$$\nabla_{\mathbf{w}} (\mathbf{w}^T \mathbf{A} \mathbf{w}) = \mathbf{A} \mathbf{w} + \mathbf{A}^T \mathbf{w} \quad (= 2\mathbf{A} \mathbf{w} \text{ for symmetric } \mathbf{A})$$

$$\begin{aligned}\mathbf{w}^T \mathbf{A} \mathbf{w} &= \sum_{i=0}^D \sum_{j=0}^D w_i w_j A_{ij} \\ \frac{\partial (\mathbf{w}^T \mathbf{A} \mathbf{w})}{\partial w_k} &= \sum_{i=0}^D w_i A_{ik} + \sum_{j=0}^D A_{kj} w_j = \mathbf{A}_{[:,k]}^T \mathbf{w} + \mathbf{A}_{[k,:]} \mathbf{w} \\ \nabla_{\mathbf{w}} (\mathbf{w}^T \mathbf{A} \mathbf{w}) &= \mathbf{A}^T \mathbf{w} + \mathbf{A} \mathbf{w}\end{aligned}\quad (4)$$

Deriving the Least Squares Estimate

$$\mathcal{L}(\mathbf{w}) = \frac{1}{2N} \sum_{i=1}^N (\mathbf{x}_i^\top \mathbf{w} - y_i)^2 = \frac{1}{2N} \left(\mathbf{w}^\top (\mathbf{X}^\top \mathbf{X}) \mathbf{w} - 2 \cdot \mathbf{y}^\top \mathbf{X} \mathbf{w} + \mathbf{y}^\top \mathbf{y} \right)$$

Deriving the Least Squares Estimate

$$\mathcal{L}(\mathbf{w}) = \frac{1}{2N} \sum_{i=1}^N (\mathbf{x}_i^\top \mathbf{w} - y_i)^2 = \frac{1}{2N} \left(\mathbf{w}^\top (\mathbf{X}^\top \mathbf{X}) \mathbf{w} - 2 \cdot \mathbf{y}^\top \mathbf{X} \mathbf{w} + \mathbf{y}^\top \mathbf{y} \right)$$

We compute the gradient $\nabla_{\mathbf{w}} \mathcal{L} = \mathbf{0}$ using the matrix differentiation rules,

$$\nabla_{\mathbf{w}} \mathcal{L} = \frac{1}{N} \left((\mathbf{X}^\top \mathbf{X}) \mathbf{w} - \mathbf{X}^\top \mathbf{y} \right)$$

Deriving the Least Squares Estimate

$$\mathcal{L}(\mathbf{w}) = \frac{1}{2N} \sum_{i=1}^N (\mathbf{x}_i^\top \mathbf{w} - y_i)^2 = \frac{1}{2N} \left(\mathbf{w}^\top (\mathbf{X}^\top \mathbf{X}) \mathbf{w} - 2 \cdot \mathbf{y}^\top \mathbf{X} \mathbf{w} + \mathbf{y}^\top \mathbf{y} \right)$$

We compute the gradient $\nabla_{\mathbf{w}} \mathcal{L} = \mathbf{0}$ using the matrix differentiation rules,

$$\nabla_{\mathbf{w}} \mathcal{L} = \frac{1}{N} \left((\mathbf{X}^\top \mathbf{X}) \mathbf{w} - \mathbf{X}^\top \mathbf{y} \right)$$

By setting $\nabla_{\mathbf{w}} \mathcal{L} = \mathbf{0}$ and solving we get,

$$(\mathbf{X}^\top \mathbf{X}) \mathbf{w} = \mathbf{X}^\top \mathbf{y}$$

$$\mathbf{w} = (\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{y} \quad (\text{Assuming inverse exists})$$

Deriving the Least Squares Estimate

$$\mathcal{L}(\mathbf{w}) = \frac{1}{2N} \sum_{i=1}^N (\mathbf{x}_i^\top \mathbf{w} - y_i)^2 = \frac{1}{2N} \left(\mathbf{w}^\top (\mathbf{X}^\top \mathbf{X}) \mathbf{w} - 2 \cdot \mathbf{y}^\top \mathbf{X} \mathbf{w} + \mathbf{y}^\top \mathbf{y} \right)$$

We compute the gradient $\nabla_{\mathbf{w}} \mathcal{L} = \mathbf{0}$ using the matrix differentiation rules,

$$\nabla_{\mathbf{w}} \mathcal{L} = \frac{1}{N} \left((\mathbf{X}^\top \mathbf{X}) \mathbf{w} - \mathbf{X}^\top \mathbf{y} \right)$$

By setting $\nabla_{\mathbf{w}} \mathcal{L} = \mathbf{0}$ and solving we get,

$$(\mathbf{X}^\top \mathbf{X}) \mathbf{w} = \mathbf{X}^\top \mathbf{y}$$

$$\mathbf{w} = (\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{y} \quad (\text{Assuming inverse exists})$$

The predictions made by the model on the data \mathbf{X} are given by

$$\hat{\mathbf{y}} = \mathbf{X} \mathbf{w} = \mathbf{X} (\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{y}$$

For this reason the matrix $\mathbf{X} (\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top$ is called the “hat” matrix

Least Squares Estimate

$$\mathbf{w} = \left(\mathbf{X}^\top \mathbf{X} \right)^{-1} \mathbf{X}^\top \mathbf{y}$$

- ▶ When do we expect $\mathbf{X}^\top \mathbf{X}$ to be invertible?

Least Squares Estimate

$$\mathbf{w} = \left(\mathbf{X}^T \mathbf{X} \right)^{-1} \mathbf{X}^T \mathbf{y}$$

- When do we expect $\mathbf{X}^T \mathbf{X}$ to be invertible?

$$\text{rank}(\mathbf{X}^T \mathbf{X}) = \text{rank}(\mathbf{X}) \leq \min\{D + 1, N\}$$

As $\mathbf{X}^T \mathbf{X}$ is $(D + 1) \times (D + 1)$, invertible if $\text{rank}(\mathbf{X}) = D + 1$

Least Squares Estimate

$$\mathbf{w} = \left(\mathbf{X}^T \mathbf{X} \right)^{-1} \mathbf{X}^T \mathbf{y}$$

- ▶ When do we expect $\mathbf{X}^T \mathbf{X}$ to be invertible?

$$\text{rank}(\mathbf{X}^T \mathbf{X}) = \text{rank}(\mathbf{X}) \leq \min\{D + 1, N\}$$

As $\mathbf{X}^T \mathbf{X}$ is $(D + 1) \times (D + 1)$, invertible if $\text{rank}(\mathbf{X}) = D + 1$

- ▶ What if we use one-hot encoding for a feature like **day**?

Least Squares Estimate

$$\mathbf{w} = \left(\mathbf{X}^T \mathbf{X} \right)^{-1} \mathbf{X}^T \mathbf{y}$$

- ▶ When do we expect $\mathbf{X}^T \mathbf{X}$ to be invertible?

$$\text{rank}(\mathbf{X}^T \mathbf{X}) = \text{rank}(\mathbf{X}) \leq \min\{D + 1, N\}$$

As $\mathbf{X}^T \mathbf{X}$ is $(D + 1) \times (D + 1)$, invertible if $\text{rank}(\mathbf{X}) = D + 1$

- ▶ What if we use one-hot encoding for a feature like **day**?

Suppose $x_{\text{mon}}, \dots, x_{\text{sun}}$ stand for 0-1 valued variables in the one-hot encoding

We always have $x_{\text{mon}} + \dots + x_{\text{sun}} = 1$

This introduces a linear dependence in the columns of \mathbf{X} reducing the rank

In this case, we can drop some features to adjust rank. We'll see alternative approaches later in the course.

Least Squares Estimate

$$\mathbf{w} = \left(\mathbf{X}^T \mathbf{X} \right)^{-1} \mathbf{X}^T \mathbf{y}$$

- ▶ When do we expect $\mathbf{X}^T \mathbf{X}$ to be invertible?

$$\text{rank}(\mathbf{X}^T \mathbf{X}) = \text{rank}(\mathbf{X}) \leq \min\{D + 1, N\}$$

As $\mathbf{X}^T \mathbf{X}$ is $(D + 1) \times (D + 1)$, invertible if $\text{rank}(\mathbf{X}) = D + 1$

- ▶ What if we use one-hot encoding for a feature like **day**?

Suppose $x_{\text{mon}}, \dots, x_{\text{sun}}$ stand for 0-1 valued variables in the one-hot encoding

We always have $x_{\text{mon}} + \dots + x_{\text{sun}} = 1$

This introduces a linear dependence in the columns of \mathbf{X} reducing the rank

In this case, we can drop some features to adjust rank. We'll see alternative approaches later in the course.

- ▶ What is the computational complexity of computing \mathbf{w} ?

Least Squares Estimate

$$\mathbf{w} = (\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{y}$$

- ▶ When do we expect $\mathbf{X}^\top \mathbf{X}$ to be invertible?

$$\text{rank}(\mathbf{X}^\top \mathbf{X}) = \text{rank}(\mathbf{X}) \leq \min\{D + 1, N\}$$

As $\mathbf{X}^\top \mathbf{X}$ is $(D + 1) \times (D + 1)$, invertible if $\text{rank}(\mathbf{X}) = D + 1$

- ▶ What if we use one-hot encoding for a feature like **day**?

Suppose $x_{\text{mon}}, \dots, x_{\text{sun}}$ stand for 0-1 valued variables in the one-hot encoding


We always have $x_{\text{mon}} + \dots + x_{\text{sun}} = 1$


This introduces a linear dependence in the columns of \mathbf{X} reducing the rank


In this case, we can drop some features to adjust rank. We'll see alternative approaches later in the course.


- ▶ What is the computational complexity of computing \mathbf{w} ?

Relatively easy to get $O(D^2 N)$ bound









Recap : Predicting Commute Time

Goal

Recap : Predicting Commute Time

Goal

- ▶ Predict the time taken for commute given distance and day of week

Recap : Predicting Commute Time

Goal

- ▶ Predict the time taken for commute given distance and day of week
- ▶ Do we only wish to make predictions or also suggestions?

Recap : Predicting Commute Time

Goal

- ▶ Predict the time taken for commute given distance and day of week
- ▶ Do we only wish to make predictions or also suggestions?

Model and Choice of Loss Function

Recap : Predicting Commute Time

Goal

- ▶ Predict the time taken for commute given distance and day of week
- ▶ Do we only wish to make predictions or also suggestions?

Model and Choice of Loss Function

- ▶ Use a linear model

$$y = w_0 + w_1x_1 + \cdots + w_Dx_D + \epsilon = \hat{y} + \epsilon$$

- ▶ Minimise average squared error $\frac{1}{2N} \sum (y_i - \hat{y}_i)^2$

Recap : Predicting Commute Time

Goal

- ▶ Predict the time taken for commute given distance and day of week
- ▶ Do we only wish to make predictions or also suggestions?

Model and Choice of Loss Function

- ▶ Use a linear model

$$y = w_0 + w_1x_1 + \cdots + w_Dx_D + \epsilon = \hat{y} + \epsilon$$

- ▶ Minimise average squared error $\frac{1}{2N} \sum (y_i - \hat{y}_i)^2$

Algorithm to Fit Model

Recap : Predicting Commute Time

Goal

- ▶ Predict the time taken for commute given distance and day of week
- ▶ Do we only wish to make predictions or also suggestions?

Model and Choice of Loss Function

- ▶ Use a linear model

$$y = w_0 + w_1x_1 + \cdots + w_Dx_D + \epsilon = \hat{y} + \epsilon$$

- ▶ Minimise average squared error $\frac{1}{2N} \sum (y_i - \hat{y}_i)^2$

Algorithm to Fit Model

- ▶ Simple matrix operations using closed-form solution

Model and Loss Function Choice

“Optimisation” View of Machine Learning

- ▶ Pick model that you expect may fit the data well enough
- ▶ Pick a measure of performance that makes “sense” and can be optimised
- ▶ Run optimisation algorithm to obtain model parameters

Model and Loss Function Choice

“Optimisation” View of Machine Learning

- ▶ Pick model that you expect may fit the data well enough
- ▶ Pick a measure of performance that makes “sense” and can be optimised
- ▶ Run optimisation algorithm to obtain model parameters

Probabilistic View of Machine Learning

- ▶ Pick a model for data and explicitly formulate the deviation (or uncertainty) from the model using the language of probability
- ▶ Use notions from probability to define suitability of various models
- ▶ “Find” the parameters or make predictions on unseen data using these suitability criteria (Frequentist vs Bayesian viewpoints)