

In-Class Lab 4

ECON 4223 (Prof. Tyler Ransom, U of Oklahoma)

September 4, 2018

The purpose of this in-class lab is to further practice your regression skills. The lab should be completed in your group. To get credit, upload your .R script to the appropriate place on Canvas.

For starters

Open up a new R script (named ICL3_XYZ.R, where XYZ are your initials) and add the usual “preamble” to the top:

```
# Add names of group members HERE
library(tidyverse)
library(broom)
library(wooldridge)
```

For this lab, let’s use data on house prices. This is located in the `hprice1` data set in the `wooldridge` package. Each observation is a house.

```
df <- as_tibble(hprice1)
```

Check out what’s in `df` by typing

```
glimpse(hprice1)
```

Multiple Regression

Let’s estimate the following regression model:

$$price = \beta_0 + \beta_1 sqft + \beta_2 bdrms + u$$

where *price* is the house price in thousands of dollars.

The code to do so is:

```
est <- lm(price ~ sqft + bdrms, data=df)
tidy(est)
glance(est)
```

You should get a coefficient of 0.128 on `sqft` and 15.2 on `bdrms`. Interpret these coefficients. (You can type the interpretation as a comment in your .R script.) Do these numbers seem reasonable?

You should get $R^2 = 0.632$. Based on that number, do you think this is a good model of house prices?

Check that the average of the residuals is zero:

```
mean(est$residuals)
```

Adding in non-linearities

The previous regression model had an estimated intercept of -19.3, meaning that a home with no bedrooms and no square footage would be expected to have a sales price of -\$19,300.

To fix this, let's instead use $\log(\text{price})$ as the dependent variable, and let's also add quadratic terms for `sqrft` and `bdrms`.

First, let's use `mutate()` to add these new variables:

```
df <- df %>% mutate(logprice = log(price), sqrftSq = sqrft^2, bdrmSq = bdrms^2)
```

Now run the new model:

```
est <- lm(logprice ~ sqrft + sqrftSq + bdrms + bdrmSq, data=df)
tidy(est)
glance(est)
```

The new coefficients have much smaller magnitudes. Explain why that might be.

The new $R^2 = 0.595$ which is less than 0.632 from before. Does that mean this model is worse?

Using the Frisch-Waugh Theorem to obtain partial effects

Let's experiment with the Frisch-Waugh Theorem, which says:

$$\hat{\beta}_1 = \frac{\sum_{i=1}^N \hat{r}_{i1} y_i}{\sum_{i=1}^N \hat{r}_{i1}^2}$$

where \hat{r}_{i1} is the residual from a regression of x_1 on x_2, \dots, x_k

Let's do this for the model we just ran. First, regress `sqrft` on the other X 's and store the residuals as a new column in `df`.

```
est <- lm(sqrft ~ sqrftSq + bdrms + bdrmSq, data=df)
df <- df %>% mutate(sqrft.resid = est$residuals)
```

Now, if we run a simple regression of `logprice` on `sqrft.resid` we should get the same coefficient as that of `sqrft` in the original regression ($=3.74\text{e-}4$).

```
est <- lm(logprice ~ sqrft.resid, data=df)
tidy(est)
```

Frisch-Waugh by hand

We can also compute the Frisch-Waugh formula by hand:

```
beta1 <- sum(df$sqrft.resid*df$logprice)/sum(df$sqrft.resid^2)
print(beta1)
```

Which indeed gives us what we expected.

References