

## 2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

# 2: Three Different Fourier Transforms

## Signal or Function Representation.

In science and engineering, one often resorts to representation of functions in terms of a *linear combination of elementary basis functions*.

## Signal or Function Representation.

In science and engineering, one often resorts to representation of functions in terms of a linear combination of elementary basis functions.

These are representations of the form,

$$f(x) = \sum_{k \in \mathbb{Z}} c_k B_k(x),$$

where, (a)  $B_k(x)$  are the basis functions and (b)  $c_k$  are expansion coefficients, namely, the amount of  $f$  present in  $B_k$ .

## Signal or Function Representation.

In science and engineering, one often resorts to representation of functions in terms of a *linear combination of elementary basis functions*.

These are representations of the form,

$$f(x) = \sum_{k \in \mathbb{Z}} c_k B_k(x),$$

where, (a)  $B_k(x)$  are the basis functions and (b)  $c_k$  are expansion coefficients, namely, the amount of  $f$  present in  $B_k$ .

Example:

$$f(x) = \underbrace{e^{-x^2}}_{B_1(x)} - 2 \underbrace{x e^{-x^2}}_{B_2(x)} = B_1(x) - 2B_2(x) \equiv B_1(x) - \frac{d}{dx} B_1(x)$$

## Signal or Function Representation.

In science and engineering, one often resorts to representation of functions in terms of a *linear combination of elementary basis functions*.

These are representations of the form,



$$f(x) = \sum_{k \in \mathbb{Z}} c_k B_k(x),$$

where, (a)  $B_k(x)$  are the basis functions and (b)  $c_k$  are expansion coefficients, namely, the amount of  $f$  present in  $B_k$ .

Example:

$$f(x) = \underbrace{e^{-x^2}}_{B_1(x)} - 2 \underbrace{x e^{-x^2}}_{B_2(x)} = B_1(x) - 2B_2(x) \equiv B_1(x) - \frac{d}{dx} B_1(x)$$

combination of two basis



## Signal or Function Representation.

In science and engineering, one often resorts to representation of functions in terms of a *linear combination of elementary basis functions*.

## Signal or Function Representation.

In science and engineering, one often resorts to representation of functions in terms of a *linear combination of elementary basis functions*.

Classical examples include.

- ① Maclaurin Series.

*taylor*

$$f(x) = c_0 + c_1x + c_2x^2 + \dots, \quad c_k = \frac{1}{k!} f^{(k)}(0), k \geq 0.$$

Here the basis functions are  $B = \{1, x, x^2, \dots\}$ .

## Signal or Function Representation.

In science and engineering, one often resorts to representation of functions in terms of a *linear combination of elementary basis functions*.

Classical examples include.

- ① Maclaurin Series.

$$f(x) = c_0 + c_1 x + c_2 x^2 + \cdots, \quad c_k = \frac{1}{k!} f^{(k)}(0), k \geq 0.$$

Here the basis functions are  $B = \{1, x, x^2, \dots\}$ .

- ② Shannon Series.

$$f(x) = c_0 \operatorname{sinc}(x) + c_{\pm 1} \operatorname{sinc}(x \pm 1) + c_{\pm 2} \operatorname{sinc}(x \pm 2) + \cdots, \quad c_k = f(k), k \in \mathbb{Z}$$

where the basis functions are  $B = \{\operatorname{sinc}(x - k)\}_{k \in \mathbb{Z}}$ .

## Signal or Function Representation.

In science and engineering, one often resorts to representation of functions in terms of a *linear combination of elementary basis functions*.

Classical examples include.

- ① Maclaurin Series.

$$f(x) = c_0 + c_1 x + c_2 x^2 + \cdots, \quad c_k = \frac{1}{k!} f^{(k)}(0), k \geq 0.$$

Here the basis functions are  $B = \{1, x, x^2, \dots\}$ .

- ② Shannon Series.

$$f(x) = c_0 \operatorname{sinc}(x) + c_{\pm 1} \operatorname{sinc}(x \pm 1) + c_{\pm 2} \operatorname{sinc}(x \pm 2) + \cdots, \quad c_k = f(k), k \in \mathbb{Z}$$

where the basis functions are  $B = \{\operatorname{sinc}(x - k)\}_{k \in \mathbb{Z}}$ .

- ③ Fourier Series.

$$f(x) = c_0 + c_{\pm 1} \exp(\pm j\omega_0 t) + c_{\pm 2} \exp(\pm j2\omega_0 t) + \cdots, \quad c_k = \frac{1}{T} \int_0^T f(x) e^{-j\frac{2\pi}{T} kt} dt, k \in \mathbb{Z}$$

where the basis functions are  $B = \left\{ e^{j\frac{2\pi}{T} kt} \right\}_{k \in \mathbb{Z}}$ .

## Signal or Function Representation.

In the area of signal processing,

- Shannon Series (Digital Representation)
- Fourier Series (Periodic Representation)





are the most important representations.

## Signal or Function Representation.

In the area of signal processing,

- Shannon Series (Digital Representation)
- Fourier Series (Periodic Representation)

are the most important representations.



## Signal Representation in Orthogonal Basis.


When the basis functions are orthogonal, that is,

$$\langle B_m, B_n \rangle = \int B_m(x) B_n^*(x) dx = \boxed{\delta_{m-n}}$$

then the function can be represented by the expansion,

$$f(x) = \sum_k \langle f, B_k \rangle B_k(x).$$

This simple yet powerful method is at the heart of the Shannon's Sampling Series as well as the Fourier Series.



## Fourier Series

In the case of Fourier Series,

- ① The basis functions are defined by,

$$B_k = \left\{ e^{j \frac{2\pi}{T} kt} \right\}_{k \in \mathbb{Z}}.$$

## Fourier Series

In the case of Fourier Series,

- ① The basis functions are defined by,

$$B_k = \left\{ e^{j \frac{2\pi}{T} kt} \right\}_{k \in \mathbb{Z}}.$$

- ② The basis functions are orthogonal. This is because,

$$\langle B_m, B_n \rangle = \int B_m(x) B_n^*(x) dx = \frac{1}{T} \int_0^T e^{j \frac{2\pi}{T} mx} e^{-j \frac{2\pi}{T} nx} dx = \frac{1}{T} \int_0^T e^{j \frac{2\pi}{T} (m-n)x} dx = \delta_{m-n}.$$

## Fourier Series

In the case of Fourier Series,

- ① The basis functions are defined by,

$$B_k = \left\{ e^{j \frac{2\pi}{T} kt} \right\}_{k \in \mathbb{Z}}.$$

- ② The basis functions are orthogonal. This is because,

$$\langle B_m, B_n \rangle = \int B_m(x) B_n^*(x) dx = \frac{1}{T} \int_0^T e^{j \frac{2\pi}{T} mx} e^{-j \frac{2\pi}{T} nx} dx = \frac{1}{T} \int_0^T e^{j \frac{2\pi}{T} (m-n)x} dx = \delta_{m-n}.$$

- ③ Because of this, we can write,

$$f(x) = \sum_k \langle f, B_k \rangle B_k(x).$$

where, the Fourier Series coefficients are given by,

$$\langle f, B_k \rangle = \int f(x) B_k^*(x) dx = \int f(x) e^{-j \frac{2\pi}{T} kx} dx = \hat{f}\left(\frac{2\pi}{T} k\right).$$

## 2: Three Different Fourier Transforms

---

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

# Fourier Transforms

---

## Three different Fourier Transforms:

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

# Fourier Transforms

Three different Fourier Transforms:

- CTFT (Continuous-Time Fourier Transform):  $x(t) \rightarrow X(j\Omega)$

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

# Fourier Transforms

Three different Fourier Transforms:

- CTFT (Continuous-Time Fourier Transform):  $x(t) \rightarrow X(j\Omega)$
- DTFT (Discrete-Time Fourier Transform):  $x[n] \rightarrow X(e^{j\omega})$

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

# Fourier Transforms

## Three different Fourier Transforms:

- CTFT (Continuous-Time Fourier Transform):  $x(t) \rightarrow X(j\Omega)$
- DTFT (Discrete-Time Fourier Transform):  $x[n] \rightarrow X(e^{j\omega})$
- DFT a.k.a. FFT (Discrete Fourier Transform):  $x[n] \rightarrow X[k]$

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

# Fourier Transforms

Three different Fourier Transforms:

- CTFT (Continuous-Time Fourier Transform):  $x(t) \rightarrow X(j\Omega)$
- DTFT (Discrete-Time Fourier Transform):  $x[n] \rightarrow X(e^{j\omega})$
- DFT a.k.a. FFT (Discrete Fourier Transform):  $x[n] \rightarrow X[k]$

	Forward Transform	Inverse Transform
CTFT	$X(j\Omega) = \int_{-\infty}^{\infty} x(t)e^{-j\Omega t} dt$	$x(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} X(j\Omega)e^{j\Omega t} d\Omega$

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

# Fourier Transforms

Three different Fourier Transforms:

- CTFT (Continuous-Time Fourier Transform):  $x(t) \rightarrow X(j\Omega)$
- DTFT (Discrete-Time Fourier Transform):  $x[n] \rightarrow X(e^{j\omega})$
- DFT a.k.a. FFT (Discrete Fourier Transform):  $x[n] \rightarrow X[k]$

	Forward Transform	Inverse Transform
CTFT	$X(j\Omega) = \int_{-\infty}^{\infty} x(t)e^{-j\Omega t} dt$	$x(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} X(j\Omega)e^{j\Omega t} d\Omega$
DTFT	$X(e^{j\omega}) = \sum_{-\infty}^{\infty} x[n]e^{-j\omega n}$	$x[n] = \frac{1}{2\pi} \int_{-\pi}^{\pi} X(e^{j\omega})e^{j\omega n} d\omega$

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

# Fourier Transforms

Three different Fourier Transforms:

- CTFT (Continuous-Time Fourier Transform):  $x(t) \rightarrow X(j\Omega)$
- DTFT (Discrete-Time Fourier Transform):  $x[n] \rightarrow X(e^{j\omega})$
- DFT a.k.a. FFT (Discrete Fourier Transform):  $x[n] \rightarrow X[k]$

Forward Transform	Inverse Transform
CTFT $X(j\Omega) = \int_{-\infty}^{\infty} x(t)e^{-j\Omega t} dt$	$x(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} X(j\Omega)e^{j\Omega t} d\Omega$
DTFT $X(e^{j\omega}) = \sum_{-\infty}^{\infty} x[n]e^{-j\omega n}$	$x[n] = \frac{1}{2\pi} \int_{-\pi}^{\pi} X(e^{j\omega})e^{j\omega n} d\omega$

We use  $\Omega$  for “real” and  $\omega = \Omega T$  for “normalized” angular frequency.  
Nyquist frequency is at  $\Omega_{\text{Nyq}} = 2\pi \frac{f_s}{2} = \frac{\pi}{T}$  and  $\omega_{\text{Nyq}} = \pi$ .

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

# Fourier Transforms

Three different Fourier Transforms:

- CTFT (Continuous-Time Fourier Transform):  $x(t) \rightarrow X(j\Omega)$
- DTFT (Discrete-Time Fourier Transform):  $x[n] \rightarrow X(e^{j\omega})$
- DFT a.k.a. FFT (Discrete Fourier Transform):  $x[n] \rightarrow X[k]$

Forward Transform	Inverse Transform
CTFT $X(j\Omega) = \int_{-\infty}^{\infty} x(t)e^{-j\Omega t} dt$	$x(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} X(j\Omega)e^{j\Omega t} d\Omega$
DTFT $X(e^{j\omega}) = \sum_{-\infty}^{\infty} x[n]e^{-j\omega n}$	$x[n] = \frac{1}{2\pi} \int_{-\pi}^{\pi} X(e^{j\omega})e^{j\omega n} d\omega$
DFT $X[k] = \sum_0^{N-1} x[n]e^{-j2\pi \frac{kn}{N}}$	$x[n] = \frac{1}{N} \sum_0^{N-1} X[k]e^{j2\pi \frac{kn}{N}}$

We use  $\Omega$  for “real” and  $\omega = \Omega T$  for “normalized” angular frequency.  
Nyquist frequency is at  $\Omega_{\text{Nyq}} = 2\pi \frac{f_s}{2} = \frac{\pi}{T}$  and  $\omega_{\text{Nyq}} = \pi$ .

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

# Fourier Transforms

Three different Fourier Transforms:

- CTFT (Continuous-Time Fourier Transform):  $x(t) \rightarrow X(j\Omega)$
- DTFT (Discrete-Time Fourier Transform):  $x[n] \rightarrow X(e^{j\omega})$
- DFT a.k.a. FFT (Discrete Fourier Transform):  $x[n] \rightarrow X[k]$

	Forward Transform	Inverse Transform
CTFT	$X(j\Omega) = \int_{-\infty}^{\infty} x(t)e^{-j\Omega t} dt$	$x(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} X(j\Omega)e^{j\Omega t} d\Omega$
DTFT	$X(e^{j\omega}) = \sum_{-\infty}^{\infty} x[n]e^{-j\omega n}$	$x[n] = \frac{1}{2\pi} \int_{-\pi}^{\pi} X(e^{j\omega})e^{j\omega n} d\omega$
DFT	$X[k] = \sum_0^{N-1} x[n]e^{-j2\pi \frac{kn}{N}}$	$x[n] = \frac{1}{N} \sum_0^{N-1} X[k]e^{j2\pi \frac{kn}{N}}$

We use  $\Omega$  for “real” and  $\omega = \Omega T$  for “normalized” angular frequency.  
Nyquist frequency is at  $\Omega_{\text{Nyq}} = 2\pi \frac{f_s}{2} = \frac{\pi}{T}$  and  $\omega_{\text{Nyq}} = \pi$ .

For “power signals” (energy  $\propto$  duration), CTFT & DTFT are unbounded.  
Fix this by normalizing:

$$X(j\Omega) = \lim_{A \rightarrow \infty} \frac{1}{2A} \int_{-A}^A x(t)e^{-j\Omega t} dt$$
$$X(e^{j\omega}) = \lim_{A \rightarrow \infty} \frac{1}{2A+1} \sum_{-A}^A x[n]e^{-j\omega n}$$

2: Three Different Fourier Transforms

- Fourier Transforms
- **Convergence of DTFT**
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## Convergence of DTFT

DTFT:  $X(e^{j\omega}) = \sum_{-\infty}^{\infty} x[n]e^{-j\omega n}$  does not converge for all  $x[n]$ .

Consider the finite sum:  $X_K(e^{j\omega}) = \sum_{-K}^K x[n]e^{-j\omega n}$

- Fourier Transforms
- **Convergence of DTFT**
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## Convergence of DTFT

DTFT:  $X(e^{j\omega}) = \sum_{-\infty}^{\infty} x[n]e^{-j\omega n}$  does not converge for all  $x[n]$ .

Consider the finite sum:  $X_K(e^{j\omega}) = \sum_{-K}^K x[n]e^{-j\omega n}$

**Strong Convergence:**

$x[n]$  absolutely summable  $\Rightarrow X(e^{j\omega})$  converges **uniformly**

- Fourier Transforms
- **Convergence of DTFT**
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## Convergence of DTFT

DTFT:  $X(e^{j\omega}) = \sum_{-\infty}^{\infty} x[n]e^{-j\omega n}$  does not converge for all  $x[n]$ .

Consider the finite sum:  $X_K(e^{j\omega}) = \sum_{-K}^K x[n]e^{-j\omega n}$

**Strong Convergence:**

$x[n]$  absolutely summable  $\Rightarrow X(e^{j\omega})$  converges **uniformly**  
 $\sum_{-\infty}^{\infty} |x[n]| < \infty \Rightarrow \sup_{\omega} |X(e^{j\omega}) - X_K(e^{j\omega})| \xrightarrow[K \rightarrow \infty]{} 0$

- Fourier Transforms
- **Convergence of DTFT**
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## Convergence of DTFT

DTFT:  $X(e^{j\omega}) = \sum_{-\infty}^{\infty} x[n]e^{-j\omega n}$  does not converge for all  $x[n]$ .

Consider the finite sum:  $X_K(e^{j\omega}) = \sum_{-K}^K x[n]e^{-j\omega n}$

**Strong Convergence:**

$x[n]$  absolutely summable  $\Rightarrow X(e^{j\omega})$  converges **uniformly**  
 $\sum_{-\infty}^{\infty} |x[n]| < \infty \Rightarrow \sup_{\omega} |X(e^{j\omega}) - X_K(e^{j\omega})| \xrightarrow[K \rightarrow \infty]{} 0$

**Weaker convergence:**

$x[n]$  finite energy  $\Rightarrow X(e^{j\omega})$  converges in **mean square**

- Fourier Transforms
- **Convergence of DTFT**
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## Convergence of DTFT

DTFT:  $X(e^{j\omega}) = \sum_{-\infty}^{\infty} x[n]e^{-j\omega n}$  does not converge for all  $x[n]$ .

Consider the finite sum:  $X_K(e^{j\omega}) = \sum_{-K}^K x[n]e^{-j\omega n}$

**Strong Convergence:**

$x[n]$  absolutely summable  $\Rightarrow X(e^{j\omega})$  converges **uniformly**

$$\sum_{-\infty}^{\infty} |x[n]| < \infty \Rightarrow \sup_{\omega} |X(e^{j\omega}) - X_K(e^{j\omega})| \xrightarrow[K \rightarrow \infty]{} 0$$

**Weaker convergence:**

$x[n]$  finite energy  $\Rightarrow X(e^{j\omega})$  converges in **mean square**

$$\sum_{-\infty}^{\infty} |x[n]|^2 < \infty \Rightarrow \frac{1}{2\pi} \int_{-\pi}^{\pi} |X(e^{j\omega}) - X_K(e^{j\omega})|^2 d\omega \xrightarrow[K \rightarrow \infty]{} 0$$

# Convergence of DTFT

2: Three Different Fourier Transforms

- Fourier Transforms
- **Convergence of DTFT**
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

DTFT:  $X(e^{j\omega}) = \sum_{-\infty}^{\infty} x[n]e^{-j\omega n}$  does not converge for all  $x[n]$ .

Consider the finite sum:  $X_K(e^{j\omega}) = \sum_{-K}^K x[n]e^{-j\omega n}$

**Strong Convergence:**

$x[n]$  absolutely summable  $\Rightarrow X(e^{j\omega})$  converges **uniformly**


$$\sum_{-\infty}^{\infty} |x[n]| < \infty \Rightarrow \sup_{\omega} |X(e^{j\omega}) - X_K(e^{j\omega})| \xrightarrow[K \rightarrow \infty]{} 0$$

**Weaker convergence:**

$x[n]$  finite energy  $\Rightarrow X(e^{j\omega})$  converges in **mean square**

$$\sum_{-\infty}^{\infty} |x[n]|^2 < \infty \Rightarrow \frac{1}{2\pi} \int_{-\pi}^{\pi} |X(e^{j\omega}) - X_K(e^{j\omega})|^2 d\omega \xrightarrow[K \rightarrow \infty]{} 0$$

**Example:**  $x[n] = \frac{\sin 0.5\pi n}{\pi n}$



# Convergence of DTFT

2: Three Different Fourier Transforms

- Fourier Transforms
- **Convergence of DTFT**
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

DTFT:  $X(e^{j\omega}) = \sum_{-\infty}^{\infty} x[n]e^{-j\omega n}$  does not converge for all  $x[n]$ .

Consider the finite sum:  $X_K(e^{j\omega}) = \sum_{-K}^K x[n]e^{-j\omega n}$

**Strong Convergence:**

$x[n]$  absolutely summable  $\Rightarrow X(e^{j\omega})$  converges **uniformly**


$$\sum_{-\infty}^{\infty} |x[n]| < \infty \Rightarrow \sup_{\omega} |X(e^{j\omega}) - X_K(e^{j\omega})| \xrightarrow[K \rightarrow \infty]{} 0$$

**Weaker convergence:**

$x[n]$  finite energy  $\Rightarrow X(e^{j\omega})$  converges in **mean square**

$$\sum_{-\infty}^{\infty} |x[n]|^2 < \infty \Rightarrow \frac{1}{2\pi} \int_{-\pi}^{\pi} |X(e^{j\omega}) - X_K(e^{j\omega})|^2 d\omega \xrightarrow[K \rightarrow \infty]{} 0$$

**Example:**  $x[n] = \frac{\sin 0.5\pi n}{\pi n}$



**Gibbs phenomenon:**

Converges at each  $\omega$  as  $K \rightarrow \infty$  but peak error does not get smaller.

- Fourier Transforms
- Convergence of DTFT
- **DTFT Properties**
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## DTFT Properties

$$\text{DTFT: } X(e^{j\omega}) = \sum_{-\infty}^{\infty} x[n]e^{-j\omega n}$$

2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## DTFT Properties

$$\text{DTFT: } X(e^{j\omega}) = \sum_{-\infty}^{\infty} x[n]e^{-j\omega n}$$

- DTFT is periodic in  $\omega$ :  $X(e^{j(\omega+2m\pi)}) = X(e^{j\omega})$  for integer  $m$ .

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## DTFT Properties

$$\text{DTFT: } X(e^{j\omega}) = \sum_{-\infty}^{\infty} x[n]e^{-j\omega n}$$

- DTFT is periodic in  $\omega$ :  $X(e^{j(\omega+2m\pi)}) = X(e^{j\omega})$  for integer  $m$ .
- DTFT is the  $z$ -Transform evaluated at the point  $e^{j\omega}$ :

$$X(z) = \sum_{-\infty}^{\infty} x[n]z^{-n}$$

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## DTFT Properties

$$\text{DTFT: } X(e^{j\omega}) = \sum_{-\infty}^{\infty} x[n]e^{-j\omega n}$$

- DTFT is **periodic** in  $\omega$ :  $X(e^{j(\omega+2m\pi)}) = X(e^{j\omega})$  for integer  $m$ .
- DTFT is the  **$z$ -Transform** evaluated at the point  $e^{j\omega}$ :

$$X(z) = \sum_{-\infty}^{\infty} x[n]z^{-n}$$

DTFT converges iff the ROC includes  $|z| = 1$ .

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## DTFT Properties

$$\text{DTFT: } X(e^{j\omega}) = \sum_{-\infty}^{\infty} x[n]e^{-j\omega n}$$

- DTFT is **periodic** in  $\omega$ :  $X(e^{j(\omega+2m\pi)}) = X(e^{j\omega})$  for integer  $m$ .

- DTFT is the  **$z$ -Transform** evaluated at the point  $e^{j\omega}$ :

$$X(z) = \sum_{-\infty}^{\infty} x[n]z^{-n}$$

DTFT converges iff the ROC includes  $|z| = 1$ .

- DTFT is the same as the CTFT of a signal comprising **impulses at the sample times** (Dirac  $\delta$  functions) of appropriate heights:

$$x_{\delta}(t) = \sum x[n]\delta(t - nT)$$

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## DTFT Properties

$$\text{DTFT: } X(e^{j\omega}) = \sum_{n=-\infty}^{\infty} x[n]e^{-j\omega n}$$

- DTFT is **periodic** in  $\omega$ :  $X(e^{j(\omega+2m\pi)}) = X(e^{j\omega})$  for integer  $m$ .

- DTFT is the  **$z$ -Transform** evaluated at the point  $e^{j\omega}$ :

$$X(z) = \sum_{n=-\infty}^{\infty} x[n]z^{-n}$$

DTFT converges iff the ROC includes  $|z| = 1$ .

- DTFT is the same as the CTFT of a signal comprising **impulses at the sample times** (Dirac  $\delta$  functions) of appropriate heights:

$$x_{\delta}(t) = \sum x[n]\underbrace{\delta(t - nT)}_{\text{Sampling}}$$

Equivalent to multiplying a continuous  $x(t)$  by an impulse train.

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## DTFT Properties

$$\text{DTFT: } X(e^{j\omega}) = \sum_{-\infty}^{\infty} x[n]e^{-j\omega n}$$

- DTFT is **periodic** in  $\omega$ :  $X(e^{j(\omega+2m\pi)}) = X(e^{j\omega})$  for integer  $m$ .

- DTFT is the  **$z$ -Transform** evaluated at the point  $e^{j\omega}$ :

$$X(z) = \sum_{-\infty}^{\infty} x[n]z^{-n}$$

DTFT converges iff the ROC includes  $|z| = 1$ .

- DTFT is the same as the CTFT of a signal comprising **impulses at the sample times** (Dirac  $\delta$  functions) of appropriate heights:

$$x_{\delta}(t) = \sum x[n]\delta(t - nT) = x(t) \times \sum_{-\infty}^{\infty} \delta(t - nT)$$

Equivalent to multiplying a continuous  $x(t)$  by an impulse train.

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## DTFT Properties

$$\text{DTFT: } X(e^{j\omega}) = \sum_{-\infty}^{\infty} x[n]e^{-j\omega n}$$

- DTFT is **periodic** in  $\omega$ :  $X(e^{j(\omega+2m\pi)}) = X(e^{j\omega})$  for integer  $m$ .

- DTFT is the  **$z$ -Transform** evaluated at the point  $e^{j\omega}$ :

$$X(z) = \sum_{-\infty}^{\infty} x[n]z^{-n}$$

DTFT converges iff the ROC includes  $|z| = 1$ .

- DTFT is the same as the CTFT of a signal comprising **impulses at the sample times** (Dirac  $\delta$  functions) of appropriate heights:

$$x_{\delta}(t) = \sum x[n]\delta(t - nT) = x(t) \times \sum_{-\infty}^{\infty} \delta(t - nT)$$

Equivalent to multiplying a continuous  $x(t)$  by an impulse train.

$$\text{Proof: } X(e^{j\omega}) = \sum_{-\infty}^{\infty} x[n]e^{-j\omega n}$$

- Fourier Transforms
- Convergence of DTFT
- **DTFT Properties**
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## DTFT Properties

$$\text{DTFT: } X(e^{j\omega}) = \sum_{n=-\infty}^{\infty} x[n]e^{-j\omega n}$$

- DTFT is **periodic** in  $\omega$ :  $X(e^{j(\omega+2m\pi)}) = X(e^{j\omega})$  for integer  $m$ .

- DTFT is the  **$z$ -Transform** evaluated at the point  $e^{j\omega}$ :

$$X(z) = \sum_{n=-\infty}^{\infty} x[n]z^{-n}$$

DTFT converges iff the ROC includes  $|z| = 1$ .

- DTFT is the same as the CTFT of a signal comprising **impulses at the sample times** (Dirac  $\delta$  functions) of appropriate heights:

$$x_{\delta}(t) = \sum x[n]\delta(t - nT) = x(t) \times \sum_{n=-\infty}^{\infty} \delta(t - nT)$$

Equivalent to multiplying a continuous  $x(t)$  by an impulse train.

$$\text{Proof: } X(e^{j\omega}) = \sum_{n=-\infty}^{\infty} x[n]e^{-j\omega n}$$

$$\sum_{n=-\infty}^{\infty} x[n] \int_{-\infty}^{\infty} \delta(t - nT)e^{-j\omega \frac{t}{T}} dt$$

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## DTFT Properties

$$\text{DTFT: } X(e^{j\omega}) = \sum_{n=-\infty}^{\infty} x[n]e^{-j\omega n}$$

- DTFT is **periodic** in  $\omega$ :  $X(e^{j(\omega+2m\pi)}) = X(e^{j\omega})$  for integer  $m$ .

- DTFT is the  **$z$ -Transform** evaluated at the point  $e^{j\omega}$ :

$$X(z) = \sum_{n=-\infty}^{\infty} x[n]z^{-n}$$

DTFT converges iff the ROC includes  $|z| = 1$ .

- DTFT is the same as the CTFT of a signal comprising **impulses at the sample times** (Dirac  $\delta$  functions) of appropriate heights:

$$x_{\delta}(t) = \sum x[n]\delta(t - nT) = x(t) \times \sum_{n=-\infty}^{\infty} \delta(t - nT)$$

Equivalent to multiplying a continuous  $x(t)$  by an impulse train.

$$\text{Proof: } X(e^{j\omega}) = \sum_{n=-\infty}^{\infty} x[n]e^{-j\omega n}$$

$$\sum_{n=-\infty}^{\infty} x[n] \int_{-\infty}^{\infty} \delta(t - nT)e^{-j\omega \frac{t}{T}} dt$$

$$\stackrel{(i)}{=} \int_{-\infty}^{\infty} \sum_{n=-\infty}^{\infty} x[n]\delta(t - nT)e^{-j\omega \frac{t}{T}} dt$$

(i) OK if  $\sum_{n=-\infty}^{\infty} |x[n]| < \infty$ .

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## DTFT Properties

$$\text{DTFT: } X(e^{j\omega}) = \sum_{n=-\infty}^{\infty} x[n]e^{-j\omega n}$$

- DTFT is **periodic** in  $\omega$ :  $X(e^{j(\omega+2m\pi)}) = X(e^{j\omega})$  for integer  $m$ .

- DTFT is the  **$z$ -Transform** evaluated at the point  $e^{j\omega}$ :

$$X(z) = \sum_{n=-\infty}^{\infty} x[n]z^{-n}$$

DTFT converges iff the ROC includes  $|z| = 1$ .

- DTFT is the same as the CTFT of a signal comprising **impulses at the sample times** (Dirac  $\delta$  functions) of appropriate heights:

$$x_{\delta}(t) = \sum x[n]\delta(t - nT) = x(t) \times \sum_{n=-\infty}^{\infty} \delta(t - nT)$$

Equivalent to multiplying a continuous  $x(t)$  by an impulse train.

$$\begin{aligned} \text{Proof: } X(e^{j\omega}) &= \sum_{n=-\infty}^{\infty} x[n]e^{-j\omega n} \\ &= \sum_{n=-\infty}^{\infty} x[n] \int_{-\infty}^{\infty} \delta(t - nT)e^{-j\omega \frac{t}{T}} dt \\ &\stackrel{(i)}{=} \int_{-\infty}^{\infty} \sum_{n=-\infty}^{\infty} x[n]\delta(t - nT)e^{-j\omega \frac{t}{T}} dt \\ &\stackrel{(ii)}{=} \int_{-\infty}^{\infty} x_{\delta}(t)e^{-j\omega t} dt \end{aligned}$$

(i) OK if  $\sum_{n=-\infty}^{\infty} |x[n]| < \infty$ . (ii) use  $\omega = \Omega T$ .

2: Three Different Fourier  
Transforms

---

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- **DFT Properties**
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## DFT Properties

---

$$\text{DFT: } X[k] = \sum_0^{N-1} x[n]e^{-j2\pi \frac{kn}{N}}$$

## 2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- **DFT Properties**
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

# DFT Properties

$$\text{DFT: } X[k] = \sum_0^{N-1} x[n]e^{-j2\pi \frac{kn}{N}}$$

Case 1:  $x[n] = 0$  for  $n \notin [0, N - 1]$

DFT is the same as DTFT at  $\omega_k = \frac{2\pi}{N}k$ .

## 2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- **DFT Properties**
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

# DFT Properties

$$\text{DFT: } X[k] = \sum_0^{N-1} x[n]e^{-j2\pi \frac{kn}{N}}$$

$$\text{DTFT: } X(e^{j\omega}) = \sum_{-\infty}^{\infty} x[n]e^{-j\omega n}$$

Case 1:  $x[n] = 0$  for  $n \notin [0, N - 1]$

DFT is the same as DTFT at  $\omega_k = \frac{2\pi}{N}k$ .

2: Three Different Fourier Transforms

---

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- **DFT Properties**
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## DFT Properties

---

$$\text{DFT: } X[k] = \sum_0^{N-1} x[n]e^{-j2\pi \frac{kn}{N}}$$

$$\text{DTFT: } X(e^{j\omega}) = \sum_{-\infty}^{\infty} x[n]e^{-j\omega n}$$

Case 1:  $x[n] = 0$  for  $n \notin [0, N - 1]$

DFT is the same as DTFT at  $\omega_k = \frac{2\pi}{N}k$ .

The  $\{\omega_k\}$  are uniformly spaced from  $\omega = 0$  to  $\omega = 2\pi \frac{N-1}{N}$ .

## 2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- **DFT Properties**
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

# DFT Properties

$$\text{DFT: } X[k] = \sum_0^{N-1} x[n]e^{-j2\pi \frac{kn}{N}}$$

$$\text{DTFT: } X(e^{j\omega}) = \sum_{-\infty}^{\infty} x[n]e^{-j\omega n}$$

Case 1:  $x[n] = 0$  for  $n \notin [0, N - 1]$

DFT is the same as DTFT at  $\omega_k = \frac{2\pi}{N}k$ .

The  $\{\omega_k\}$  are uniformly spaced from  $\omega = 0$  to  $\omega = 2\pi \frac{N-1}{N}$ .  
DFT is the  $z$ -Transform evaluated at  $N$  equally spaced points around the unit circle beginning at  $z = 1$ .

## 2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- **DFT Properties**
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

# DFT Properties

$$\text{DFT: } X[k] = \sum_0^{N-1} x[n]e^{-j2\pi \frac{kn}{N}}$$

$$\text{DTFT: } X(e^{j\omega}) = \sum_{-\infty}^{\infty} x[n]e^{-j\omega n}$$

Case 1:  $x[n] = 0$  for  $n \notin [0, N - 1]$

DFT is the same as DTFT at  $\omega_k = \frac{2\pi}{N}k$ .

The  $\{\omega_k\}$  are uniformly spaced from  $\omega = 0$  to  $\omega = 2\pi \frac{N-1}{N}$ .

DFT is the  $z$ -Transform evaluated at  $N$  equally spaced points around the unit circle beginning at  $z = 1$ .

Case 2:  $x[n]$  is periodic with period  $N$

DFT equals the normalized DTFT

## 2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- **DFT Properties**
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

# DFT Properties

$$\text{DFT: } X[k] = \sum_0^{N-1} x[n]e^{-j2\pi \frac{kn}{N}}$$

$$\text{DTFT: } X(e^{j\omega}) = \sum_{-\infty}^{\infty} x[n]e^{-j\omega n}$$

Case 1:  $x[n] = 0$  for  $n \notin [0, N - 1]$

DFT is the same as DTFT at  $\omega_k = \frac{2\pi}{N}k$ .

The  $\{\omega_k\}$  are uniformly spaced from  $\omega = 0$  to  $\omega = 2\pi \frac{N-1}{N}$ .

DFT is the  $z$ -Transform evaluated at  $N$  equally spaced points around the unit circle beginning at  $z = 1$ .

Case 2:  $x[n]$  is periodic with period  $N$

DFT equals the normalized DTFT

$$X[k] = \lim_{K \rightarrow \infty} \frac{N}{2K+1} \times X_K(e^{j\omega_k})$$

## 2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- **DFT Properties**
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

# DFT Properties

$$\text{DFT: } X[k] = \sum_0^{N-1} x[n] e^{-j2\pi \frac{kn}{N}}$$

$$\text{DTFT: } X(e^{j\omega}) = \sum_{-\infty}^{\infty} x[n] e^{-j\omega n}$$

Case 1:  $x[n] = 0$  for  $n \notin [0, N - 1]$

DFT is the same as DTFT at  $\omega_k = \frac{2\pi}{N}k$ .

The  $\{\omega_k\}$  are uniformly spaced from  $\omega = 0$  to  $\omega = 2\pi \frac{N-1}{N}$ .

DFT is the  $z$ -Transform evaluated at  $N$  equally spaced points around the unit circle beginning at  $z = 1$ .

Case 2:  $x[n]$  is periodic with period  $N$

DFT equals the normalized DTFT

$$X[k] = \lim_{K \rightarrow \infty} \frac{N}{2K+1} \times X_K(e^{j\omega_k})$$

$$\text{where } X_K(e^{j\omega}) = \sum_{-K}^K x[n] e^{-j\omega n}$$

# Symmetries

2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- **Symmetries**
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

If  $x[n]$  has a special property then  $X(e^{j\omega})$  and  $X[k]$  will have corresponding properties as shown in the table (and vice versa):

One domain	Other domain
Discrete	Periodic
Symmetric	Symmetric
Antisymmetric	Antisymmetric
Real	Conjugate Symmetric
Imaginary	Conjugate Antisymmetric
Real + Symmetric	Real + Symmetric
Real + Antisymmetric	Imaginary + Antisymmetric

# Symmetries

2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- **Symmetries**
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

If  $x[n]$  has a special property then  $X(e^{j\omega})$  and  $X[k]$  will have corresponding properties as shown in the table (and vice versa):

One domain	Other domain
Discrete	Periodic
Symmetric	Symmetric
Antisymmetric	Antisymmetric
Real	Conjugate Symmetric
Imaginary	Conjugate Antisymmetric
Real + Symmetric	Real + Symmetric
Real + Antisymmetric	Imaginary + Antisymmetric

**Symmetric:**  $x[n] = x[-n]$

# Symmetries

2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- **Symmetries**
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

If  $x[n]$  has a special property then  $X(e^{j\omega})$  and  $X[k]$  will have corresponding properties as shown in the table (and vice versa):

One domain	Other domain
Discrete	Periodic
Symmetric	Symmetric
Antisymmetric	Antisymmetric
Real	Conjugate Symmetric
Imaginary	Conjugate Antisymmetric
Real + Symmetric	Real + Symmetric
Real + Antisymmetric	Imaginary + Antisymmetric

**Symmetric:**  $x[n] = x[-n]$   
 $X(e^{j\omega}) = X(e^{-j\omega})$

# Symmetries

2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- **Symmetries**
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

If  $x[n]$  has a special property then  $X(e^{j\omega})$  and  $X[k]$  will have corresponding properties as shown in the table (and vice versa):

One domain	Other domain
Discrete	Periodic
Symmetric	Symmetric
Antisymmetric	Antisymmetric
Real	Conjugate Symmetric
Imaginary	Conjugate Antisymmetric
Real + Symmetric	Real + Symmetric
Real + Antisymmetric	Imaginary + Antisymmetric

**Symmetric:**  $x[n] = x[-n]$   
 $X(e^{j\omega}) = X(e^{-j\omega})$   
 $X[k] = X[(-k)_{\text{mod } N}] = X[N - k]$  for  $k > 0$

# Symmetries

2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- **Symmetries**
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

If  $x[n]$  has a special property then  $X(e^{j\omega})$  and  $X[k]$  will have corresponding properties as shown in the table (and vice versa):

One domain	Other domain
Discrete	Periodic
Symmetric	Symmetric
Antisymmetric	Antisymmetric
Real	Conjugate Symmetric
Imaginary	Conjugate Antisymmetric
Real + Symmetric	Real + Symmetric
Real + Antisymmetric	Imaginary + Antisymmetric

**Symmetric:**  $x[n] = x[-n]$   
 $X(e^{j\omega}) = X(e^{-j\omega})$   
 $X[k] = X[(-k) \bmod N] = X[N - k]$  for  $k > 0$

**Conjugate Symmetric:**  $x[n] = x^*[-n]$

## 2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- **Symmetries**
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

# Symmetries

If  $x[n]$  has a special property then  $X(e^{j\omega})$  and  $X[k]$  will have corresponding properties as shown in the table (and vice versa):

One domain	Other domain
Discrete	Periodic
Symmetric	Symmetric
Antisymmetric	Antisymmetric
Real	Conjugate Symmetric
Imaginary	Conjugate Antisymmetric
Real + Symmetric	Real + Symmetric
Real + Antisymmetric	Imaginary + Antisymmetric

**Symmetric:**  $x[n] = x[-n]$   
 $X(e^{j\omega}) = X(e^{-j\omega})$   
 $X[k] = X[(-k) \bmod N] = X[N - k]$  for  $k > 0$

**Conjugate Symmetric:**  $x[n] = x^*[-n]$   
**Conjugate Antisymmetric:**  $x[n] = -x^*[-n]$

## 2: Three Different Fourier Transforms

---

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

# Parseval's Theorem

---

Fourier transforms preserve “energy”

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## Parseval's Theorem

Fourier transforms preserve “energy”

CTFT       $\int |x(t)|^2 dt = \frac{1}{2\pi} \int |X(j\Omega)|^2 d\Omega$

2: Three Different Fourier  
Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## Parseval's Theorem

Fourier transforms preserve “energy”

$$\text{CTFT} \quad \int |x(t)|^2 dt = \frac{1}{2\pi} \int |X(j\Omega)|^2 d\Omega$$

$$\text{DTFT} \quad \sum_{-\infty}^{\infty} |x[n]|^2 = \frac{1}{2\pi} \int_{-\pi}^{\pi} |X(e^{j\omega})|^2 d\omega$$

2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## Parseval's Theorem

Fourier transforms preserve “energy”

$$\text{CTFT} \quad \int |x(t)|^2 dt = \frac{1}{2\pi} \int |X(j\Omega)|^2 d\Omega$$

$$\text{DTFT} \quad \sum_{-\infty}^{\infty} |x[n]|^2 = \frac{1}{2\pi} \int_{-\pi}^{\pi} |X(e^{j\omega})|^2 d\omega$$

$$\text{DFT} \quad \sum_0^{N-1} |x[n]|^2 = \frac{1}{N} \sum_0^{N-1} |X[k]|^2$$

2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## Parseval's Theorem

Fourier transforms preserve “energy”

$$\text{CTFT} \quad \int |x(t)|^2 dt = \frac{1}{2\pi} \int |X(j\Omega)|^2 d\Omega$$

$$\text{DTFT} \quad \sum_{-\infty}^{\infty} |x[n]|^2 = \frac{1}{2\pi} \int_{-\pi}^{\pi} |X(e^{j\omega})|^2 d\omega$$

$$\text{DFT} \quad \sum_0^{N-1} |x[n]|^2 = \frac{1}{N} \sum_0^{N-1} |X[k]|^2$$

More generally, they actually preserve **complex inner products**:

$$\sum_0^{N-1} x[n]y^*[n] = \frac{1}{N} \sum_0^{N-1} X[k]Y^*[k]$$

2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## Parseval's Theorem

Fourier transforms preserve “energy”

$$\text{CTFT} \quad \int |x(t)|^2 dt = \frac{1}{2\pi} \int |X(j\Omega)|^2 d\Omega$$

$$\text{DTFT} \quad \sum_{-\infty}^{\infty} |x[n]|^2 = \frac{1}{2\pi} \int_{-\pi}^{\pi} |X(e^{j\omega})|^2 d\omega$$

$$\text{DFT} \quad \sum_0^{N-1} |x[n]|^2 = \frac{1}{N} \sum_0^{N-1} |X[k]|^2$$

More generally, they actually preserve **complex inner products**:

$$\sum_0^{N-1} x[n]y^*[n] = \frac{1}{N} \sum_0^{N-1} X[k]Y^*[k]$$

Unitary matrix viewpoint for DFT:

If we regard  $\mathbf{x}$  and  $\mathbf{X}$  as vectors, then  $\mathbf{X} = \mathbf{F}\mathbf{x}$  where  $\mathbf{F}$  is a symmetric matrix defined by  $f_{k+1,n+1} = e^{-j2\pi \frac{kn}{N}}$ .

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## Parseval's Theorem

Fourier transforms preserve “energy”

$$\text{CTFT} \quad \int |x(t)|^2 dt = \frac{1}{2\pi} \int |X(j\Omega)|^2 d\Omega$$

$$\text{DTFT} \quad \sum_{-\infty}^{\infty} |x[n]|^2 = \frac{1}{2\pi} \int_{-\pi}^{\pi} |X(e^{j\omega})|^2 d\omega$$

$$\text{DFT} \quad \sum_0^{N-1} |x[n]|^2 = \frac{1}{N} \sum_0^{N-1} |X[k]|^2$$

More generally, they actually preserve **complex inner products**:

$$\sum_0^{N-1} x[n]y^*[n] = \frac{1}{N} \sum_0^{N-1} X[k]Y^*[k]$$

Unitary matrix viewpoint for DFT:

If we regard  $\mathbf{x}$  and  $\mathbf{X}$  as vectors, then  $\mathbf{X} = \mathbf{F}\mathbf{x}$  where  $\mathbf{F}$  is a symmetric matrix defined by  $f_{k+1,n+1} = e^{-j2\pi \frac{kn}{N}}$ .

The inverse DFT matrix is  $\mathbf{F}^{-1} = \frac{1}{N} \mathbf{F}^H$   
equivalently,  $\mathbf{G} = \frac{1}{\sqrt{N}} \mathbf{F}$  is a **unitary matrix** with  $\mathbf{G}^H \mathbf{G} = \mathbf{I}$ .

# Convolution

2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## DTFT: Convolution → Product

# Convolution

2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

DTFT: Convolution → Product

$$x[n] = g[n] * h[n] = \sum_{k=-\infty}^{\infty} g[k]h[n - k]$$

*weighted sum of shifted sequence*

2: Three Different Fourier  
Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## Convolution

DTFT: Convolution → Product

$$x[n] = g[n] * h[n] = \sum_{k=-\infty}^{\infty} g[k]h[n-k]$$
$$\Rightarrow X(e^{j\omega}) = G(e^{j\omega})H(e^{j\omega})$$

2: Three Different Fourier  
Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## Convolution

DTFT: Convolution → Product

$$x[n] = g[n] * h[n] = \sum_{k=-\infty}^{\infty} g[k]h[n-k]$$
$$\Rightarrow X(e^{j\omega}) = G(e^{j\omega})H(e^{j\omega})$$

DFT: Circular convolution → Product

$$x[n] = g[n] \circledast_N h[n] = \sum_{k=0}^{N-1} g[k]h[(n-k) \bmod N]$$

2: Three Different Fourier  
Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## Convolution

### DTFT: Convolution → Product

$$x[n] = g[n] * h[n] = \sum_{k=-\infty}^{\infty} g[k]h[n-k]$$
$$\Rightarrow X(e^{j\omega}) = G(e^{j\omega})H(e^{j\omega})$$

### DFT: Circular convolution → Product

$$x[n] = g[n] \circledast_N h[n] = \sum_{k=0}^{N-1} g[k]h[(n-k) \bmod N]$$
$$\Rightarrow X[k] = G[k]H[k]$$

## 2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines


# Convolution

DTFT: Convolution → Product

$$x[n] = g[n] * h[n] = \sum_{k=-\infty}^{\infty} g[k]h[n-k]$$
$$\Rightarrow X(e^{j\omega}) = G(e^{j\omega})H(e^{j\omega})$$

DFT: Circular convolution → Product

$$x[n] = g[n] \circledast_N h[n] = \sum_{k=0}^{N-1} g[k]h[(n-k) \bmod N]$$
$$\Rightarrow X[k] = G[k]H[k]$$



# Convolution

2: Three Different Fourier Transforms


- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

DTFT: Convolution → Product

$$x[n] = g[n] * h[n] = \sum_{k=-\infty}^{\infty} g[k]h[n-k]$$
$$\Rightarrow X(e^{j\omega}) = G(e^{j\omega})H(e^{j\omega})$$

DFT: Circular convolution → Product

$$x[n] = g[n] \circledast_N h[n] = \sum_{k=0}^{N-1} g[k]h[(n-k) \bmod N]$$
$$\Rightarrow X[k] = G[k]H[k]$$



# Convolution

2: Three Different Fourier Transforms


- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

DTFT: Convolution → Product

$$x[n] = g[n] * h[n] = \sum_{k=-\infty}^{\infty} g[k]h[n-k]$$
$$\Rightarrow X(e^{j\omega}) = G(e^{j\omega})H(e^{j\omega})$$

DFT: Circular convolution → Product

$$x[n] = g[n] \circledast_N h[n] = \sum_{k=0}^{N-1} g[k]h[(n-k) \bmod N]$$
$$\Rightarrow X[k] = G[k]H[k]$$



# Convolution

2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

DTFT: Convolution → Product


$$x[n] = g[n] * h[n] = \sum_{k=-\infty}^{\infty} g[k]h[n-k]$$
$$\Rightarrow X(e^{j\omega}) = G(e^{j\omega})H(e^{j\omega})$$

DFT: Circular convolution → Product

$$x[n] = g[n] \circledast_N h[n] = \sum_{k=0}^{N-1} g[k]h[(n-k) \bmod N]$$
$$\Rightarrow X[k] = G[k]H[k]$$

DTFT: Product → Circular Convolution  $\div 2\pi$

$$y[n] = g[n]h[n]$$



# Convolution

2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

DTFT: Convolution → Product


$$x[n] = g[n] * h[n] = \sum_{k=-\infty}^{\infty} g[k]h[n-k]$$
$$\Rightarrow X(e^{j\omega}) = G(e^{j\omega})H(e^{j\omega})$$

DFT: Circular convolution → Product

$$x[n] = g[n] \circledast_N h[n] = \sum_{k=0}^{N-1} g[k]h[(n-k) \bmod N]$$
$$\Rightarrow X[k] = G[k]H[k]$$

DTFT: Product → Circular Convolution  $\div 2\pi$

$$y[n] = g[n]h[n]$$
$$\Rightarrow Y(e^{j\omega}) = \frac{1}{2\pi} G(e^{j\omega}) \circledast_{\pi} H(e^{j\omega}) =$$
$$\frac{1}{2\pi} \int_{-\pi}^{\pi} G(e^{j\theta})H(e^{j(\omega-\theta)})d\theta$$



# Convolution

2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

DTFT: Convolution → Product

$$x[n] = g[n] * h[n] = \sum_{k=-\infty}^{\infty} g[k]h[n-k]$$
$$\Rightarrow X(e^{j\omega}) = G(e^{j\omega})H(e^{j\omega})$$

DFT: Circular convolution → Product


$$x[n] = g[n] \circledast_N h[n] = \sum_{k=0}^{N-1} g[k]h[(n-k) \bmod N]$$
$$\Rightarrow X[k] = G[k]H[k]$$

DTFT: Product → Circular Convolution  $\div 2\pi$

$$y[n] = g[n]h[n]$$
$$\Rightarrow Y(e^{j\omega}) = \frac{1}{2\pi} G(e^{j\omega}) \circledast_{\pi} H(e^{j\omega}) =$$
$$\frac{1}{2\pi} \int_{-\pi}^{\pi} G(e^{j\theta})H(e^{j(\omega-\theta)})d\theta$$

DFT: Product → Circular Convolution  $\div N$

$$y[n] = g[n]h[n]$$



# Convolution

2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

DTFT: Convolution → Product

$$x[n] = g[n] * h[n] = \sum_{k=-\infty}^{\infty} g[k]h[n-k]$$
$$\Rightarrow X(e^{j\omega}) = G(e^{j\omega})H(e^{j\omega})$$

DFT: Circular convolution → Product


$$x[n] = g[n] \circledast_N h[n] = \sum_{k=0}^{N-1} g[k]h[(n-k) \bmod N]$$
$$\Rightarrow X[k] = G[k]H[k]$$

DTFT: Product → Circular Convolution  $\div 2\pi$

$$y[n] = g[n]h[n]$$
$$\Rightarrow Y(e^{j\omega}) = \frac{1}{2\pi} G(e^{j\omega}) \circledast_{\pi} H(e^{j\omega}) =$$
$$\frac{1}{2\pi} \int_{-\pi}^{\pi} G(e^{j\theta})H(e^{j(\omega-\theta)})d\theta$$

DFT: Product → Circular Convolution  $\div N$

$$y[n] = g[n]h[n]$$
$$\Rightarrow Y[k] = \frac{1}{N} G[k] \circledast_N H[k]$$




# Sampling Process




# Sampling Process




# Sampling Process



# Sampling Process



# Sampling Process



## 2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- **Zero-Padding**
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

# Zero-Padding

Zero padding means added extra zeros onto the end of  $x[n]$  before performing the DFT.

Windowed Signal

Time  $x[n]$



2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- **Zero-Padding**
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines


# Zero-Padding

Zero padding means added extra zeros onto the end of  $x[n]$  before performing the DFT.

Windowed Signal

With zero-padding

Time  $x[n]$



## 2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- **Zero-Padding**
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines


# Zero-Padding

Zero padding means added extra zeros onto the end of  $x[n]$  before performing the DFT.


Windowed Signal

With zero-padding

Time  $x[n]$



Frequency  $|X[k]|$



## 2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- **Zero-Padding**
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines


# Zero-Padding

Zero padding means added extra zeros onto the end of  $x[n]$  before performing the DFT.


Windowed Signal

With zero-padding

Time  $x[n]$



Frequency  $|X[k]|$



# Zero-Padding

2: Three Different Fourier Transforms


- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- **Zero-Padding**
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

Zero padding means added extra zeros onto the end of  $x[n]$  before performing the DFT.


Windowed Signal

With zero-padding

Time  $x[n]$



Frequency  $|X[k]|$



- Zero-padding causes the DFT to evaluate the DTFT at more values of  $\omega_k$ . Denser frequency samples.

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- **Zero-Padding**
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines


## Zero-Padding

Zero padding means added extra zeros onto the end of  $x[n]$  before performing the DFT.


Windowed Signal

With zero-padding

Time  $x[n]$



Frequency  $|X[k]|$



- Zero-padding causes the DFT to evaluate the DTFT at more values of  $\omega_k$ . Denser frequency samples.
- Width of the peaks remains constant: determined by the length and shape of the window.

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- **Zero-Padding**
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines


## Zero-Padding

Zero padding means added extra zeros onto the end of  $x[n]$  before performing the DFT.


Windowed Signal

With zero-padding

Time  $x[n]$



Frequency  $|X[k]|$




- Zero-padding causes the DFT to evaluate the DTFT at more values of  $\omega_k$ . Denser frequency samples.
- Width of the peaks remains constant: determined by the length and shape of the window.
- Smoother graph but increased frequency resolution is an illusion.

# Phase Unwrapping


2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

Phase of a DTFT is only defined to within an integer multiple of  $2\pi$ .



$x[n]$



$|X[k]|$

# Phase Unwrapping


2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines


Phase of a DTFT is only defined to within an integer multiple of  $2\pi$ .



$x[n]$



$|X[k]|$



$\angle X[k]$

# Phase Unwrapping


2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines


Phase of a DTFT is only defined to within an integer multiple of  $2\pi$ .



$x[n]$



$|X[k]|$



$\angle X[k]$

Phase unwrapping adds multiples of  $2\pi$  onto each  $\angle X[k]$  to make the phase as continuous as possible.

# Phase Unwrapping


2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines


Phase of a DTFT is only defined to within an integer multiple of  $2\pi$ .




$x[n]$



$|X[k]|$



$\angle X[k]$



$\angle X[k]$  unwrapped

Phase unwrapping adds multiples of  $2\pi$  onto each  $\angle X[k]$  to make the phase as continuous as possible.

2: Three Different Fourier  
Transforms

---

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- **Uncertainty principle**
- Summary
- MATLAB routines

## Uncertainty principle

**CTFT uncertainty principle:**  $\left( \frac{\int t^2 |x(t)|^2 dt}{\int |x(t)|^2 dt} \right)^{\frac{1}{2}} \left( \frac{\int \omega^2 |X(j\omega)|^2 d\omega}{\int |X(j\omega)|^2 d\omega} \right)^{\frac{1}{2}} \geq \frac{1}{2}$

## Fourier Uncertainty Relation

A non-zero signal can not be simultaneously localized in time and frequency domains.

---

<sup>1</sup> "Uncertainty measures and uncertainty relations for angle observables," *Foundations of Physics*, vol. 15, pp. 353–364, 1983.

## Fourier Uncertainty Relation

A non-zero signal can not be simultaneously localized in time and frequency domains.

For an intuitive understanding, consider the case of a Gaussian function. This is an example of a self-Fourier transform. That is,

$$g_\sigma(t) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{t^2}{2\sigma^2}} \xrightarrow{\text{Fourier}} \hat{g}_\sigma(\omega) = e^{-\frac{\sigma^2\omega^2}{2}}.$$

And hence,

$$g_\sigma(t)\hat{g}_\sigma(\omega) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\left(\frac{t^2}{2\sigma^2} + \frac{\sigma^2\omega^2}{2}\right)}, \quad \underbrace{\frac{t^2}{2\sigma^2} + \frac{\sigma^2\omega^2}{2}}_{\text{Ellipse}}.$$

Shrinking  $\sigma$  in one domain leads to fattening in another domain.

---

<sup>1</sup> "Uncertainty measures and uncertainty relations for angle observables," *Foundations of Physics*, vol. 15, pp. 353–364, 1983.

## Fourier Uncertainty Relation

A non-zero signal can not be simultaneously localized in time and frequency domains.

For an intuitive understanding, consider the case of a Gaussian function. This is an example of a self-Fourier transform. That is,

$$g_\sigma(t) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{t^2}{2\sigma^2}} \xrightarrow{\text{Fourier}} \hat{g}_\sigma(\omega) = e^{-\frac{\sigma^2\omega^2}{2}}.$$

And hence,

$$g_\sigma(t)\hat{g}_\sigma(\omega) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\left(\frac{t^2}{2\sigma^2} + \frac{\sigma^2\omega^2}{2}\right)}, \quad \underbrace{\frac{t^2}{2\sigma^2} + \frac{\sigma^2\omega^2}{2}}_{\text{Ellipse}}.$$

Shrinking  $\sigma$  in one domain leads to fattening in another domain.

- Interesting topic with historical roots and interpretations in Quantum mechanics.
- There is a variant for discrete sequences that can measure “localization” and goes back to the work of Breitenberger<sup>1</sup>.

---

<sup>1</sup> “Uncertainty measures and uncertainty relations for angle observables,” *Foundations of Physics*, vol. 15, pp. 353–364, 1983.

## Fourier Uncertainty Relation


The “localization” or spread is measured by,

$$\text{spread}^2 = \int_{\mathbb{R}} (t - \text{center})f(t)dt \quad \text{where} \quad \text{center} = \int_{\mathbb{R}} tf(t)dt.$$

## Fourier Uncertainty Relation

The “localization” or spread is measured by,


$$\text{spread}^2 = \int_{\mathbb{R}} (t - \text{center})f(t)dt \quad \text{where} \quad \text{center} = \int_{\mathbb{R}} tf(t)dt.$$



## Fourier Uncertainty Relation

The “localization” or spread is measured by,

$$\text{spread}^2 = \int_{\mathbb{R}} (t - \text{center})f(t)dt \quad \text{where} \quad \text{center} = \int_{\mathbb{R}} tf(t)dt.$$




Not surprisingly, for  $f(t) = e^{-\frac{(t-\mu)^2}{2\sigma^2}}$  (a Gaussian function), its center and spread are its mean and variance, respectively,

$$\mu \triangleq \int_{\mathbb{R}} tf(t)dt \quad \text{and} \quad \sigma^2 \triangleq \int_{\mathbb{R}} (t - \mu)^2 f(t) dt.$$

## Fourier Uncertainty Relation

The “localization” or spread is measured by,

$$\text{spread}^2 = \int_{\mathbb{R}} (t - \text{center})f(t)dt \quad \text{where} \quad \text{center} = \int_{\mathbb{R}} tf(t)dt.$$



Not surprisingly, for  $f(t) = e^{-\frac{(t-\mu)^2}{2\sigma^2}}$  (a Gaussian function), its center and spread are its mean and variance, respectively,

$$\mu \triangleq \int_{\mathbb{R}} tf(t)dt \quad \text{and} \quad \sigma^2 \triangleq \int_{\mathbb{R}} (t - \mu)^2 f(t)dt.$$

When dealing with Fourier Uncertainty Principles, we work with **squared functions**, that is,  $|f(t)|^2$ .

2: Three Different Fourier  
Transforms

---

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- **Uncertainty principle**
- Summary
- MATLAB routines

## Uncertainty principle

$$\text{CTFT uncertainty principle: } \left( \frac{\int t^2 |x(t)|^2 dt}{\int |x(t)|^2 dt} \right)^{\frac{1}{2}} \left( \frac{\int \omega^2 |X(j\omega)|^2 d\omega}{\int |X(j\omega)|^2 d\omega} \right)^{\frac{1}{2}} \geq \frac{1}{2}$$

The first term measures the “width” of  $x(t)$  around  $t = 0$ .

It is like  $\sigma$  if  $|x(t)|^2$  was a zero-mean probability distribution.

2: Three Different Fourier  
Transforms

---

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## Uncertainty principle

$$\text{CTFT uncertainty principle: } \left( \frac{\int t^2 |x(t)|^2 dt}{\int |x(t)|^2 dt} \right)^{\frac{1}{2}} \left( \frac{\int \omega^2 |X(j\omega)|^2 d\omega}{\int |X(j\omega)|^2 d\omega} \right)^{\frac{1}{2}} \geq \frac{1}{2}$$

The first term measures the “width” of  $x(t)$  around  $t = 0$ .

It is like  $\sigma$  if  $|x(t)|^2$  was a zero-mean probability distribution.

The second term is similarly the “width” of  $X(j\omega)$  in frequency.

A signal **cannot be concentrated in both time and frequency.**

2: Three Different Fourier  
Transforms

---

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## Uncertainty principle

$$\text{CTFT uncertainty principle: } \left( \frac{\int t^2 |x(t)|^2 dt}{\int |x(t)|^2 dt} \right)^{\frac{1}{2}} \left( \frac{\int \omega^2 |X(j\omega)|^2 d\omega}{\int |X(j\omega)|^2 d\omega} \right)^{\frac{1}{2}} \geq \frac{1}{2}$$

The first term measures the “width” of  $x(t)$  around  $t = 0$ .

It is like  $\sigma$  if  $|x(t)|^2$  was a zero-mean probability distribution.

The second term is similarly the “width” of  $X(j\omega)$  in frequency.

A signal **cannot be concentrated in both time and frequency.**

**Proof Outline:**

Assume  $\int |x(t)|^2 dt = 1 \Rightarrow \int |X(j\omega)|^2 d\omega = 2\pi$  [Parseval]

2: Three Different Fourier  
Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## Uncertainty principle

$$\text{CTFT uncertainty principle: } \left( \frac{\int t^2 |x(t)|^2 dt}{\int |x(t)|^2 dt} \right)^{\frac{1}{2}} \left( \frac{\int \omega^2 |X(j\omega)|^2 d\omega}{\int |X(j\omega)|^2 d\omega} \right)^{\frac{1}{2}} \geq \frac{1}{2}$$

The first term measures the “width” of  $x(t)$  around  $t = 0$ .

It is like  $\sigma$  if  $|x(t)|^2$  was a zero-mean probability distribution.

The second term is similarly the “width” of  $X(j\omega)$  in frequency.

A signal **cannot be concentrated in both time and frequency.**

### Proof Outline:

Assume  $\int |x(t)|^2 dt = 1 \Rightarrow \int |X(j\omega)|^2 d\omega = 2\pi$  [Parseval]

Set  $v(t) = \frac{dx}{dt} \Rightarrow V(j\omega) = j\omega X(j\omega)$  [by parts]

2: Three Different Fourier  
Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## Uncertainty principle

$$\text{CTFT uncertainty principle: } \left( \frac{\int t^2 |x(t)|^2 dt}{\int |x(t)|^2 dt} \right)^{\frac{1}{2}} \left( \frac{\int \omega^2 |X(j\omega)|^2 d\omega}{\int |X(j\omega)|^2 d\omega} \right)^{\frac{1}{2}} \geq \frac{1}{2}$$

The first term measures the “width” of  $x(t)$  around  $t = 0$ .

It is like  $\sigma$  if  $|x(t)|^2$  was a zero-mean probability distribution.

The second term is similarly the “width” of  $X(j\omega)$  in frequency.

A signal **cannot be concentrated in both time and frequency.**

### Proof Outline:

Assume  $\int |x(t)|^2 dt = 1 \Rightarrow \int |X(j\omega)|^2 d\omega = 2\pi$  [Parseval]

Set  $v(t) = \frac{dx}{dt} \Rightarrow V(j\omega) = j\omega X(j\omega)$  [by parts]

Now  $\int tx \frac{dx}{dt} dt = \frac{1}{2}tx^2(t) \Big|_{t=-\infty}^{\infty} - \int \frac{1}{2}x^2 dt = 0 - \frac{1}{2}$  [by parts]

2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## Uncertainty principle

$$\text{CTFT uncertainty principle: } \left( \frac{\int t^2 |x(t)|^2 dt}{\int |x(t)|^2 dt} \right)^{\frac{1}{2}} \left( \frac{\int \omega^2 |X(j\omega)|^2 d\omega}{\int |X(j\omega)|^2 d\omega} \right)^{\frac{1}{2}} \geq \frac{1}{2}$$

The first term measures the “width” of  $x(t)$  around  $t = 0$ .

It is like  $\sigma$  if  $|x(t)|^2$  was a zero-mean probability distribution.

The second term is similarly the “width” of  $X(j\omega)$  in frequency.

A signal **cannot be concentrated in both time and frequency.**

### Proof Outline:

Assume  $\int |x(t)|^2 dt = 1 \Rightarrow \int |X(j\omega)|^2 d\omega = 2\pi$  [Parseval]

Set  $v(t) = \frac{dx}{dt} \Rightarrow V(j\omega) = j\omega X(j\omega)$  [by parts]

Now  $\int tx \frac{dx}{dt} dt = \frac{1}{2}tx^2(t) \Big|_{t=-\infty}^{\infty} - \int \frac{1}{2}x^2 dt = 0 - \frac{1}{2}$  [by parts]

So  $\frac{1}{4} = \left| \int tx \frac{dx}{dt} dt \right|^2 \leq \left( \int t^2 x^2 dt \right) \left( \int \left| \frac{dx}{dt} \right|^2 dt \right)$  [Schwartz]

2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## Uncertainty principle

$$\text{CTFT uncertainty principle: } \left( \frac{\int t^2 |x(t)|^2 dt}{\int |x(t)|^2 dt} \right)^{\frac{1}{2}} \left( \frac{\int \omega^2 |X(j\omega)|^2 d\omega}{\int |X(j\omega)|^2 d\omega} \right)^{\frac{1}{2}} \geq \frac{1}{2}$$

The first term measures the “width” of  $x(t)$  around  $t = 0$ .

It is like  $\sigma$  if  $|x(t)|^2$  was a zero-mean probability distribution.

The second term is similarly the “width” of  $X(j\omega)$  in frequency.

A signal **cannot be concentrated in both time and frequency.**

### Proof Outline:

Assume  $\int |x(t)|^2 dt = 1 \Rightarrow \int |X(j\omega)|^2 d\omega = 2\pi$  [Parseval]

Set  $v(t) = \frac{dx}{dt} \Rightarrow V(j\omega) = j\omega X(j\omega)$  [by parts]

Now  $\int tx \frac{dx}{dt} dt = \frac{1}{2}tx^2(t) \Big|_{t=-\infty}^{\infty} - \int \frac{1}{2}x^2 dt = 0 - \frac{1}{2}$  [by parts]

So  $\frac{1}{4} = \left| \int tx \frac{dx}{dt} dt \right|^2 \leq \left( \int t^2 x^2 dt \right) \left( \int \left| \frac{dx}{dt} \right|^2 dt \right)$  [Schwartz]  
 $= \left( \int t^2 x^2 dt \right) \left( \int |v(t)|^2 dt \right)$

2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## Uncertainty principle

$$\text{CTFT uncertainty principle: } \left( \frac{\int t^2 |x(t)|^2 dt}{\int |x(t)|^2 dt} \right)^{\frac{1}{2}} \left( \frac{\int \omega^2 |X(j\omega)|^2 d\omega}{\int |X(j\omega)|^2 d\omega} \right)^{\frac{1}{2}} \geq \frac{1}{2}$$

The first term measures the “width” of  $x(t)$  around  $t = 0$ .

It is like  $\sigma$  if  $|x(t)|^2$  was a zero-mean probability distribution.

The second term is similarly the “width” of  $X(j\omega)$  in frequency.

A signal **cannot be concentrated in both time and frequency.**

### Proof Outline:

Assume  $\int |x(t)|^2 dt = 1 \Rightarrow \int |X(j\omega)|^2 d\omega = 2\pi$  [Parseval]

Set  $v(t) = \frac{dx}{dt} \Rightarrow V(j\omega) = j\omega X(j\omega)$  [by parts]

Now  $\int tx \frac{dx}{dt} dt = \frac{1}{2}tx^2(t) \Big|_{t=-\infty}^{\infty} - \int \frac{1}{2}x^2 dt = 0 - \frac{1}{2}$  [by parts]

So  $\frac{1}{4} = \left| \int tx \frac{dx}{dt} dt \right|^2 \leq \left( \int t^2 x^2 dt \right) \left( \int \left| \frac{dx}{dt} \right|^2 dt \right)$  [Schwartz]

$$= \left( \int t^2 x^2 dt \right) \left( \int |v(t)|^2 dt \right) = \left( \int t^2 x^2 dt \right) \left( \frac{1}{2\pi} \int |V(j\omega)|^2 d\omega \right)$$

2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## Uncertainty principle

$$\text{CTFT uncertainty principle: } \left( \frac{\int t^2 |x(t)|^2 dt}{\int |x(t)|^2 dt} \right)^{\frac{1}{2}} \left( \frac{\int \omega^2 |X(j\omega)|^2 d\omega}{\int |X(j\omega)|^2 d\omega} \right)^{\frac{1}{2}} \geq \frac{1}{2}$$

The first term measures the “width” of  $x(t)$  around  $t = 0$ .

It is like  $\sigma$  if  $|x(t)|^2$  was a zero-mean probability distribution.

The second term is similarly the “width” of  $X(j\omega)$  in frequency.

A signal **cannot be concentrated in both time and frequency.**

### Proof Outline:

Assume  $\int |x(t)|^2 dt = 1 \Rightarrow \int |X(j\omega)|^2 d\omega = 2\pi$  [Parseval]

Set  $v(t) = \frac{dx}{dt} \Rightarrow V(j\omega) = j\omega X(j\omega)$  [by parts]

Now  $\int tx \frac{dx}{dt} dt = \frac{1}{2}tx^2(t) \Big|_{t=-\infty}^{\infty} - \int \frac{1}{2}x^2 dt = 0 - \frac{1}{2}$  [by parts]

So  $\frac{1}{4} = \left| \int tx \frac{dx}{dt} dt \right|^2 \leq \left( \int t^2 x^2 dt \right) \left( \int \left| \frac{dx}{dt} \right|^2 dt \right)$  [Schwartz]

$$= \left( \int t^2 x^2 dt \right) \left( \int |v(t)|^2 dt \right) = \left( \int t^2 x^2 dt \right) \left( \frac{1}{2\pi} \int |V(j\omega)|^2 d\omega \right)$$
$$= \left( \int t^2 x^2 dt \right) \left( \frac{1}{2\pi} \int \omega^2 |X(j\omega)|^2 d\omega \right)$$

2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## Uncertainty principle

$$\text{CTFT uncertainty principle: } \left( \frac{\int t^2 |x(t)|^2 dt}{\int |x(t)|^2 dt} \right)^{\frac{1}{2}} \left( \frac{\int \omega^2 |X(j\omega)|^2 d\omega}{\int |X(j\omega)|^2 d\omega} \right)^{\frac{1}{2}} \geq \frac{1}{2}$$

The first term measures the “width” of  $x(t)$  around  $t = 0$ .

It is like  $\sigma$  if  $|x(t)|^2$  was a zero-mean probability distribution.

The second term is similarly the “width” of  $X(j\omega)$  in frequency.

A signal **cannot be concentrated in both time and frequency.**

### Proof Outline:

Assume  $\int |x(t)|^2 dt = 1 \Rightarrow \int |X(j\omega)|^2 d\omega = 2\pi$  [Parseval]

Set  $v(t) = \frac{dx}{dt} \Rightarrow V(j\omega) = j\omega X(j\omega)$  [by parts]

Now  $\int tx \frac{dx}{dt} dt = \frac{1}{2}tx^2(t) \Big|_{t=-\infty}^{\infty} - \int \frac{1}{2}x^2 dt = 0 - \frac{1}{2}$  [by parts]

So  $\frac{1}{4} = \left| \int tx \frac{dx}{dt} dt \right|^2 \leq \left( \int t^2 x^2 dt \right) \left( \int \left| \frac{dx}{dt} \right|^2 dt \right)$  [Schwartz]

$$= \left( \int t^2 x^2 dt \right) \left( \int |v(t)|^2 dt \right) = \left( \int t^2 x^2 dt \right) \left( \frac{1}{2\pi} \int |V(j\omega)|^2 d\omega \right)$$
$$= \left( \int t^2 x^2 dt \right) \left( \frac{1}{2\pi} \int \omega^2 |X(j\omega)|^2 d\omega \right)$$

No exact equivalent for DTFT/DFT but a similar effect is true

2: Three Different Fourier  
Transforms

---

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- **Summary**
- MATLAB routines

## Summary

- Three types: CTFT, DTFT, DFT

2: Three Different Fourier  
Transforms

---

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- **Summary**
- MATLAB routines

## Summary

- Three types: CTFT, DTFT, DFT
  - $\text{DTFT} = \text{CTFT of continuous signal} \times \text{impulse train}$

2: Three Different Fourier  
Transforms

---

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- **Summary**
- MATLAB routines

## Summary

- Three types: CTFT, DTFT, DFT
  - DTFT = CTFT of continuous signal  $\times$  impulse train
  - DFT = DTFT of periodic or finite support signal

2: Three Different Fourier Transforms

---

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- **Summary**
- MATLAB routines

## Summary

- Three types: CTFT, DTFT, DFT
  - DTFT = CTFT of continuous signal  $\times$  impulse train
  - DFT = DTFT of periodic or finite support signal
  - DFT is a scaled unitary transform

2: Three Different Fourier Transforms

---

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- **Summary**
- MATLAB routines

## Summary

- Three types: CTFT, DTFT, DFT
  - DTFT = CTFT of continuous signal  $\times$  impulse train
  - DFT = DTFT of periodic or finite support signal
    - DFT is a scaled unitary transform
- DTFT: Convolution  $\rightarrow$  Product; Product  $\rightarrow$  Circular Convolution

## 2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- **Summary**
- MATLAB routines

# Summary

- Three types: CTFT, DTFT, DFT
  - DTFT = CTFT of continuous signal  $\times$  impulse train
  - DFT = DTFT of periodic or finite support signal
    - DFT is a scaled unitary transform
- DTFT: Convolution  $\rightarrow$  Product; Product  $\rightarrow$  Circular Convolution
- DFT: Product  $\leftrightarrow$  Circular Convolution

## 2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- **Summary**
- MATLAB routines

# Summary

- Three types: CTFT, DTFT, DFT
  - DTFT = CTFT of continuous signal  $\times$  impulse train
  - DFT = DTFT of periodic or finite support signal
    - DFT is a scaled unitary transform
- DTFT: Convolution  $\rightarrow$  Product; Product  $\rightarrow$  Circular Convolution
- DFT: Product  $\leftrightarrow$  Circular Convolution
- DFT: Zero Padding  $\rightarrow$  Denser freq sampling but same resolution

## 2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- **Summary**
- MATLAB routines

# Summary

- Three types: CTFT, DTFT, DFT
  - DTFT = CTFT of continuous signal  $\times$  impulse train
  - DFT = DTFT of periodic or finite support signal
    - DFT is a scaled unitary transform
- DTFT: Convolution  $\rightarrow$  Product; Product  $\rightarrow$  Circular Convolution
- DFT: Product  $\leftrightarrow$  Circular Convolution
- DFT: Zero Padding  $\rightarrow$  Denser freq sampling but same resolution
- Phase is only defined to within a multiple of  $2\pi$ .

## 2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- **Summary**
- MATLAB routines

# Summary

- Three types: CTFT, DTFT, DFT
  - DTFT = CTFT of continuous signal  $\times$  impulse train
  - DFT = DTFT of periodic or finite support signal
    - DFT is a scaled unitary transform
- DTFT: Convolution  $\rightarrow$  Product; Product  $\rightarrow$  Circular Convolution
- DFT: Product  $\leftrightarrow$  Circular Convolution
- DFT: Zero Padding  $\rightarrow$  Denser freq sampling but same resolution
- Phase is only defined to within a multiple of  $2\pi$ .
- Whenever you integrate over frequency you need a **scale factor**
  - $\frac{1}{2\pi}$  for CTFT and DTFT or  $\frac{1}{N}$  for DFT
  - e.g. Inverse transform, Parseval, frequency domain convolution

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- **Summary**
- MATLAB routines

## Summary

- Three types: CTFT, DTFT, DFT
  - DTFT = CTFT of continuous signal  $\times$  impulse train
  - DFT = DTFT of periodic or finite support signal
    - DFT is a scaled unitary transform
- DTFT: Convolution  $\rightarrow$  Product; Product  $\rightarrow$  Circular Convolution
- DFT: Product  $\leftrightarrow$  Circular Convolution
- DFT: Zero Padding  $\rightarrow$  Denser freq sampling but same resolution
- Phase is only defined to within a multiple of  $2\pi$ .
- Whenever you integrate over frequency you need a **scale factor**
  - $\frac{1}{2\pi}$  for CTFT and DTFT or  $\frac{1}{N}$  for DFT
  - e.g. Inverse transform, Parseval, frequency domain convolution

For further details see Mitra: 3 & 5.

## 2: Three Different Fourier Transforms

- Fourier Transforms
- Convergence of DTFT
- DTFT Properties
- DFT Properties
- Symmetries
- Parseval's Theorem
- Convolution
- Sampling Process
- Zero-Padding
- Phase Unwrapping
- Uncertainty principle
- Summary
- MATLAB routines

## MATLAB routines

fft, ifft	DFT with optional zero-padding
fftshift	swap the two halves of a vector
conv	convolution or polynomial multiplication (not circular)
$x[n] \circledast y[n]$	<code>real(ifft(fft(x).*fft(y)))</code>
unwrap	remove $2\pi$ jumps from phase spectrum