

Solutions 2015

[A] application

[B] book word

[T] new theory

[E] new example

1. a) Just expand RHS

$$H(\mathbf{q}) + f_2 H\left(\frac{p_2}{f_2}, \frac{p_3}{f_2}\right) \\ = -f_1 \log f_1 - f_2 \log f_2 - \cancel{f_2 \log \frac{p_2}{f_2}} - \cancel{f_2 \log \frac{p_3}{f_2}} \quad [2E]$$

$$= -p_1 \log p_1 - p_2 \log p_2 - p_3 \log p_3 \\ + \underbrace{p_2 \log f_2 + p_3 \log f_2 - f_2 \log f_2}_{=0 \text{ since } f_2 = p_2 + p_3} \quad [2E]$$

$$= -p_1 \log p_1 - p_2 \log p_2 - p_3 \log p_3 \quad [2E] \\ = H(\mathbf{p})$$

b) We have

| $x_1 x_2$ | 00 | 01 | 10 | 11 |
|-----------|----|----|----|----|
| y | 0 | 0 | 0 | 1 |

each with prob. $\frac{1}{4}$ [1E]

i) Thus

$$p(y=0) = \frac{3}{4} \quad p(y=1) = \frac{1}{4} \\ H(y) = -\frac{3}{4} \log \frac{3}{4} - \frac{1}{4} \log \frac{1}{4} \quad [2E] \\ = \frac{1}{2} + 0.31 \\ = 0.81$$

ii) We have the conditional distribution

| $x_1 \backslash y$ | 0 | 1 |
|--------------------|---------------|---------------|
| 0 | 1 | 0 |
| 1 | $\frac{1}{2}$ | $\frac{1}{2}$ |

[3E]

Thus

$$H(y|x_1) = \frac{1}{2} H(1) + \frac{1}{2} H\left(\frac{1}{2}\right) = \frac{1}{2} \\ I(x_1; y) = H(y) - H(y|x_1) \\ = 0.81 - \frac{1}{2} = 0.31$$

$$\text{iii) } I(x_{1:2}; y) = H(y) - H(y|x_{1:2})$$

$$= H(y) - 0 \quad y \text{ is a function of } x_{1:2} \\ = H(y) = 0.81 \quad [3E]$$

c) X takes values in $\{1, 2, 3, \dots\}$

$X = n$ means that Tail occurs for the first $n-1$ flips,
while Head occurs for the n -th flip. [3 A]

Thus

$$P(X = n) = \left(\frac{1}{2}\right)^{n-1} \frac{1}{2} = \left(\frac{1}{2}\right)^n \quad [3 A]$$

$$H(X) = \sum_{n=1}^{\infty} 2^{-n} \log 2^n$$

$$= \sum_{n=1}^{\infty} n \cdot 2^{-n} \cdot \log 2$$

[4 A]

$$= \frac{\frac{1}{2}}{\left(1 - \frac{1}{2}\right)^2}$$

use the second formula for $r = \frac{1}{2}$

$$= 2 \text{ bits}$$

2. a)

i) The marginal distributions are given by

$$P_X(X) = P_Y(Y) = \left\{ \frac{1}{4}, \frac{3}{4} \right\} \quad [1E]$$

So their entropy is

$$\begin{aligned} H(X) = H(Y) &= H\left(\frac{1}{4}, \frac{3}{4}\right) \quad [2E] \\ &= -\frac{1}{4} \log \frac{1}{4} - \frac{3}{4} \log \frac{3}{4} \\ &= 0.81 \end{aligned}$$

Since ~~the~~ ^{for} sequences ~~X~~ and ~~Y~~

$$\begin{aligned} -\frac{1}{8} \log p(X) &= -\frac{1}{8} \log p(Y) \\ &= -\frac{1}{8} \log \left(\frac{1}{4}\right)^2 \left(\frac{3}{4}\right)^6 \quad [2E] \\ &= -\frac{1}{4} \log \frac{1}{4} - \frac{3}{4} \log \frac{3}{4} \\ &= H(X) = H(Y), \end{aligned}$$

both of them are typical. [1E]

The joint distribution is $P_{XY}(X, Y) = \left\{ \frac{1}{8}, \frac{1}{8}, \frac{1}{8}, \frac{5}{8} \right\}$

$$H(X, Y) = -\frac{3}{8} \log \frac{1}{8} - \frac{5}{8} \log \frac{5}{8} = 1.55 \quad [1E]$$

For sequence (X, Y) , we check

$$\begin{aligned} -\frac{1}{8} \log p(X, Y) &= -\frac{1}{8} \log \left(\frac{1}{8}\right)^4 \left(\frac{5}{8}\right)^4 \quad [2E] \\ &= -\frac{1}{2} \log \frac{1}{8} - \frac{1}{2} \log \frac{5}{8} \\ &= 1.84 > H(X, Y) + \epsilon = 1.55 + 0.2 \end{aligned}$$

So they are not jointly typical. [1E]

ii) c1) total probability theorem [1B]

(2) first term: taking maximum yields an upper bound

second term: $P(A \cap B) \leq P(B)$ [1B]

(3) $|S^{(n)}| < 2^{n(H(X) - 2\epsilon)}$ is given

$$p(X) < 2^{-n(H - \epsilon)} \text{ if typical} \quad [2B]$$

probability of atypical set $< \epsilon$

(4) algebra

$$(5) 2^{-n\epsilon} < \epsilon \quad \text{because } n > -\epsilon^{-1} \log \epsilon \quad [1B]$$

b)

i) If only one source is encoded,


$$3D = D_1 + D_2 + D_3$$

$$> \sigma_1^2 + \sigma_2^2 + \sigma_3^2$$

$$= 2\sigma_2^2 + \sigma_3^2 \Rightarrow D > \frac{2\sigma_2^2 + \sigma_3^2}{3}$$

$$R_1 = \frac{1}{2} \log \frac{\sigma_1^2}{D_1}$$

$$= \frac{1}{2} \log \frac{\sigma_1^2}{3D - \sigma_2^2 - \sigma_3^2}$$



[3A]

$$D_1 = 3D - D_2 - D_3$$

$$= 3D - \sigma_2^2 - \sigma_3^2$$

$$R_2 = R_3 = 0$$

ii) If two sources are encoded


$$3D = D_1 + D_2 + D_3$$

$$> 3\sigma_3^2$$

$$3D < 2\sigma_2^2 + \sigma_3^2$$

$$\sigma_3^2 < D < \frac{2\sigma_2^2 + \sigma_3^2}{3}$$

$$D_1 = D_2 = \frac{3D - \sigma_3^2}{2}$$



[3A]

$$R_1 = \frac{1}{2} \log \frac{\sigma_1^2}{D_1} = \frac{1}{2} \log \left(\frac{2\sigma_1^2}{3D - \sigma_3^2} \right)$$

$$R_2 = \frac{1}{2} \log \frac{\sigma_2^2}{D_2} = \frac{1}{2} \log \left(\frac{2\sigma_2^2}{3D - \sigma_3^2} \right)$$

$$R_3 = 0$$

iii) If all three sources are encoded,


$$D < \sigma_3^2$$

$$D_1 = D_2 = D_3 = D/3$$

$$R_1 = \frac{1}{2} \log \frac{\sigma_1^2}{D_1} = \frac{1}{2} \log \frac{3\sigma_1^2}{D}$$

$$R_2 = \frac{1}{2} \log \frac{3\sigma_2^2}{D}$$

$$R_3 = \frac{1}{2} \log \frac{3\sigma_3^2}{D}$$



[3A]

3. a) (1) average over codewords, average over codes

[18]
Each

(2) Exchange order of summation

(3) for random coding, $\sum_C p(C) \lambda_w(C)$ doesn't depend on index w , so w can be 1 w.o.g

(4) average error prob. of $w=1$

(5) definition of error prob.

(6) Union bound

(7) prob. of atypical set $\leq \epsilon$

prob. of $x(w)$ and y jointly typical $\leq 2^{-n(I(x,y) - 3\epsilon)}$

(8) algebra: $2^{nR} - 1 < 2^{nR}$

(9) $I(X;Y) \leq C$

(10) $n > -\frac{\log \epsilon}{C - R - 3\epsilon} \Rightarrow 2^{-n(C - R - 3\epsilon)} < \epsilon$

(11) by contradiction

(12) by contradiction

(13) Since half of the codewords are gone,

$$\text{rate} = \frac{1}{n} \log(2^{nR}/2) = \frac{nR-1}{n} = R - n^{-1}$$

b)

$$\begin{aligned} \text{i)} \quad Y &= \alpha Y_1 + (1-\alpha) Y_2 \\ &= \alpha(X + Z_1) + (1-\alpha)(X + Z_2) \\ &= X + \alpha Z_1 + (1-\alpha) Z_2 \end{aligned}$$

[BA]

So this is an equivalent Gaussian channel with noise variance $\alpha^2 N_1 + (1-\alpha)^2 N_2$.

Capacity

$$C = \frac{1}{2} \log \left(1 + \frac{P}{\alpha^2 N_1 + (1-\alpha)^2 N_2} \right)$$

[BA]

ii) To maximize C , we minimize noise variance

Let $\frac{\partial}{\partial \alpha} (\alpha^2 N_1 + (1-\alpha)^2 N_2) = 0$, we get

$$\begin{aligned} 2\alpha N_1 - 2(1-\alpha)N_2 &= 0 \\ \alpha &= \frac{N_2}{N_1 + N_2} \end{aligned}$$

[BA]

Capacity

$$C = \frac{1}{2} \log \left(1 + \frac{P}{N_1} + \frac{P}{N_2} \right)$$

[BA]

4.

a) Slepian-Wolf region

$$R_1 \geq H(X|Y)$$

$$R_2 \geq H(Y|X)$$

[3B]

$$R_1 + R_2 \geq H(X, Y)$$

We have

$$H(X) = H(p) = 1$$

$$H(Y) = H(p * r) = 1 \text{ where } p * r = p(1-r) + r(1-p) = 0.5$$

$$H(X, Y) = H(X, Z)$$

$$= H(X) + H(Z)$$

$$= H(p) + H(r) = 1.47$$

[1E] x 5


$$H(Y|X) = H(\frac{r}{p}) = H(Z) = 0.47$$

each

$$H(X|Y) = H(X, Y) - H(Y)$$

$$= H(p) + H(r) - H(p * r)$$

$$= 0.47$$



[1E]

b)

i) Since it is an erasure channel from X to Y_1 ,

$$C(X \rightarrow Y_1) = 1 - \alpha_1$$

[2A]

ii) It is easy to show that the link from X to Y_2

is an equivalent erasure channel with erasure [2A]


probability $\alpha_1 + \alpha_2 - \alpha_1 \alpha_2$. Hence

$$C(X \rightarrow Y_2) = (1 - \alpha_1)(1 - \alpha_2)$$

$$= 1 - \alpha_1 - \alpha_2 + \alpha_1 \alpha_2$$

[2A]

iii) The equivalent channel from U to Y_2 is depicted as follows



It is easy to write down the transitional prob. [2T]

| $U \backslash Y_2$ | 0 | 1 |
|--------------------|-----------------------|-----------------------|
| 0 | $(1-\beta)(1-\alpha)$ | α |
| 1 | $\beta(1-\alpha)$ | $(1-\beta)(1-\alpha)$ |

and the marginal distribution of Y_2

$$P_{Y_2}(y) = \left\{ \frac{1-\alpha}{2}, \alpha, \frac{1-\alpha}{2} \right\}$$

Hence,

$$R_2 = I(U; Y_2) = H(Y_2) - H(Y_2|U)$$

$$\text{Since } H(Y_2) = H\left(\frac{1-\alpha}{2}, \alpha, \frac{1-\alpha}{2}\right) = H(\alpha) + (1-\alpha)$$

$$H(Y_2|U) = H((1-\beta)(1-\alpha), \alpha, \beta(1-\alpha)) = H(\alpha) + (1-\alpha)H(\beta)$$

We obtain

$$R_2 = (1-\alpha)(1-H(\beta)).$$

Similarly,

$$\begin{aligned} R_1 &= I(X; Y_1|U) \\ &= H(Y_1|U) - H(Y_1|U, X) \\ &= H(Y_1|U) - H(Y_1|X) \end{aligned}$$

$U \rightarrow X \rightarrow Y_1$ is Markov chain

We know

$$H(Y_1|X) = H(\alpha_1)$$

Following the same procedure as that for $H(Y_2|U)$,

$$H(Y_1|U) = H(\alpha_1) + (1-\alpha_1)H(\beta)$$

Thus,

$$R_1 = (1-\alpha_1)H(\beta)$$

Capacity region:

