

6-Field Models (V3.4.4)

Tianyang Xia^{1,2,*}, Jianguo Chen^{2,#}, Xueqiao Xu³

February 9, 2018

¹Institute of Plasma Physics, Chinese Academy of Sciences, Hefei, China

²School of Physics, Peking University, Beijing, 100871, China

³Lawrence Livermore National Laboratory, Livermore, CA 94550, USA

Email: *xiaty@ipp.ac.cn, #cjgls@pku.edu.cn

Contents

1	MHD equations	1
2	Full set of equations	3
2.1	Curvature vector	3
2.2	Radial Force Balance	4
2.3	Vorticity	5
2.4	Density equation	7
2.5	Parallel velocity equations	9
2.6	Ohm's Law	11
2.7	Ion temperature equation	12
2.8	Electron temperature equation	15
2.9	Magnetic Flutter in Parallel Thermal Conduction	17
3	Normalization	19
4	Normalization by \bar{B} (Recommended)	24
5	Results	29

6	Conclusion	29
7	Reduced to 3Field model	34
7.1	Pressure Equation	34
7.2	Ohm's Law	34
7.3	Vorticity Equation	34

1 MHD equations

The original equations we want to use in the simulations are[1]:

$$\frac{\partial}{\partial t} n_i + \nabla \cdot (n_i \mathbf{V}_i) = 0,$$

$$\frac{\partial}{\partial t} (m_j n_j \mathbf{V}_j) + \nabla P_j + \nabla \cdot (\overleftrightarrow{\pi}_j + m_j n_j \mathbf{V}_j \mathbf{V}_j) = Z_j e n_j (\mathbf{E} + \mathbf{V}_j \times \mathbf{B}) + \mathbf{R}_j,$$

$$\frac{\partial}{\partial t} \left(\frac{3}{2} P_j + \frac{1}{2} m_j n_j V_j^2 \right) + \nabla \cdot \mathbf{Q}_j = Z_j e n_j \mathbf{V}_j \cdot \mathbf{E} + \mathbf{V}_i \cdot \mathbf{R}_i + W_j.$$

Here, the energy flux is given by Branginskii

$$\mathbf{Q}_j = \left(\frac{5}{2} k_B n_j T_j + \frac{1}{2} m_j n_j V_j^2 \right) \mathbf{V}_j + \overleftrightarrow{\pi} \cdot \mathbf{V}_j + \mathbf{q}_j$$

with

$$\mathbf{q}_i = -\kappa_{\parallel i} \nabla_{\parallel} T_i + \frac{5P_i}{2m_i \Omega_i} \hat{\mathbf{b}} \times \nabla T_i - \kappa_{\perp i} \nabla_{\perp} T_i \quad (1.1)$$

$$\mathbf{q}_e = -\kappa_{\parallel e} \nabla_{\parallel} T_e + \frac{5P_e}{2m_e \Omega_e} \hat{\mathbf{b}} \times \nabla T_e - \kappa_{\perp e} \nabla_{\perp} T_e - 0.71 \frac{T_e}{e} J_{\parallel} + \frac{3\nu_e}{2\Omega_e} \frac{T_e \hat{\mathbf{b}} \times \mathbf{J}}{e} \quad (1.2)$$

and the friction force

$$\mathbf{R}_e = -\mathbf{R}_i = e n_e \left(\frac{\mathbf{J}_{\parallel}}{\sigma_{\parallel}} + \frac{\mathbf{J}_{\perp}}{\sigma_{\perp}} \right) - 0.71 k_B n_e \nabla_{\parallel} T_e + \frac{3\nu_e}{2\Omega_e} k_B n_e \hat{\mathbf{b}} \times \nabla T_e.$$

The energy exchange term

$$\begin{aligned} W_i &= \frac{3m_e}{m_i} \frac{k_B n_e}{\tau_e} (T_e - T_i), \\ W_e &= -W_i - \mathbf{R}_e \cdot (\mathbf{V}_e - \mathbf{V}_i). \\ \tau_e &= \frac{3\sqrt{m_e} (k_B T_e)^{\frac{3}{2}}}{4\sqrt{2\pi} n_e e^4 \ln \Lambda} \end{aligned}$$

The viscous stress tensor $\overleftrightarrow{\pi}_j$ is composed by three parts, sp-caled “parallel” part $\overleftrightarrow{\pi}_{cj}$, “perpendicular” part $\overleftrightarrow{\pi}_{\perp j}$ and “gyroviscous” part $\overleftrightarrow{\pi}_{gj}$. The “parallel” viscous stress tensor yields magnetic pumping term which can damps the plasma flow shear [2]. The expression of $\overleftrightarrow{\pi}_{cj}$ is given by Ref. [1] [3] as

$$\overleftrightarrow{\pi}_{cj} = \left(\mathbf{b}\mathbf{b} - \frac{1}{3} \overleftrightarrow{\mathbf{I}} \right) \pi_{cj}, \quad (1.3)$$

and

$$\begin{aligned} \pi_{cj} &= \eta_j^0 \left[(\mathbf{V}_E + \mathbf{V}_{Pj}) \cdot \boldsymbol{\kappa} - \frac{2}{\sqrt{B}} \nabla_{\parallel} (\sqrt{B} V_{\parallel j}) \right] \\ &+ \eta_j^0 \left[\left(1.61 k_B \frac{\mathbf{b} \times \nabla T_j}{Z_j e B} \right) \cdot \boldsymbol{\kappa} + \left(\frac{7.09}{5 P_{ij} \sqrt{B}} \right) \nabla_{\parallel} (\sqrt{B} \kappa_{\parallel i} \nabla_{\parallel} T_i) \right] \\ &- \eta_j^0 \left[\frac{2.44 \kappa_{\parallel i} \nabla_{\parallel} T_i}{5 P_j} (2.27 \nabla_{\parallel} \ln T_j - \nabla_{\parallel} \ln P_j) \right] \\ &- \eta_j^0 \left[\left(\frac{\mu_0}{B^3} \right) \mathbf{b} \times \nabla (P_i + P_e) \cdot \left(\nabla \Phi + \frac{\nabla P_j}{Z_j e n} + 1.61 \frac{\nabla T_j}{Z_j e} \right) \right]. \end{aligned} \quad (1.4)$$

The “perpendicular” part is a factor of ν_j/Ω_j smaller than the “gyroviscous” part [1].

For those equations, we can get the momentum and energy of Braginskii equations as

$$m_j n_j \frac{d}{dt} \mathbf{V}_j + \nabla P_j + \nabla \cdot \overleftrightarrow{\pi}_j = Z_j e n_j (\mathbf{E} + \mathbf{V}_j \times \mathbf{B}) + \mathbf{R}_j, \quad (1.5)$$

$$\frac{d}{dt} \left(\frac{3}{2} P_j \right) + \frac{5}{2} P_j \nabla \cdot \mathbf{V}_i = -\overleftrightarrow{\pi}_j : \nabla \mathbf{V}_j - \nabla \cdot \mathbf{q}_j + W_j, \quad (1.6)$$

Here $d/dt = \partial/\partial t + \mathbf{V}_j \cdot \nabla$. The electron velocity V_e is written as

$$V_{\parallel e} = V_{\parallel i} - \frac{J_{\parallel}}{en_e}. \quad (1.7)$$

With the quasineutral condition,

$$Z_i n_i = n_e. \quad (1.8)$$

if the ion and electron have same temperature $T_i = T_e$:

$$P_0 = P_i + P_e = n_i * T_i + n_e * T_e = P_i * (1 + Z_i) \quad (1.9)$$

The linearizing expressions use the definitions of variables as

$$\begin{aligned} n_j &= n_{j0} + n_{j1}, \\ P_j &= P_{j0} + p_{j1}, \\ P = P_i + P_e &= P_0 + p_1 = (P_{i0} + P_{e0}) + (p_{i1} + p_{e1}), \\ \Phi &= \Phi_0 + \phi, \\ J_{\parallel} &= J_{\parallel 0} + J_{\parallel 1}, \\ \mathbf{b} &= \mathbf{b}_0 - \mathbf{b}_0 \times \nabla \psi. \end{aligned}$$

Notice that the flute reduction $k_{\parallel} \sim \epsilon k_{\perp}$ is adopted in this work.

2 Full set of equations

2.1 Curvature vector

The field-line curvature vector κ is defined as

$$\kappa \equiv \mathbf{b} \cdot \nabla \mathbf{b} = (\nabla \times \mathbf{b}) \times \mathbf{b} \quad (2.1.1)$$

due to

$$\nabla(\mathbf{b} \cdot \mathbf{b}) = 2[\mathbf{b} \times (\nabla \times \mathbf{b}) + (\mathbf{b} \cdot \nabla)\mathbf{b}] = 0$$

This can be re-written as

$$\kappa = \left[\frac{1}{B} \nabla \times \mathbf{B} - \mathbf{B} \times \nabla \left(\frac{1}{B} \right) \right] \times \mathbf{b}$$

Using $\nabla \left(\frac{1}{B} \right) = -\frac{\nabla B}{B^2}$, this becomes

$$\begin{aligned} \kappa &= \frac{\mu_0}{B^2} \mathbf{J} \times \mathbf{B} + \frac{1}{B} (\mathbf{b} \times \nabla B) \times \mathbf{b} \\ &= \frac{\mu_0}{B^2} \mathbf{J} \times \mathbf{B} + \frac{1}{B} \nabla_{\perp} B \end{aligned} \quad (2.1.2)$$

$$\begin{aligned}
\mathbf{b} \cdot \nabla \times \mathbf{b} &= \mathbf{b} \cdot \nabla \times \frac{\mathbf{B}}{B} = \mathbf{b} \cdot \left[\frac{1}{B} \nabla \times \mathbf{B} + \frac{1}{B^2} \mathbf{B} \times \nabla B \right] \\
&= \mathbf{b} \cdot \frac{\nabla \times \mathbf{B}}{B}
\end{aligned} \tag{2.1.3}$$

$$\frac{\nabla \times \mathbf{b}}{B} = \frac{1}{B^2} \nabla \times \mathbf{B} + \frac{1}{B^2} \mathbf{b} \times \nabla B \tag{2.1.4}$$

$$\begin{aligned}
\frac{2}{B} \mathbf{b} \times \kappa &= \frac{2}{B} \mathbf{b} \times [(\nabla \times \mathbf{b}) \times \mathbf{b}] \\
&= \frac{2}{B} \nabla \times \mathbf{b} - \frac{2}{B^2} \mathbf{b} \cdot (\nabla \times \mathbf{B}) \mathbf{b}
\end{aligned} \tag{2.1.5}$$

$$\begin{aligned}
\nabla \times \frac{\mathbf{b}}{B} &= \frac{1}{B} \nabla \times \mathbf{b} + \frac{1}{B^2} \mathbf{b} \times \nabla B \\
&= \frac{2}{B} \nabla \times \mathbf{b} - \frac{1}{B^2} \nabla \times \mathbf{B} \\
&= \frac{2}{B} \mathbf{b} \times \kappa - \frac{1}{B^2} \nabla \times \mathbf{B} + \frac{2}{B^2} \mathbf{b} \cdot (\nabla \times \mathbf{B}) \mathbf{b} \\
&= \frac{2}{B} \mathbf{b} \times \kappa + \frac{\mu_0(\mathbf{J}_{\parallel} - \mathbf{J}_{\perp})}{B^2}
\end{aligned} \tag{2.1.6}$$

$$\simeq \frac{2}{B} \mathbf{b} \times \kappa + O(\beta) \tag{2.1.7}$$

For the last step (2.1.7), looking into [1] APPENDIX A for details.

2.2 Radial Force Balance

According to the radial force balance, the radial electric field can be expressed as:

$$E_r = \left(\frac{\nabla P_i}{Z_i e n_i} \right)_r - V_{\theta i} B_{\phi} + V_{\phi i} B_{\theta} \tag{2.2.1}$$

We define part of equilibrium E_{r0} :

$$E_{r,dia0} = -(\nabla \phi_{dia0})_r = \left(\frac{\nabla P_i}{Z_i e n_i} \right)_r \tag{2.2.2}$$

which makes the $\mathbf{E}_{r,dia0} \times \mathbf{B}$ flow balance ion diamagnetic flow:

$$\frac{\mathbf{E}_{r,dia0} \times \mathbf{B}}{B^2} = -\frac{\mathbf{b} \times \nabla P_i}{Z_i e n_i B} \tag{2.2.3}$$

The remained part of E_{r0} is called net $E_{r0,net}$ introducing the net flow:

$$\mathbf{V}_{e0,net} = \frac{\mathbf{E}_{r0,net} \times \mathbf{B}}{B^2} = \frac{\mathbf{B} \times \nabla \Phi_{0,net}}{B^2} \tag{2.2.4}$$

2.3 Vorticity

First, we can add the ion and electron momentum equations together and obtain

$$m_i n_i \frac{d}{dt} \mathbf{V}_i + \nabla \cdot \overleftrightarrow{\boldsymbol{\pi}} + \nabla P = \mathbf{J} \times \mathbf{B}. \quad (2.3.1)$$

Here the electron momentum is neglected for $m_e \ll m_i$ and the viscosity is kept. Multiply the term $\mathbf{b} \cdot \nabla \times$ on Eq. (2.3.1), we derive

$$\begin{aligned} \mathbf{b} \cdot \nabla \times \left[m_i n_i \frac{d}{dt} \mathbf{V}_i + \nabla \cdot \overleftrightarrow{\boldsymbol{\pi}} \right] &= \mathbf{b} \cdot \nabla \times (\mathbf{J} \times \mathbf{B}) \\ &= B^2 \mathbf{b} \cdot \nabla \left(\frac{J_{\parallel}}{B} \right) - \frac{1}{B^2} \mathbf{b} \times (\mathbf{J} \times \mathbf{B}) \cdot \nabla_{\perp} B^2 \\ &= B^2 \mathbf{b} \cdot \nabla \left(\frac{J_{\parallel}}{B} \right) - \frac{1}{\mu_0} \mathbf{b} \times \boldsymbol{\kappa} \cdot \nabla_{\perp} B^2. \end{aligned}$$

Because $\boldsymbol{\kappa} \equiv \mathbf{b} \cdot \nabla \mathbf{b} = \frac{\mu_0}{B^2} \mathbf{J} \times \mathbf{B} + \frac{1}{B} \nabla_{\perp} B$, so the last equation becomes

$$\mathbf{b} \cdot (\nabla - 2\boldsymbol{\kappa}) \times \left[m_i n_i \frac{d}{dt} \mathbf{V}_i + \nabla \cdot \overleftrightarrow{\boldsymbol{\pi}} \right] = B^2 \mathbf{b} \cdot \nabla \left(\frac{J_{\parallel}}{B} \right) + 2\mathbf{b} \times \boldsymbol{\kappa} \cdot \nabla P.$$

Neglect the “parallel” viscous term and the curvature on momentum, then take the time derivative operator out of the spatial differential, the vorticity equation is obtained

$$\begin{aligned} \frac{d}{dt} \varpi &= \frac{\partial}{\partial t} \varpi + (\mathbf{V}_E + \mathbf{V}_{\parallel i} + \mathbf{V}_{Pi}) \cdot \nabla \varpi \\ &= B^2 \mathbf{b} \cdot \nabla \left(\frac{J_{\parallel}}{B} \right) + 2\mathbf{b} \times \boldsymbol{\kappa} \cdot \nabla P_1 \\ &\quad - \frac{1}{2\Omega_i} [n_i Z_i e \mathbf{V}_{Di} \cdot \nabla (\nabla_{\perp}^2 \Phi) - m_i \Omega_i \mathbf{b} \times \nabla n_i \cdot \nabla V_E^2] \\ &\quad + \frac{1}{2\Omega_i} [\mathbf{V}_E \cdot \nabla (\nabla_{\perp}^2 P_i) - \nabla_{\perp}^2 (\mathbf{V}_E \cdot \nabla P_i)]. \end{aligned}$$

Here the vorticity is defined as

$$\begin{aligned} \varpi &= \mathbf{b} \cdot \nabla \times m_i n_i \mathbf{V}_i \\ &\simeq n_{i0} \frac{m_i}{B_0} \left(\nabla_{\perp}^2 \phi + \frac{1}{n_{i0}} \nabla_{\perp} \phi \cdot \nabla_{\perp} n_{i0} + \frac{1}{n_{i0} Z_i e} \nabla_{\perp}^2 p_{i1} \right). \end{aligned} \quad (2.3.2)$$

The ion velocity is composed by $E \times B$ drift \mathbf{V}_E , diamagnetic drift \mathbf{V}_D , polarization drift \mathbf{V}_{Pi} (Eq.(2.4.1)) and perturbed parallel velocity $V_{\parallel i} \mathbf{b}$

$$\begin{aligned}
\mathbf{V}_i &= \mathbf{V}_{\perp i} + V_{\parallel i} \mathbf{b} = \mathbf{V}_E + \mathbf{V}_D + V_{\parallel i} \mathbf{b} + \mathbf{V}_{Pi} = \\
&= \frac{1}{B_0} \mathbf{b}_0 \times \nabla_{\perp} \Phi + \frac{1}{Z_i e n_i B_0} \mathbf{b}_0 \times \nabla_{\perp} P_i + V_{\parallel i} \mathbf{b} + \mathbf{V}_{Pi} \\
&= \frac{1}{B_0} \mathbf{b}_0 \times \nabla_{\perp} \phi + \frac{1}{B_0} \mathbf{b}_0 \times \nabla_{\perp} \Phi_{0,net} + \frac{1}{Z_i e n_i B_0} \mathbf{b}_0 \times \nabla_{\perp} p_{i1} + V_{\parallel i} \mathbf{b}_0 + \mathbf{V}_{Pi} \quad (2.3.3)
\end{aligned}$$

$$\begin{aligned}
\mathbf{V}_{E \times B}^2 &= \left(\frac{\mathbf{b} \times \nabla \Phi}{B} \right)^2 = \left(\frac{\nabla_{\perp} \Phi}{B} \right)^2 = \frac{(\nabla \Phi)^2 - (\nabla_{\parallel} \Phi)^2}{B^2} \\
&= \frac{1}{B^2} [(\nabla \Phi_0)^2 + 2 \nabla \Phi_0 \cdot \nabla \phi + (\nabla \phi)^2 - (\nabla_{\parallel} \Phi_0)^2 - 2 \nabla_{\parallel} \Phi_0 \cdot \nabla_{\parallel} \phi - (\nabla_{\parallel} \phi)^2] \\
&= \frac{1}{B^2} [(\nabla_{\perp} \Phi_0)^2 + 2 \nabla_{\perp} \Phi_0 \cdot \nabla_{\perp} \phi + (\nabla_{\perp} \phi)^2] \quad (2.3.4)
\end{aligned}$$

$$\mathbf{V}_{0,net} = \frac{1}{B_0} \mathbf{b}_0 \times \nabla_{\perp} \Phi_{0,net} \quad (2.3.5)$$

$$\begin{aligned}
\varpi_0 &\simeq n_{i0} \frac{m_i}{B_0} \left(\nabla_{\perp}^2 \Phi_0 + \frac{1}{n_{i0}} \nabla_{\perp} \Phi_0 \cdot \nabla_{\perp} n_{i0} + \frac{1}{n_{i0} Z_i e} \nabla_{\perp}^2 P_{i0} \right) \\
&= n_{i0} \frac{m_i}{B_0} \left(\nabla_{\perp}^2 \Phi_{0,net} + \frac{1}{n_{i0}} \nabla_{\perp} \Phi_0 \cdot \nabla_{\perp} n_{i0} \right) \quad (2.3.6)
\end{aligned}$$

$$\nabla_{\perp} \Phi = \frac{\mathbf{b} \times \nabla_{\perp} \Phi}{B} \times \mathbf{B} = \mathbf{V}_{E \times B} \times \mathbf{B} \quad (2.3.7)$$

Notice that the cancellation of zero-order of $E \times B$ and diamagnetic drift is applied here for equilibrium. For the last step ,we use the linearized expression

$$\begin{aligned}
\frac{\partial}{\partial t} \varpi &= B_0^2 \nabla_{\parallel 0} \left(\frac{J_{\parallel 1}}{B} \right) - B_0^2 \mathbf{b}_0 \times \nabla \psi \cdot \nabla \left(\frac{J_{\parallel 0}}{B_0} \right) + 2 \mathbf{b}_0 \times \boldsymbol{\kappa} \cdot \nabla p_1 \\
&\quad - \frac{1}{B_0} \mathbf{b}_0 \times \nabla_{\perp} \Phi_0 \cdot \nabla \varpi - \frac{1}{B_0} \mathbf{b}_0 \times \nabla_{\perp} \phi \cdot \nabla \varpi_0 - V_{\parallel i} \nabla_{\parallel 0} \varpi_0 \\
&\quad - \frac{1}{2\Omega_i} \frac{1}{B_0} \mathbf{b}_0 \times \nabla p_{i1} \cdot \nabla (\nabla_{\perp}^2 \Phi_0) - \frac{1}{2\Omega_i} \frac{1}{B_0} \mathbf{b}_0 \times \nabla P_{i0} \cdot \nabla (\nabla_{\perp}^2 \phi) \\
&\quad + \frac{1}{2\Omega_i} Z_i e B_0 \mathbf{b}_0 \times \nabla n_{i1} \cdot \nabla \left(\frac{\nabla \Phi_0}{B_0} \right)^2 + \frac{1}{\Omega_i} Z_i e B_0 \mathbf{b}_0 \times \nabla n_{i0} \cdot \nabla \left(\frac{\nabla \Phi_0 \cdot \nabla \phi}{B_0^2} \right) \\
&\quad - \frac{1}{2\Omega_i} Z_i e B_0 \mathbf{b}_0 \times \nabla n_{i1} \cdot \nabla \left(\frac{\nabla_{\parallel} \Phi_0}{B_0} \right)^2 - \frac{1}{\Omega_i} Z_i e B_0 \mathbf{b}_0 \times \nabla n_{i0} \cdot \nabla \left(\frac{\nabla_{\parallel} \Phi_0 \nabla_{\parallel} \phi}{B_0^2} \right) \\
&\quad + \frac{1}{2\Omega_i} \frac{1}{B_0} \mathbf{b}_0 \times \nabla \phi \cdot \nabla (\nabla_{\perp}^2 P_{i0}) + \frac{1}{2\Omega_i} \frac{1}{B_0} \mathbf{b}_0 \times \nabla \Phi_0 \cdot \nabla (\nabla_{\perp}^2 p_{i1}) \\
&\quad - \frac{1}{2\Omega_i} \nabla_{\perp}^2 \left(\frac{1}{B_0} \mathbf{b}_0 \times \nabla \Phi_0 \cdot \nabla p_{i1} \right) - \frac{1}{2\Omega_i} \nabla_{\perp}^2 \left(\frac{1}{B_0} \mathbf{b}_0 \times \nabla \phi \cdot \nabla P_{i0} \right) \\
&\quad - B_0^2 \mathbf{b}_0 \times \nabla \psi \cdot \nabla \left(\frac{J_{\parallel 1}}{B} \right) \\
&\quad - \frac{1}{B_0} \mathbf{b}_0 \times \nabla_{\perp} \phi \cdot \nabla \varpi - V_{\parallel i} \nabla_{\parallel 0} \varpi + V_{\parallel i} \mathbf{b}_0 \times \nabla \psi \cdot \nabla \varpi_0 \\
&\quad - \frac{1}{2\Omega_i} \frac{1}{B_0} \mathbf{b}_0 \times \nabla p_{i1} \cdot \nabla (\nabla_{\perp}^2 \phi) \\
&\quad + \frac{1}{2\Omega_i} Z_i e B_0 \mathbf{b}_0 \times \nabla n_{i0} \cdot \nabla \left(\frac{\nabla \phi}{B_0} \right)^2 + \frac{1}{\Omega_i} Z_i e B_0 \mathbf{b}_0 \times \nabla n_{i1} \cdot \nabla \left(\frac{\nabla \Phi_0 \cdot \nabla \phi}{B_0^2} \right) \\
&\quad - \frac{1}{2\Omega_i} Z_i e B_0 \mathbf{b}_0 \times \nabla n_{i0} \cdot \nabla \left(\frac{\nabla_{\parallel} \phi}{B_0} \right)^2 - \frac{1}{\Omega_i} Z_i e B_0 \mathbf{b}_0 \times \nabla n_{i1} \cdot \nabla \left(\frac{\nabla_{\parallel} \Phi_0 \nabla_{\parallel} \phi}{B_0^2} \right) \\
&\quad + \frac{1}{2\Omega_i} \frac{1}{B_0} \mathbf{b}_0 \times \nabla \phi \cdot \nabla (\nabla_{\perp}^2 p_{i1}) \\
&\quad - \frac{1}{2\Omega_i} \nabla_{\perp}^2 \left(\frac{1}{B_0} \mathbf{b}_0 \times \nabla \phi \cdot \nabla p_{i1} \right). \tag{2.3.8}
\end{aligned}$$

2.4 Density equation

The density equation can be written

$$\frac{\partial}{\partial t} n_i + \mathbf{V}_i \cdot \nabla n_i = -n_i \nabla \cdot \mathbf{V}_i.$$

It is convient to use the exact identities

$$\begin{aligned}
\nabla \cdot \mathbf{V}_E &= \left(\nabla \times \frac{\mathbf{b}}{B} \right) \cdot \nabla \Phi \\
&\simeq \frac{2}{B} \mathbf{b} \times \boldsymbol{\kappa} \cdot \nabla \Phi, \\
\nabla \cdot (n_i \mathbf{V}_D) &= \frac{1}{Z_i e} \left(\nabla \times \frac{\mathbf{b}}{B} \right) \cdot \nabla P_i \\
&\simeq \frac{2}{Z_i e B} \mathbf{b} \times \boldsymbol{\kappa} \cdot \nabla P_i, \\
\nabla \cdot (V_{\parallel i} \mathbf{b}) &= B \nabla_{\parallel} \left(\frac{V_{\parallel i}}{B} \right).
\end{aligned}$$

Here, $\nabla \times \frac{\mathbf{b}}{B} \simeq \frac{2}{B} \mathbf{b} \times \boldsymbol{\kappa}$ (2.1.7). The polarization drift is defined as

$$\mathbf{V}_P = \frac{m_j}{Z_j e B^2} \frac{d\mathbf{E}}{dt} \quad (2.4.1)$$

and it is the small correction to the $E \times B$ and diamagnetic velocities. For ion it is of the order δ^2 and for electron $\sqrt{m_e/m_i}$ smaller than the other two. ????

We can rewrite the density equation as

$$\begin{aligned}
\frac{\partial}{\partial t} n_i &= -\frac{1}{B_0} \mathbf{b} \times \nabla_{\perp} \Phi \cdot \nabla n_i - \frac{2n_i}{B} \mathbf{b} \times \boldsymbol{\kappa} \cdot \nabla_{\perp} \Phi \\
&\quad - \frac{2}{Z_i e B} \mathbf{b} \times \boldsymbol{\kappa} \cdot \nabla_{\perp} P_i \\
&\quad - V_{\parallel i} \mathbf{b} \cdot \nabla n_i - n_i B \nabla_{\parallel} \left(\frac{V_{\parallel i}}{B} \right) \\
&\quad - \nabla \cdot (n_i \mathbf{V}_{Pi})
\end{aligned} \quad (2.4.2)$$

After Linearizing

$$\begin{aligned}
\frac{\partial}{\partial t} n_i &= -\frac{1}{B_0} \mathbf{b}_0 \times \nabla_{\perp} \phi \cdot \nabla n_{i0} - \frac{1}{B_0} \mathbf{b}_0 \times \nabla_{\perp} \Phi_0 \cdot \nabla n_i \\
&\quad - \frac{2n_{i0}}{B_0} \mathbf{b}_0 \times \boldsymbol{\kappa} \cdot \nabla_{\perp} \phi - \frac{2n_i}{B_0} \mathbf{b}_0 \times \boldsymbol{\kappa} \cdot \nabla_{\perp} \Phi_{0,net} \\
&\quad - \frac{2}{Z_i e B_0} \mathbf{b}_0 \times \boldsymbol{\kappa} \cdot \nabla_{\perp} p_{i1}
\end{aligned} \tag{2.4.3}$$

$$\begin{aligned}
&\quad - V_{\parallel i} \nabla_{\parallel 0} n_{i0} - n_{i0} B_0 \nabla_{\parallel 0} \left(\frac{V_{\parallel i}}{B_0} \right) \\
&\quad - \frac{1}{B_0} \mathbf{b}_0 \times \nabla_{\perp} \phi \cdot \nabla n_i - \frac{2n_i}{B_0} \mathbf{b}_0 \times \boldsymbol{\kappa} \cdot \nabla_{\perp} \phi \\
&\quad - V_{\parallel i} \nabla_{\parallel 0} n_i + V_{\parallel i} \mathbf{b}_0 \times \nabla \psi \cdot \nabla n_{i0} \\
&\quad - n_i B_0 \nabla_{\parallel 0} \left(\frac{V_{\parallel i}}{B_0} \right) + n_{i0} B_0 \mathbf{b}_0 \times \nabla \psi \cdot \nabla \left(\frac{V_{\parallel i}}{B_0} \right) \\
&\quad - \nabla \cdot (n_i \mathbf{V}_{Pi}).
\end{aligned} \tag{2.4.4}$$

2.5 Parallel velocity equations

Start from the momentum equation, \mathbf{b} ·Eq. (2.3.1) can get

$$m_i n_i \frac{d}{dt} V_{\parallel i} + \mathbf{b} \cdot \nabla P + \mathbf{b} \cdot (\nabla \cdot \overleftrightarrow{\boldsymbol{\pi}}_i) = 0.$$

Here we neglect $m_i n_i \mathbf{V}_i \cdot \partial \mathbf{b} / \partial t$ and $m_i n_i \nabla \mathbf{b} \cdot \mathbf{V}_i$ for they are smaller than the other terms. Because the “perpendicular” part of viscous stress tensor is much smaller than the other two parts, it is neglected in our discussion. According to Ref. [1], from the expression of parallel viscosity (1.3),

$$\nabla \cdot \overleftrightarrow{\boldsymbol{\pi}}_{ci} = [\mathbf{b} (\nabla \cdot \mathbf{b}) + \boldsymbol{\kappa}] \pi_{ci} + \mathbf{b} \nabla_{\parallel} \pi_{ci} - \frac{1}{3} \nabla \pi_{ci},$$

so that

$$\mathbf{b} \cdot (\nabla \cdot \overleftrightarrow{\boldsymbol{\pi}}_{ci}) = \frac{2}{3} B^{\frac{3}{2}} \nabla_{\parallel} \left(\frac{\pi_{ci}}{B^{\frac{3}{2}}} \right).$$

For the “gyroviscous” contributions, we only consider the case of a straight, homogeneous, time-independent magnetic field and neglect parallel derivatives ($\mathbf{b} \cdot \nabla \rightarrow 0$) to obtain

$$\begin{aligned}
\nabla \cdot \overleftrightarrow{\boldsymbol{\pi}}_{gi} &\simeq -m_i n_i \mathbf{V}_{Di} \cdot \nabla \mathbf{V}_i + \mathbf{b} \times \nabla \left(\frac{P_i}{2\Omega_i} \nabla \cdot \mathbf{V}_{\perp} + \frac{1}{5\Omega_i} \nabla \cdot \mathbf{q}_{\perp i} \right) \\
&\quad + \nabla_{\perp} \left[\frac{P_i}{2\Omega_i} \nabla \cdot (\mathbf{b} \times \mathbf{V}_i) + \frac{1}{5\Omega_i} \nabla \cdot (\mathbf{b} \times \mathbf{q}_i) \right].
\end{aligned} \tag{2.5.1}$$

It is easy to see that

$$\begin{aligned}\frac{P_i}{2\Omega_i}\nabla \cdot \mathbf{V}_\perp + \frac{1}{5\Omega_i}\nabla \cdot \mathbf{q}_{\perp i} &\simeq -\frac{P_i}{2\Omega_i}\mathbf{V}_{Di} \cdot \nabla \ln n_i - \frac{n_i k_B}{2\Omega_i}\mathbf{V}_{Di} \cdot \nabla T_i \\ &\simeq 0,\end{aligned}$$

and

$$\frac{P_i}{2\Omega_i}\nabla \cdot (\mathbf{b} \times \mathbf{V}_i) + \frac{1}{5\Omega_i}\nabla \cdot (\mathbf{b} \times \mathbf{q}_i) \simeq -\frac{1}{2m_i\Omega_i} \left[P_i \nabla_\perp \cdot \left(e \nabla_\perp \Phi + \frac{\nabla_\perp P_i}{n_i} \right) + \nabla_\perp \cdot (P_i \nabla_\perp T_i) \right]$$

So the parallel part of the divergence of the ion “gyroviscous” is

$$\begin{aligned}\nabla \cdot \overleftrightarrow{\boldsymbol{\pi}}_{gi} &\simeq -m_i n_i \mathbf{V}_{Di} \cdot \nabla \mathbf{V}_i - \nabla_\perp \cdot \left\{ \frac{1}{2m_i\Omega_i^2} \left[P_i \nabla_\perp \cdot \left(e \nabla_\perp \Phi + \frac{\nabla_\perp P_i}{n_i} \right) + \nabla_\perp \cdot (P_i \nabla_\perp T_i) \right] \right\} \\ \mathbf{b} \cdot (\nabla \cdot \overleftrightarrow{\boldsymbol{\pi}}_{gi}) &\simeq -m_i n_i \mathbf{V}_{Di} \cdot \nabla V_{\parallel i}.\end{aligned}$$

The parallel divergence of viscous term can be written as

$$\begin{aligned}\mathbf{b} \cdot (\nabla \cdot \overleftrightarrow{\boldsymbol{\pi}}_i) &\simeq \mathbf{b} \cdot (\nabla \cdot \overleftrightarrow{\boldsymbol{\pi}}_{ci}) + \mathbf{b} \cdot (\nabla \cdot \overleftrightarrow{\boldsymbol{\pi}}_{gi}) \\ &\simeq \frac{2}{3} B^{\frac{3}{2}} \nabla_\parallel \left(\frac{\pi_{ci}}{B^{\frac{3}{2}}} \right) - m_i n_i \mathbf{V}_{Di} \cdot \nabla V_{\parallel i}.\end{aligned}$$

Therefore, we can get the parallel velocity as

$$\begin{aligned}\frac{\partial}{\partial t} V_{\parallel i} + \mathbf{V}_i \cdot \nabla V_{\parallel i} &= -\frac{1}{m_i n_i} \mathbf{b} \cdot \nabla P - \mathbf{b} \cdot (\nabla \cdot \overleftrightarrow{\boldsymbol{\pi}}_i), \\ \frac{\partial}{\partial t} V_{\parallel i} &= -\left(\frac{1}{B_0} \mathbf{b} \times \nabla_\perp \Phi + V_{\parallel i} \mathbf{b} \right) \cdot \nabla V_{\parallel i} - \frac{1}{m_i n_i} \mathbf{b} \cdot \nabla P \\ &\quad - \frac{2}{3m_i n_i} B^{\frac{3}{2}} \nabla_\parallel \left(\frac{\pi_{ci}}{B^{\frac{3}{2}}} \right) - \mathbf{V}_{Pi} \cdot \nabla V_{\parallel i}.\end{aligned}\tag{2.5.2}$$

Linearizing derives

$$\begin{aligned}
\frac{\partial}{\partial t} V_{\parallel i} = & -\frac{1}{B_0} \mathbf{b}_0 \times \nabla_{\perp} \Phi_0 \cdot \nabla V_{\parallel i} \\
& -\frac{1}{m_i n_{i0}} \nabla_{\parallel 0} p_1 + \frac{1}{m_i n_i} \mathbf{b}_0 \times \nabla \psi \cdot \nabla P_0 \\
& -\frac{1}{B_0} \mathbf{b}_0 \times \nabla_{\perp} \phi \cdot \nabla V_{\parallel i} - V_{\parallel i} \nabla_{\parallel} V_{\parallel i} \\
& +\frac{1}{m_i n_{i0}} \mathbf{b}_0 \times \nabla \psi \cdot \nabla p_1 \\
& -\mathbf{V}_{Pi} \cdot \nabla V_{\parallel i} - \frac{2}{3m_i n_i} B^{\frac{3}{2}} \nabla_{\parallel} \left(\frac{\pi_{ci}}{B^{\frac{3}{2}}} \right).
\end{aligned} \tag{2.5.3}$$

2.6 Ohm's Law

For the parallel part of Eq. (1.5) for electron,

$$m_e n_e \frac{d}{dt} V_{\parallel e} + \mathbf{b} \cdot \nabla P_e = -en_e E_{\parallel} + R_{\parallel e} - \mathbf{b} \cdot (\nabla \cdot \overleftrightarrow{\pi}_e).$$

Here the electron-ion friction is written as

$$\mathbf{R}_e = en_e \left(\frac{\mathbf{J}_{\parallel}}{\sigma_{\parallel}} + \frac{\mathbf{J}_{\perp}}{\sigma_{\perp}} \right) - 0.71 k_B n_e \nabla_{\parallel} T_e + \frac{3\nu_e}{2\Omega_e} k_B n_e \mathbf{b} \times \nabla T_e.$$

So neglect the inertial terms and viscous stress tensor, we have

$$\nabla_{\parallel} P_e = -en_e \left(-\nabla_{\parallel} \Phi - \frac{\partial}{\partial t} A_{\parallel} \right) + en_e \frac{J_{\parallel 1}}{\sigma_{\parallel}} - 0.71 k_B n_e \nabla_{\parallel} T_e.$$

Then we can obtain the Ohm's law with the substitution $A_{\parallel} = B\psi$ and $J_{\parallel 1} = -\frac{1}{\mu_0} B_0 \nabla_{\perp}^2 \psi$,

$$\frac{\partial}{\partial t} \psi = -\frac{1}{B} \nabla_{\parallel} \Phi + \frac{\eta_{\parallel}}{\mu_0} \nabla_{\perp}^2 \psi + \frac{1}{en_e B} \nabla_{\parallel} P_e + \frac{0.71 k_B}{eB} \nabla_{\parallel} T_e. \tag{2.6.1}$$

After linearized,

$$\begin{aligned}
\frac{\partial}{\partial t} \psi &= -\frac{1}{B_0} \nabla_{\parallel 0} \phi + \frac{1}{B_0} \mathbf{b}_0 \times \nabla \psi \cdot \nabla \Phi_0 \\
&+ \frac{\eta_{\parallel}}{\mu_0} \nabla_{\perp}^2 \psi \\
&+ \frac{1}{en_{e0} B_0} \nabla_{\parallel 0} p_{e1} - \frac{1}{en_e B_0} \mathbf{b}_0 \times \nabla \psi \cdot \nabla P_{e0} \\
&+ \frac{0.71 k_B}{e B_0} \nabla_{\parallel 0} T_{e1} - \frac{0.71 k_B}{e B_0} \mathbf{b}_0 \times \nabla \psi \cdot \nabla T_{e0} \\
&+ \frac{1}{B_0} \mathbf{b}_0 \times \nabla \psi \cdot \nabla \phi - \frac{1}{en_e B_0} \mathbf{b}_0 \times \nabla \psi \cdot \nabla p_{e1} \\
&- \frac{0.71 k_B}{e B_0} \mathbf{b}_0 \times \nabla \psi \cdot \nabla T_{e1}.
\end{aligned} \tag{2.6.2}$$

2.7 Ion temperature equation

For simplicity, we treat the time partial differential on kinetic term simply on velocity since it is 2nd order quantity at least. Then we have

$$\begin{aligned}
\frac{d}{dt} \left(\frac{3}{2} P_j \right) + \frac{5}{2} P_j \nabla \cdot \mathbf{V}_i &= - \langle \vec{\pi}_i : \nabla \mathbf{V}_i - \nabla \cdot \mathbf{q}_j + W_j, \\
\frac{d}{dt} \left(\frac{3}{2} P_i \right) &= -\frac{5}{2} P_i \nabla \cdot \mathbf{V}_i - \langle \vec{\pi}_i : \nabla \mathbf{V}_i - \nabla \cdot \left(\frac{5 P_i}{2 m_i \Omega_i} \mathbf{b} \times \nabla T_i \right) \\
&+ \nabla_{\parallel} (\kappa_{\parallel i} \nabla_{\parallel} T_i) + \nabla_{\perp} (\kappa_{\perp i} \nabla_{\perp} T_i) \\
&+ \frac{3 m_e k_B n_e}{m_i \tau_e} (T_e - T_i).
\end{aligned}$$

The ion temperature can be written as

$$\begin{aligned}
\frac{\partial}{\partial t} T_i + \mathbf{V}_i \cdot \nabla T_i &= -\frac{2}{3} T_i \nabla \cdot \mathbf{V} + \frac{2}{3 n_i k_B} \nabla_{\parallel} (\kappa_{\parallel i} \nabla_{\parallel} T_i) + \frac{2}{3 n_i k_B} \nabla_{\perp} (\kappa_{\perp i} \nabla_{\perp} T_i) \\
&+ \frac{2 m_e Z_i}{m_i \tau_e} (T_e - T_i) - \frac{2}{3 k_B n_i} \langle \vec{\pi}_i : \nabla \mathbf{V}_i - \frac{5}{3 k_B n_i} \nabla \cdot \left(\frac{P_i}{m_i \Omega_i} \hat{\mathbf{b}} \times \nabla T_i \right) \rangle.
\end{aligned}$$

For the gyrofrequency term in the energy flux,

$$\begin{aligned}
\nabla \cdot \left(\frac{5 P_i}{2 m_i \Omega_i} \mathbf{b} \times \nabla T_i \right) &= \frac{5}{2 Z_i e} \left[P_i \nabla \cdot \left(\frac{\mathbf{b} \times \nabla T_i}{B} \right) + \frac{\mathbf{b} \times \nabla T_i}{B} \cdot \nabla P_i \right] \\
&= \frac{5}{2 Z_i e} P_i \frac{2}{B} \mathbf{b} \times \boldsymbol{\kappa} \cdot \nabla T_i - \frac{5}{2} \frac{\mathbf{b} \times \nabla P_i}{Z_i e B} \cdot \nabla T_i
\end{aligned}$$

The last term of RHS is canceled by both the convectional term and the $\nabla \cdot \mathbf{V}_{Di}$.

For the viscous term, the “perpendicular” part is neglected for ions, then we can write the “parallel” viscous term as

$$\begin{aligned}
\overleftrightarrow{\pi}_{ci} : \nabla \mathbf{V}_i &= \pi_{ci} \left(\mathbf{b}\mathbf{b} - \frac{1}{3} \overleftrightarrow{\mathbf{I}} \right) : \nabla \mathbf{V}_i \\
&\simeq \frac{\pi_{ci}}{3} [2\nabla_{\parallel} V_{\parallel i} - 3\boldsymbol{\kappa} \cdot \mathbf{V}_{\perp i} - \nabla \cdot \mathbf{V}_{\perp i} - V_{\parallel i} (\nabla \cdot \mathbf{b})] \\
&\simeq \frac{\pi_{ci}}{3} \left[\frac{2}{\sqrt{B}} \nabla_{\parallel} \left(\sqrt{B} V_{\parallel i} \right) - \boldsymbol{\kappa} \cdot (\mathbf{V}_E + \mathbf{V}_{Di}) - \frac{k_B}{Z_i e n_i B} \mathbf{b} \cdot \nabla n_i \times \nabla T_i \right] \\
&\quad + \frac{\pi_{ci}}{3} \left[\frac{\mu_0}{B^3} \mathbf{b} \times \nabla P \cdot \nabla \phi + \frac{\mu_0}{Z_i e n_i B^3} \mathbf{b} \times \nabla P_e \cdot \nabla P_i \right].
\end{aligned} \tag{2.7.1}$$

with the leading order expression. For simplicity, the last two terms of RHS can be dropped since they come from the second term of RHS of

$$\left(\nabla \times \frac{\mathbf{b}}{B} \right) = \frac{2}{B} \mathbf{b} \times \boldsymbol{\kappa} - \frac{\mu_0 \mathbf{b} \times \nabla P}{B^3} + O(\beta).$$

For the calculation of viscous term, the approximation of streight and homogeneous magnetic field is adopted, thus Eq. (2.7.1) can be written as

$$\overleftrightarrow{\pi}_{ci} : \nabla \mathbf{V}_i = \frac{\pi_{ci}}{3} \left[\frac{2}{\sqrt{B}} \nabla_{\parallel} \left(\sqrt{B} V_{\parallel i} \right) - \frac{k_B}{Z_i e n_i B} \mathbf{b} \cdot \nabla n_i \times \nabla T_i \right]$$

The “gyroviscous” term have the relations as

$$\overleftrightarrow{\pi}_{gi} \cdot \mathbf{V}_i \simeq \frac{P_i}{2\Omega_i} \mathbf{b} \times \nabla V_{\parallel i}^2.$$

Within Eq. (2.5.1), we can get

$$\begin{aligned}
\overleftrightarrow{\pi}_{gi} : \nabla \mathbf{V}_i &\simeq \nabla \cdot (\overleftrightarrow{\pi}_{gi} \cdot \mathbf{V}_i) - (\nabla \cdot \overleftrightarrow{\pi}_{gi}) \cdot \mathbf{V}_i \\
&\simeq \frac{m_i}{Z_i e} P_i V_{\parallel i} \frac{2}{B} \mathbf{b} \times \boldsymbol{\kappa} \cdot \nabla V_{\parallel i} - m_i n_i V_{\parallel i} \mathbf{V}_{Di} \cdot \nabla V_{\parallel i} \\
&\quad + m_i n_i V_{\parallel i} \mathbf{V}_{Di} \cdot \nabla V_{\parallel i} + \mathbf{V}_{\perp i} \cdot \nabla_{\perp} \left(\frac{1}{2m_i \Omega_i^2} A_{\pi gi} \right),
\end{aligned}$$

where

$$A_{\pi gi} = eP_i \nabla_{\perp}^2 \phi + k_B T_i \nabla_{\perp}^2 P_i - k_B T_i \nabla_{\perp} P_i \cdot \frac{\nabla_{\perp} n_i}{n_i} + k_B \nabla_{\perp} P_i \cdot \nabla_{\perp} T_i + k_B P_i \nabla_{\perp}^2 T_i.$$

According to Ref. [1], we drop the $\overleftrightarrow{\pi}_{gi} : \nabla V_i$ and obtain

$$\overleftrightarrow{\pi}_{gi} : \nabla V_i \simeq \frac{m_i}{Z_i e} P_i V_{\parallel i} \frac{2}{B} \mathbf{b} \times \boldsymbol{\kappa} \cdot \nabla V_{\parallel i}$$

After all, the temperature equation can be written as

$$\begin{aligned} \frac{\partial}{\partial t} T_i = & - \left(\frac{1}{B_0} \mathbf{b} \times \nabla_{\perp} \Phi + V_{\parallel i} \mathbf{b} \right) \cdot \nabla T_i \\ & - \frac{2}{3} T_i \left[\left(\frac{2}{B} \mathbf{b} \times \boldsymbol{\kappa} \right) \cdot \left(\nabla \Phi + \frac{1}{Z_i e n_i} \nabla P_i + \frac{5}{2} \frac{k_B}{Z_i e} \nabla T_i \right) + B \nabla_{\parallel} \left(\frac{V_{\parallel i}}{B} \right) \right] \\ & + \frac{2}{3 n_i k_B} \nabla_{\parallel} (\kappa_{\parallel i} \nabla_{\parallel} T_i) + \frac{2}{3 n_i k_B} \nabla_{\perp} (\kappa_{\perp i} \nabla_{\perp} T_i) \\ & + \frac{2 m_e}{m_i} \frac{Z_i}{\tau_e} (T_e - T_i) \\ & - \frac{2 \pi c_i}{9 k_B n_i} \left[\frac{2}{\sqrt{B}} \nabla_{\parallel} (\sqrt{B} V_{\parallel i}) - \frac{k_B}{Z_i e n_i B} \mathbf{b} \cdot \nabla n_i \times \nabla T_i \right] \\ & - \frac{4}{3 \Omega_i} T_i V_{\parallel i} \mathbf{b} \times \boldsymbol{\kappa} \cdot \nabla V_{\parallel i} \\ & - \frac{2}{3} T_i \nabla \cdot \mathbf{V}_{Pi} - \mathbf{V}_{Pi} \cdot \nabla T_i. \end{aligned}$$

Linearizing

$$\begin{aligned}
\frac{\partial}{\partial t} T_i = & -\frac{1}{B_0} \mathbf{b}_0 \times \nabla_{\perp} \phi \cdot \nabla T_{i0} - \frac{1}{B_0} \mathbf{b}_0 \times \nabla_{\perp} \Phi_0 \cdot \nabla T_{i1} - V_{\parallel i} \nabla_{\parallel 0} T_{i0} \\
& -\frac{2}{3} T_{i0} \left[\left(\frac{2}{B_0} \mathbf{b}_0 \times \boldsymbol{\kappa} \right) \cdot \left(\nabla \phi + \frac{1}{Z_i e n_{i0}} \nabla p_{i1} + \frac{5}{2} \frac{k_B}{Z_i e} \nabla T_{i1} \right) + B_0 \nabla_{\parallel 0} \left(\frac{V_{\parallel i}}{B_0} \right) \right] \\
& -\frac{2}{3} T_{i1} \left(\frac{2}{B_0} \mathbf{b}_0 \times \boldsymbol{\kappa} \right) \cdot \left(\nabla \Phi_0 + \frac{1}{Z_i e n_{i0}} \nabla P_{i0} + \frac{5}{2} \frac{k_B}{Z_i e} \nabla T_{i0} \right) \\
& + \frac{2}{3 n_{i0} k_B} \nabla_{\parallel} (\boldsymbol{\kappa}_{\parallel i} \nabla_{\parallel} T_{i1}) + \frac{2}{3 n_{i0} k_B} \nabla_{\perp} (\boldsymbol{\kappa}_{\perp i} \nabla_{\perp} T_{i1}) \\
& + \frac{2 m_e}{m_i} \frac{Z_i}{\tau_e} (T_{e1} - T_{i1}) \\
& -\frac{1}{B_0} \mathbf{b}_0 \times \nabla_{\perp} \phi \cdot \nabla T_{i1} - V_{\parallel i} \nabla_{\parallel 0} T_{i1} + V_{\parallel i} \mathbf{b}_0 \times \nabla \psi \cdot \nabla T_{i0} \\
& -\frac{2}{3} T_{i1} \left[\left(\frac{2}{B_0} \mathbf{b}_0 \times \boldsymbol{\kappa} \right) \cdot \left(\nabla \phi + \frac{1}{Z_i e n_{i0}} \nabla p_{i1} + \frac{5}{2} \frac{k_B}{Z_i e} \nabla T_{i1} \right) + B_0 \nabla_{\parallel 0} \left(\frac{V_{\parallel i}}{B_0} \right) \right] \\
& + \frac{2}{3} T_{i0} B_0 \mathbf{b}_0 \times \nabla \psi \cdot \nabla \left(\frac{V_{\parallel i}}{B_0} \right) \\
& - \frac{2}{3 k_B n_i} \overleftrightarrow{\boldsymbol{\pi}}_i : \nabla \mathbf{V}_i \\
& + \frac{2}{3} T_i \nabla \cdot \mathbf{V}_{Pi} - \mathbf{V}_{Pi} \cdot \nabla T_i
\end{aligned} \tag{2.7.2}$$

2.8 Electron temperature equation

For electron temperature, the Braginskii equation is

$$\begin{aligned}
\frac{3}{2} n_e \frac{d}{dt} (k_B T_e) = & - P_e \nabla \cdot \mathbf{V}_e - \nabla \cdot \left[-0.71 \frac{k_B T_e \mathbf{J}_{\parallel}}{e} + \frac{5 P_e}{2 e B} \mathbf{b} \times \nabla T_e + \frac{3 k_B \nu_e}{2 \Omega_e} \frac{T_e \mathbf{b} \times \mathbf{J}}{e} \right] \\
& + \nabla_{\parallel} (\boldsymbol{\kappa}_{\parallel e} \nabla_{\parallel} T_e) + \nabla_{\perp} (\boldsymbol{\kappa}_{\perp e} \nabla_{\perp} T_e) \\
& - \frac{3 m_e}{m_i} \frac{n_e k_B}{\tau_e} (T_e - T_i) - \mathbf{R}_e \cdot (\mathbf{V}_e - \mathbf{V}_i) \\
& - \frac{2}{3 k_B n_e} \overleftrightarrow{\boldsymbol{\pi}}_e : \nabla \mathbf{V}_e.
\end{aligned}$$

For the terms in energy flux,

$$\begin{aligned}
\nabla \cdot \left[-0.71 \frac{k_B T_e \mathbf{J}_{\parallel}}{e} + \frac{3 k_B \nu_e}{2 \Omega_e} \frac{T_e \mathbf{b} \times \mathbf{J}}{e} \right] = & -0.71 \frac{k_B T_e}{e} B \nabla_{\parallel} \left(\frac{J_{\parallel}}{B} \right) - 0.71 \frac{k_B J_{\parallel}}{e} \nabla_{\parallel} T_e \\
& + \frac{3 k_B m_e \nu_e T_e}{e^2 B^2} \mathbf{b} \times \boldsymbol{\kappa} \cdot \nabla P + \frac{3 k_B m_e}{2 e^2 B} \mathbf{b} \times \nabla P \cdot \nabla \left(\frac{\nu_e T_e}{B} \right)
\end{aligned}$$

Notice that the last two terms in the RHD of last equation has the order of $\nu_e/\Omega_e \sim \epsilon\sqrt{m_e/m_i} \ll 1$ as the first two, so they can be neglected in the following calculations in flute reduction. The friction force term in the energy exchange term can be written as

$$\begin{aligned}
\mathbf{R}_e \cdot (\mathbf{V}_e - \mathbf{V}_i) &= \left[Z_i e n_i \left(\frac{\mathbf{J}_\parallel}{\sigma_\parallel} + \frac{\mathbf{J}_\perp}{\sigma_\perp} \right) - 0.71 k_B n_e \nabla_\parallel T_e + \frac{3\nu_e}{2\Omega_e} k_B n_e \mathbf{b} \times \nabla T_e \right] \\
&\quad \cdot \left[-\frac{1}{Z_i e n_i B} \mathbf{b} \times \nabla P - \frac{1}{Z_i e n_i} J_\parallel \mathbf{b}_\parallel \right] \\
&= -\eta_\parallel J_\parallel^2 + 0.71 \frac{k_B}{e} J_\parallel \nabla_\parallel T_e - \frac{\eta_\perp}{B} \mathbf{b} \times \nabla P \cdot \mathbf{J}_\perp - \frac{3\nu_e k_B}{2\Omega_e e B} (\mathbf{b} \times \nabla T_e) \cdot (\mathbf{b} \times \nabla P) \\
&\simeq -\eta_\parallel J_\parallel^2 + 0.71 \frac{k_B}{e} J_\parallel \nabla_\parallel T_e.
\end{aligned}$$

Notice that the polarization drift is neglected here for it is much smaller than the other velocities. The term of perpendicular current is zero because $\mathbf{J}_\perp \simeq \nabla P/B$. The last term with gyrofrequency is also dropped for the order $\nu_e/\Omega_e \sim \epsilon\sqrt{m_e/m_i}$. Then Neglect the electron viscosity, we have

$$\begin{aligned}
\frac{\partial}{\partial t} T_e + \mathbf{V}_e \cdot \nabla T_e &= -\frac{2}{3} T_e \nabla \cdot \mathbf{V}_e + 0.71 \frac{2T_e}{3en_e} B \nabla_\parallel \left(\frac{J_\parallel}{B} \right) \\
&\quad + \frac{2}{3n_e k_B} \nabla_\parallel (\kappa_{\parallel e} \nabla_\parallel T_e) + \frac{2}{3n_e k_B} \nabla_\perp (\kappa_{\perp e} \nabla_\perp T_e) \\
&\quad - \frac{2m_e}{m_i} \frac{1}{\tau_e} (T_e - T_i) + \frac{2}{3n_e k_B} \eta_\parallel J_\parallel^2,
\end{aligned}$$

and

$$\begin{aligned}
\frac{\partial}{\partial t} T_e &= -\left(\frac{1}{B_0} \mathbf{b} \times \nabla_\perp \Phi + V_{\parallel e} \mathbf{b} \right) \cdot \nabla T_e \\
&\quad - \frac{2}{3} T_e \left[\left(\frac{2}{B} \mathbf{b} \times \boldsymbol{\kappa} \right) \cdot \left(\nabla \Phi - \frac{1}{en_e} \nabla P_e - \frac{5}{2} \frac{k_B}{e} \nabla T_e \right) + B \nabla_\parallel \left(\frac{V_{\parallel e}}{B} \right) \right] \\
&\quad + 0.71 \frac{2T_e}{3en_e} B \nabla_\parallel \left(\frac{J_\parallel}{B} \right) \\
&\quad + \frac{2}{3n_e k_B} \nabla_\parallel (\kappa_{\parallel e} \nabla_\parallel T_e) + \frac{2}{3n_e k_B} \nabla_\parallel (\kappa_{\perp e} \nabla_\perp T_e) \\
&\quad - \frac{2m_e}{m_i} \frac{1}{\tau_e} (T_e - T_i) + \frac{2}{3n_e k_B} \eta_\parallel J_\parallel^2 \\
&\quad - \frac{2}{3} T_e \nabla \cdot \mathbf{V}_{Pe} - \mathbf{V}_{Pe} \cdot \nabla T_e.
\end{aligned}$$

After linearizing

$$\begin{aligned}
\frac{\partial}{\partial t} T_e = & -\frac{1}{B_0} \mathbf{b}_0 \times \nabla_{\perp} \phi \cdot \nabla T_{e0} - \frac{1}{B_0} \mathbf{b}_0 \times \nabla_{\perp} \Phi_0 \cdot \nabla T_{e1} - V_{\parallel e} \nabla_{\parallel 0} T_{e0} \\
& -\frac{2}{3} T_{e0} \left[\left(\frac{2}{B_0} \mathbf{b}_0 \times \boldsymbol{\kappa} \right) \cdot \left(\nabla \phi - \frac{1}{en_{e0}} \nabla p_{e1} - \frac{5}{2} \frac{k_B}{e} \nabla T_{e1} \right) + B_0 \nabla_{\parallel 0} \left(\frac{V_{\parallel e}}{B_0} \right) \right] \\
& -\frac{2}{3} T_{e1} \left(\frac{2}{B_0} \mathbf{b}_0 \times \boldsymbol{\kappa} \right) \cdot \left(\nabla \Phi_0 - \frac{1}{en_{e0}} \nabla P_{e0} - \frac{5}{2} \frac{k_B}{e} \nabla T_{e0} \right) \\
& +0.71 \frac{2}{3e} \frac{T_{e0}}{n_{e0}} B_0 \nabla_{\parallel 0} \left(\frac{J_{\parallel 1}}{B_0} \right) + 0.71 \frac{2}{3e} \frac{T_{e1}}{n_{e0}} B_0 \nabla_{\parallel 0} \left(\frac{J_{\parallel 0}}{B_0} \right) \\
& -0.71 \frac{2}{3e} \frac{T_{e0}}{n_{e0}} B_0 \mathbf{b}_0 \times \nabla \psi \cdot \nabla \left(\frac{J_{\parallel 0}}{B_0} \right) \\
& +\frac{2}{3n_{e0}k_B} \nabla_{\parallel} (\kappa_{\parallel e} \nabla_{\parallel} T_{e1}) + \frac{2}{3n_{e0}k_B} \nabla_{\perp} (\kappa_{\perp e} \nabla_{\perp} T_{e1}) \\
& -\frac{2m_e}{m_i} \frac{1}{\tau_e} (T_{e1} - T_{i1}) + \frac{4}{3n_{e0}k_B} \eta_{\parallel} J_{\parallel 0} J_{\parallel 1} \\
& -\frac{1}{B_0} \mathbf{b}_0 \times \nabla_{\perp} \phi \cdot \nabla T_{e1} - V_{\parallel e} \nabla_{\parallel 0} T_{e1} + V_{\parallel e} \mathbf{b}_0 \times \nabla \psi \cdot \nabla T_{e0} \\
& -\frac{2}{3} T_{e1} \left[\left(\frac{2}{B_0} \mathbf{b}_0 \times \boldsymbol{\kappa} \right) \cdot \left(\nabla \phi - \frac{1}{en_{e0}} \nabla p_{e1} - \frac{5}{2} \frac{k_B}{e} \nabla T_{e1} \right) + B_0 \nabla_{\parallel 0} \left(\frac{V_{\parallel e}}{B_0} \right) \right] \\
& +\frac{2}{3} T_{e0} B_0 \mathbf{b}_0 \times \nabla \psi \cdot \nabla \left(\frac{V_{\parallel e}}{B_0} \right) \\
& +0.71 \frac{2}{3e} \frac{T_{e1}}{n_{e0}} B_0 \nabla_{\parallel 0} \left(\frac{J_{\parallel 1}}{B_0} \right) - 0.71 \frac{2}{3e} \frac{T_{e0}}{n_{e0}} B_0 \mathbf{b}_0 \times \nabla \psi \cdot \nabla \left(\frac{J_{\parallel 1}}{B_0} \right) \\
& -0.71 \frac{2}{3e} \frac{T_{e1}}{n_{e0}} B_0 \mathbf{b}_0 \times \nabla \psi \cdot \nabla \left(\frac{J_{\parallel 0}}{B_0} \right) + \frac{2}{3n_{e0}k_B} \eta_{\parallel} J_{\parallel 1}^2 \\
& -\frac{2}{3} T_e \nabla \cdot \mathbf{V}_{Pe} - \mathbf{V}_{Pe} \cdot \nabla T_e
\end{aligned} \tag{2.8.1}$$

2.9 Magnetic Flutter in Parallel Thermal Conduction

The magnetic flutter has strong impact on the distribution of heat fluxes towards divertor targets[4]. Considering the magnetic field unit vector:

$$\mathbf{b} = \mathbf{b}_0 + \mathbf{b}_1 = \mathbf{b}_0 + \nabla A_{\parallel} \times \mathbf{b}_0 / B \simeq \mathbf{b}_0 - \mathbf{b}_0 \times \nabla \psi \tag{2.9.1}$$

$$\nabla_{\parallel} = \mathbf{b} \cdot \nabla = \mathbf{b}_0 \cdot \nabla - \mathbf{b}_0 \times \nabla \psi \cdot \nabla = \nabla_{\parallel 0} - B_0 [\psi, \cdot] \tag{2.9.2}$$

the higher order (>1) of the thermal conduction in temperature equations (2.7.2) and (2.8.1) can be expressed as:

$$\begin{aligned}
\nabla_{\parallel}(\kappa_{\parallel j} \nabla_{\parallel} T_j) &= \nabla_{\parallel 0}(\kappa_{\parallel j} \nabla_{\parallel 0} T_{j1}) \\
&\quad - \mathbf{b}_0 \times \nabla \psi \cdot \nabla [\kappa_{\parallel j} \nabla_{\parallel 0} T_{j1}] - \nabla_{\parallel 0} [\kappa_{\parallel j} \mathbf{b}_0 \times \nabla \psi \cdot \nabla (T_{j0} + T_{j1})] \\
&\quad + \mathbf{b}_0 \times \nabla \psi \cdot \nabla [\kappa_{\parallel j} \mathbf{b}_0 \times \nabla \psi \cdot \nabla (T_{j0} + T_{j1})] \\
&= \nabla_{\parallel 0} \kappa_{\parallel j} \nabla_{\parallel 0} T_{j1} + \kappa_{\parallel j} \nabla_{\parallel 0}^2 T_{j1} \\
&\quad - \mathbf{b}_0 \times \nabla \psi \cdot \nabla \kappa_{\parallel j} (\nabla_{\parallel 0} T_{j1}) - \kappa_{\parallel j} \mathbf{b}_0 \times \nabla \psi \cdot \nabla (\nabla_{\parallel 0} T_{j1}) \\
&\quad - \nabla_{\parallel 0} \kappa_{\parallel j} [\mathbf{b}_0 \times \nabla \psi \cdot \nabla (T_{j0} + T_{j1})] - \kappa_{\parallel j} \nabla_{\parallel 0} [\mathbf{b}_0 \times \nabla \psi \cdot \nabla (T_{j0} + T_{j1})] \\
&\quad + \mathbf{b}_0 \times \nabla \psi \cdot \nabla \kappa_{\parallel j} [\mathbf{b}_0 \times \nabla \psi \cdot \nabla (T_{j0} + T_{j1})] \\
&\quad + \kappa_{\parallel j} \mathbf{b}_0 \times \nabla \psi \cdot \nabla [\mathbf{b}_0 \times \nabla \psi \cdot \nabla (T_{j0} + T_{j1})]
\end{aligned} \tag{2.9.3}$$

NOTE: The equilibrium temperature profiles are flux functions $T_{j0}(\psi)$, so $\nabla_{\parallel 0} T_{j0} = 0$.

3 Normalization

For numerical simulation, the normalization are necessary

$$\begin{aligned}
\hat{T}_j &= \frac{T_j}{\bar{T}_j}, & \hat{n} &= \frac{n_i}{\bar{n}}, & \hat{L} &= \frac{L}{\bar{L}}, \\
\hat{t} &= \frac{t}{\bar{t}}, & \hat{B} &= \frac{B}{\bar{B}}, & \hat{J} &= \frac{\mu_0 \bar{L}}{B_0} J, \\
\hat{\psi} &= \frac{\psi}{\bar{L}}, & \hat{\phi} &= \frac{\bar{t}}{\bar{L}^2 \bar{B}_0} \phi, & \hat{\omega} &= \frac{\bar{t}}{m_i \bar{n}} \omega, \\
\tau &= \frac{\bar{T}_i}{\bar{T}_e}, & \hat{V} &= \frac{V}{V_A}, & V_A &= \frac{\bar{L}}{\bar{t}} = \frac{\bar{B}}{\sqrt{\mu_0 m_i n_i}},
\end{aligned} \tag{3.1}$$

$$\hat{P}_j = \frac{P_j}{k_B \bar{n} \bar{T}_j}, \quad \hat{\kappa} = \bar{L} \kappa, \quad \hat{\nabla} = \bar{L} \nabla \tag{3.2}$$

so we have

$$\begin{aligned}
\hat{P} = (\tau \hat{P}_i + \hat{P}_e) &= \frac{P}{(k_B \bar{n} \bar{T}_e)}, \\
&= \tau \hat{P}_i * (1 + Z_i), \text{ if } T_i = T_e
\end{aligned} \tag{3.3}$$

$$\hat{E} = \frac{E}{V_A \bar{B}} = -\hat{\nabla}(\hat{B}_0 \hat{\phi}) \tag{3.4}$$

$$\hat{\mathbf{V}}_{E \times B} = \frac{\mathbf{V}_{E \times B}}{V_A} = \frac{\hat{\mathbf{E}}_r \times \hat{\mathbf{B}}_0}{\hat{B}_0^2} \simeq \mathbf{b}_0 \times \hat{\nabla} \hat{\phi} \tag{3.5}$$

The last step in Eq.(3.8), the B_0 is assumed constant.

$$\hat{E} = \frac{E}{V_A \bar{B} \hat{B}_0} = -\frac{\hat{\nabla}(\hat{B}_0 \hat{\phi})}{\hat{B}_0} \simeq -\hat{\nabla} \hat{\phi} \tag{3.6}$$

$$\hat{\mathbf{V}}_{E \times B} = \frac{\mathbf{V}_{E \times B}}{V_A} = \frac{\hat{\mathbf{E}}_r \times \hat{\mathbf{B}}_0}{\hat{B}_0} \simeq \mathbf{b}_0 \times \hat{\nabla} \hat{\phi} \tag{3.7}$$

$$\hat{\nabla}_\perp(\hat{B}_0 \hat{\Phi}_0) = \left[\mathbf{b}_0 \times \hat{\nabla}(\hat{B}_0 \hat{\Phi}_0) \right] \times \mathbf{b}_0 = \hat{\mathbf{V}}_{E \times B} \times \mathbf{B}_0 \tag{3.8}$$

NOTE: In the code of version that using normalization factor B_0 , in some place the B_0 is considered as constant value. In the gyroviscous terms in vorticity equation, for example:

$$\left[\frac{\hat{\nabla}_\perp(\hat{B}_0 \hat{\Phi}_0)}{\hat{B}_0} \right]^2 \simeq \left[\hat{\nabla}_\perp \hat{\Phi}_0 \right]^2 \tag{3.9}$$

The linearized equations are

$$\hat{\omega} = \mathbf{b} \cdot \hat{\nabla} \times \hat{n}_i \hat{V}_i \simeq \frac{\hat{n}_{i0}}{\hat{B}_0} \left[\hat{\nabla}_\perp^2 (\hat{B}_0 \hat{\phi}) + \frac{1}{\hat{n}_{i0}} \hat{\nabla}_\perp (\hat{B}_0 \hat{\phi}) \cdot \hat{\nabla}_\perp \hat{n}_{i0} + \frac{k_B \bar{T}_i}{Z_i e \bar{L} \bar{B} V_A} \frac{1}{\hat{n}_{i0}} \hat{\nabla}_\perp^2 \hat{p}_{i1} \right] \quad (3.10)$$

$$U_{\text{para0}} = \frac{k_B \bar{T}_i}{Z_i e \bar{B} \bar{L} V_A}, \quad U_{\text{para1}} = \frac{k_B \bar{T}_e}{m_i V_A^2}, \quad U_{\text{para2}} = \frac{k_B \bar{T}_i}{Z_i e \bar{B} \bar{L} V_A}, \quad U_{\text{para3}} = 1.0 \quad (3.11)$$

$$\begin{aligned} \frac{\partial}{\partial t} \hat{\omega} &= \hat{B}_0^2 \hat{\nabla}_{\parallel 0} \hat{J}_{\parallel 1} - \hat{B}_0^2 \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \hat{J}_{\parallel 0} \\ &+ 2 \frac{k_B \bar{T}_e}{m_i V_A^2} \mathbf{b}_0 \times \hat{\kappa} \cdot \hat{\nabla} \hat{p}_1 - \frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla}_\perp (\hat{B}_0 \hat{\Phi}_0) \cdot \hat{\nabla} \hat{\omega} \\ &- \frac{k_B \bar{T}_i}{2 Z_i e \bar{B} \bar{L} V_A} \frac{1}{\hat{B}_0^2} \mathbf{b}_0 \times \hat{\nabla} \hat{p}_{i1} \cdot \hat{\nabla} \left[\hat{\nabla}_\perp^2 (\hat{B}_0 \hat{\Phi}_0) \right] \\ &- \frac{k_B \bar{T}_i}{2 Z_i e \bar{B} \bar{L} V_A} \frac{1}{\hat{B}_0^2} \mathbf{b}_0 \times \hat{\nabla} \hat{P}_{i0} \cdot \hat{\nabla} \left[\hat{\nabla}_\perp^2 (\hat{B}_0 \hat{\phi}) \right] \\ &+ \frac{1}{2} \mathbf{b}_0 \times \hat{\nabla} \hat{n}_{i1} \cdot \hat{\nabla} \left[\frac{\hat{\nabla}_\perp (\hat{B}_0 \hat{\Phi}_0)}{\hat{B}_0} \right]^2 + \mathbf{b}_0 \times \hat{\nabla} \hat{n}_{i0} \cdot \hat{\nabla} \left[\frac{\hat{\nabla}_\perp (\hat{B}_0 \hat{\Phi}_0) \cdot \hat{\nabla}_\perp (\hat{B}_0 \hat{\phi})}{\hat{B}_0^2} \right] \\ &+ \frac{k_B \bar{T}_i}{2 Z_i e \bar{B} \bar{L} V_A} \frac{1}{\hat{B}_0^2} \mathbf{b}_0 \times \hat{\nabla} (\hat{B}_0 \hat{\phi}) \cdot \hat{\nabla} (\hat{\nabla}_\perp^2 \hat{P}_{i0}) + \frac{k_B \bar{T}_i}{2 Z_i e \bar{B} \bar{L} V_A} \frac{1}{\hat{B}_0^2} \mathbf{b}_0 \times \hat{\nabla} (\hat{B}_0 \hat{\Phi}_0) \cdot \hat{\nabla} (\hat{\nabla}_\perp^2 \hat{p}_{i1}) \\ &- \frac{k_B \bar{T}_i}{2 Z_i e \bar{B} \bar{L} V_A} \frac{1}{\hat{B}_0} \hat{\nabla}_\perp^2 \left[\frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla} (\hat{B}_0 \hat{\Phi}_0) \cdot \hat{\nabla} \hat{p}_{i1} \right] \\ &- \frac{k_B \bar{T}_i}{2 Z_i e \bar{B} \bar{L} V_A} \frac{1}{\hat{B}_0} \hat{\nabla}_\perp^2 \left[\frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla} (\hat{B}_0 \hat{\phi}) \cdot \hat{\nabla} \hat{P}_{i0} \right] \\ &- \hat{B}_0^2 \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \hat{J}_{\parallel 1} \\ &- \frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla}_\perp (\hat{B}_0 \hat{\phi}) \cdot \hat{\nabla} \hat{\omega} - \hat{V}_{\parallel i} \hat{\nabla}_{\parallel 0} \hat{\omega} \\ &- \frac{k_B \bar{T}_i}{2 Z_i e \bar{B} \bar{L} V_A} \frac{1}{\hat{B}_0^2} \mathbf{b}_0 \times \hat{\nabla} \hat{p}_{i1} \cdot \hat{\nabla} \left[\hat{\nabla}_\perp^2 (\hat{B}_0 \hat{\phi}) \right] + \frac{1}{2} \mathbf{b}_0 \times \hat{\nabla} \hat{n}_{i0} \cdot \hat{\nabla} \left[\frac{\hat{\nabla}_\perp (\hat{B}_0 \hat{\phi})}{\hat{B}_0} \right]^2 \\ &+ \mathbf{b}_0 \times \hat{\nabla} \hat{n}_{i1} \cdot \hat{\nabla} \left[\frac{\hat{\nabla}_\perp (\hat{B}_0 \hat{\Phi}_0) \cdot \hat{\nabla}_\perp (\hat{B}_0 \hat{\phi})}{\hat{B}_0^2} \right] \\ &+ \frac{k_B \bar{T}_i}{2 Z_i e \bar{B} \bar{L} V_A} \frac{1}{\hat{B}_0^2} \mathbf{b}_0 \times \hat{\nabla} (\hat{B}_0 \hat{\phi}) \cdot \hat{\nabla} (\hat{\nabla}_\perp^2 \hat{p}_{i1}) \\ &- \frac{k_B \bar{T}_i}{2 Z_i e \bar{B} \bar{L} V_A} \frac{1}{\hat{B}_0} \hat{\nabla}_\perp^2 \left(\frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla} (\hat{B}_0 \hat{\phi}) \cdot \hat{\nabla} \hat{p}_{i1} \right), \end{aligned}$$

$$Ni_{\text{para}1} = \frac{k_B \bar{T}_i}{Z_i e \bar{B} \bar{L} V_A}, \quad Vi_{\text{para}} = \frac{\mu_0 k_B \bar{T}_e \bar{n}}{\bar{B}^2} = \frac{k_B \bar{T}_e}{m_i V_A^2}, \quad (3.12)$$

$$\begin{aligned} \frac{\partial}{\partial t} \hat{n}_i &= -\frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla}_\perp \left(\hat{B}_0 \hat{\phi} \right) \cdot \hat{\nabla} \hat{n}_{i0} \\ &\quad -\frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla}_\perp \left(\hat{B}_0 \hat{\Phi}_0 \right) \cdot \hat{\nabla} n_i - \hat{V}_{\parallel i} \hat{\nabla}_{\parallel 0} \hat{n}_{i0} \\ &\quad -\frac{2\hat{n}_{i0}}{\hat{B}_0} \mathbf{b}_0 \times \hat{\kappa} \cdot \hat{\nabla}_\perp \left(\hat{B}_0 \hat{\phi} \right) - \frac{2\hat{n}_i}{\hat{B}_0} \mathbf{b}_0 \times \hat{\kappa} \cdot \hat{\nabla}_\perp \left(\hat{B}_0 \hat{\Phi}_{0,net} \right) \\ &\quad -\frac{2k_B \bar{T}_e}{Z_i e \bar{B} \bar{L} V_A} \frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\kappa} \cdot \hat{\nabla}_\perp \hat{p}_{i1} - \hat{n}_{i0} \hat{B}_0 \hat{\nabla}_{\parallel 0} \left(\frac{V_{\parallel i}}{\hat{B}_0} \right) \\ &\quad -\frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla}_\perp \left(\hat{B}_0 \hat{\phi} \right) \cdot \hat{\nabla} \hat{n}_i - \hat{V}_{\parallel i} \hat{\nabla}_{\parallel 0} \hat{n}_i + \hat{V}_{\parallel i} \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \hat{n}_{i0} \\ &\quad -\frac{2\hat{n}_i}{\hat{B}_0} \mathbf{b}_0 \times \hat{\kappa} \cdot \hat{\nabla}_\perp \left(\hat{B}_0 \hat{\phi} \right) \\ &\quad -\hat{n}_i \hat{B}_0 \hat{\nabla}_{\parallel 0} \left(\frac{\hat{V}_{\parallel i}}{\hat{B}_0} \right) + \hat{n}_{i0} \hat{B}_0 \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \left(\frac{\hat{V}_{\parallel i}}{\hat{B}_0} \right), \\ \frac{\partial}{\partial t} \hat{V}_{\parallel i} &= -\frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla}_\perp \left(\hat{B}_0 \hat{\Phi}_0 \right) \cdot \hat{\nabla} \hat{V}_{\parallel i} \\ &\quad -\frac{k_B \bar{T}_e}{m_i V_A^2} \frac{1}{\hat{n}_{i0}} \hat{\nabla}_{\parallel 0} \hat{p}_1 + \frac{k_B \bar{T}_e}{m_i V_A^2} \frac{1}{\hat{n}_{i0}} \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \hat{P}_0 \\ &\quad -\frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla}_\perp \left(\hat{B}_0 \hat{\phi} \right) \cdot \hat{\nabla} \hat{V}_{\parallel i} \\ &\quad +\frac{k_B \bar{T}_e}{m_i V_A^2} \frac{1}{\hat{n}_{i0}} \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \hat{p}_1, \end{aligned}$$

$$\begin{aligned}
\psi_{\text{para1}} &= \frac{k_B \bar{T}_e}{e \bar{B} \bar{L} V_A}, \\
Ti_{\text{para1}} &= \frac{2}{3} \frac{1}{\bar{L} V_A}, \quad Ti_{\text{para2}} = \frac{k_B \bar{T}_i}{Z_i e \bar{B} \bar{L} V_A},
\end{aligned} \tag{3.13}$$

$$\begin{aligned}
\frac{\partial}{\partial \hat{t}} \hat{\psi} &= -\frac{1}{\hat{B}_0} \hat{\nabla}_{\parallel 0} \left(\hat{B}_0 \hat{\phi} \right) + \frac{\eta_{\parallel}}{\mu_0 V_A \bar{L}} \hat{\nabla}_{\perp}^2 \hat{\psi} \\
&+ \frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla}_{\perp} \left(\hat{B}_0 \hat{\Phi}_0 \right) \\
&+ \frac{k_B \bar{T}_e}{e \bar{B} \bar{L} V_A} \frac{1}{\hat{n}_{e0} \hat{B}_0} \hat{\nabla}_{\parallel 0} \hat{p}_{e1} - \frac{k_B \bar{T}_e}{e \bar{B} \bar{L} V_A} \frac{1}{\hat{n}_{e0} \hat{B}_0} \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \hat{P}_{e0} \\
&+ \frac{k_B \bar{T}_e}{e \bar{B} \bar{L} V_A} \frac{0.71}{\hat{B}_0} \hat{\nabla}_{\parallel 0} \hat{T}_{e1} - \frac{k_B \bar{T}_e}{e \bar{B} \bar{L} V_A} \frac{0.71}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \hat{T}_{e0} \\
&+ \frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla}_{\perp} \left(\hat{B}_0 \hat{\phi} \right) - \frac{k_B \bar{T}_e}{e \bar{B} \bar{L} V_A} \frac{1}{\hat{n}_{e0} \hat{B}_0} \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \hat{p}_{e1} \\
&- \frac{k_B \bar{T}_e}{e \bar{B} \bar{L} V_A} \frac{0.71}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \hat{T}_{e1}, \\
\frac{\partial}{\partial \hat{t}} \hat{T}_i &= -\frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla}_{\perp} \left(\hat{B}_0 \hat{\phi} \right) \cdot \hat{\nabla} \hat{T}_{i0} - \frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla}_{\perp} \left(\hat{B}_0 \hat{\Phi}_0 \right) \cdot \hat{\nabla} \hat{T}_{i1} - \hat{V}_{\parallel i} \hat{\nabla}_{\parallel 0} \hat{T}_{i0} \\
&- \frac{2}{3} \hat{T}_{i0} \left(\frac{2}{\hat{B}_0} \mathbf{b}_0 \times \hat{\kappa} \right) \cdot \left[\hat{\nabla} \left(\hat{B}_0 \hat{\phi} \right) + \frac{k_B \bar{T}_i}{Z_i e \bar{B} \bar{L} V_A} \frac{1}{\hat{n}_{i0}} \hat{\nabla} \hat{p}_{i1} + \frac{5 k_B \bar{T}_i}{2 Z_i e \bar{B} \bar{L} V_A} \hat{\nabla} \hat{T}_{i1} \right] \\
&- \frac{2}{3} \hat{T}_{i0} \hat{B}_0 \hat{\nabla}_{\parallel 0} \left(\frac{\hat{V}_{\parallel i}}{\hat{B}_0} \right) \\
&- \frac{2}{3} \hat{T}_{i1} \left(\frac{2}{\hat{B}_0} \mathbf{b}_0 \times \hat{\kappa} \right) \cdot \left[\hat{\nabla} \left(\hat{B}_0 \hat{\Phi}_0 \right) + \frac{k_B \bar{T}_i}{Z_i e \bar{B} \bar{L} V_A} \frac{1}{\hat{n}_{i0}} \hat{\nabla} \hat{P}_{i0} + \frac{5 k_B \bar{T}_i}{2 Z_i e \bar{B} \bar{L} V_A} \hat{\nabla} \hat{T}_{i0} \right] \\
&+ \frac{2}{3 \bar{n}_i \bar{L} V_A} \frac{1}{\hat{n}_{i0}} \hat{\nabla}_{\parallel} \left(\kappa_{\parallel i} \hat{\nabla}_{\parallel} \hat{T}_{i1} \right) + \frac{2}{3 \bar{n}_i \bar{L} V_A} \frac{1}{\hat{n}_{i0}} \hat{\nabla}_{\perp} \left(\kappa_{\perp i} \hat{\nabla}_{\perp} \hat{T}_{i1} \right) \\
&+ \frac{2 Z_i m_e \bar{t}}{m_i \tau_e} \left(\hat{T}_{e1} - \hat{T}_{i1} \right) \\
&- \frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla}_{\perp} \left(\hat{B}_0 \hat{\phi} \right) \cdot \hat{\nabla} \hat{T}_{i1} - \hat{V}_{\parallel i} \hat{\nabla}_{\parallel 0} \hat{T}_{i1} + \hat{V}_{\parallel i} \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \hat{T}_{i0} \\
&- \frac{2}{3} \hat{T}_{i1} \left(\frac{2}{\hat{B}_0} \mathbf{b}_0 \times \hat{\kappa} \right) \cdot \left[\hat{\nabla} \left(\hat{B}_0 \hat{\phi} \right) + \frac{k_B \bar{T}_i}{Z_i e \bar{B} \bar{L} V_A} \frac{1}{\hat{n}_{i0}} \hat{\nabla} \hat{p}_{i1} + \frac{5 k_B \bar{T}_i}{2 Z_i e \bar{B} \bar{L} V_A} \hat{\nabla} \hat{T}_{i1} \right] \\
&- \frac{2}{3} \hat{T}_{i1} \hat{B}_0 \hat{\nabla}_{\parallel 0} \left(\frac{V_{\parallel i}}{\hat{B}_0} \right) + \frac{2}{3} \hat{T}_{i0} \hat{B}_0 \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \left(\frac{\hat{V}_{\parallel i}}{\hat{B}_0} \right),
\end{aligned}$$

$$T_{e\text{para}1} = \frac{2}{3} \frac{1}{Z_i \bar{L} V_A}, \quad T_{e\text{para}2} = \frac{k_B \bar{T}_e}{e \bar{B} \bar{L} V_A}, \quad T_{e\text{para}3} = \frac{\bar{B}}{e \mu_0 \bar{n}_e \bar{L} V_A}, \quad T_{e\text{para}4} = \frac{\bar{B}^2}{\mu_0 k_B \bar{n}_e \bar{T}_e}, \quad (3.14)$$

$$\begin{aligned} \frac{\partial}{\partial t} \hat{T}_e &= -\frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla}_\perp \left(\hat{B}_0 \hat{\phi} \right) \cdot \hat{\nabla} \hat{T}_{e0} - \frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla}_\perp \left(\hat{B}_0 \hat{\Phi}_0 \right) \cdot \hat{\nabla} \hat{T}_{e1} - \hat{V}_{\parallel e} \hat{\nabla}_{\parallel 0} \hat{T}_{e0} \\ &\quad - \frac{2}{3} \hat{T}_{e0} \left(\frac{2}{\hat{B}_0} \mathbf{b}_0 \times \hat{\kappa} \right) \cdot \left[\hat{\nabla} \left(\hat{B}_0 \hat{\phi} \right) - \frac{k_B \bar{T}_e}{e \bar{B} \bar{L} V_A} \frac{1}{\hat{n}_{e0}} \hat{\nabla} \hat{p}_{e1} - \frac{5 k_B \bar{T}_e}{2 e \bar{B} \bar{L} V_A} \hat{\nabla} \hat{T}_{e1} \right] \\ &\quad - \frac{2}{3} \hat{T}_{e0} \hat{B}_0 \hat{\nabla}_{\parallel 0} \left(\frac{\hat{V}_{\parallel e}}{\hat{B}_0} \right) \\ &\quad - \frac{2}{3} \hat{T}_{e1} \left(\frac{2}{\hat{B}_0} \mathbf{b}_0 \times \hat{\kappa} \right) \cdot \left[\hat{\nabla} \left(\hat{B}_0 \hat{\Phi}_0 \right) - \frac{k_B \bar{T}_e}{e \bar{B} \bar{L} V_A} \frac{1}{\hat{n}_{e0}} \hat{\nabla} \hat{p}_{e0} - \frac{5 k_B \bar{T}_e}{2 e \bar{B} \bar{L} V_A} \hat{\nabla} \hat{T}_{e0} \right] \\ &\quad + 0.71 \frac{2 \bar{B}}{3 e \mu_0 \bar{n}_e \bar{L} V_A} \frac{\hat{T}_{e0}}{\hat{n}_{e0}} \hat{B}_0 \hat{\nabla}_{\parallel 0} \hat{J}_{\parallel 1} + 0.71 \frac{2 \bar{B}}{3 e \mu_0 \bar{n}_e \bar{L} V_A} \frac{\hat{T}_{e1}}{\hat{n}_{e0}} \hat{B}_0 \hat{\nabla}_{\parallel 0} \hat{J}_{\parallel 0} \\ &\quad - 0.71 \frac{2 \bar{B}}{3 e \mu_0 \bar{n}_e \bar{L} V_A} \frac{\hat{T}_{e0}}{\hat{n}_{e0}} \hat{B}_0 \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla}_{\parallel 0} \hat{J}_{\parallel 0} \\ &\quad + \frac{2}{3 \bar{n}_e \bar{L} V_A} \frac{1}{\hat{n}_{e0}} \hat{\nabla}_{\parallel} \left(\kappa_{\parallel e} \hat{\nabla}_{\parallel} \hat{T}_{e1} \right) + \frac{2}{3 \bar{n}_e \bar{L} V_A} \frac{1}{\hat{n}_{e0}} \hat{\nabla}_\perp \left(\kappa_{\perp e} \hat{\nabla}_\perp \hat{T}_{e1} \right) \\ &\quad - \frac{2 m_e \bar{t}}{m_i \tau_e} \left(\hat{T}_{e1} - \hat{T}_{i1} \right) + \frac{4 \bar{B}^2}{3 \mu_0 \bar{n}_e k_B \bar{T}_e} \frac{\hat{B}_0^2}{\hat{n}_{e0}} \hat{\eta}_{\parallel} \hat{J}_{\parallel 0} \hat{J}_{\parallel 1} \\ &\quad - \frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla}_\perp \left(\hat{B}_0 \hat{\phi} \right) \cdot \hat{\nabla} \hat{T}_{e1} - \hat{V}_{\parallel e} \hat{\nabla}_{\parallel 0} \hat{T}_{e1} + \hat{V}_{\parallel e} \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \hat{T}_{e0} \\ &\quad - \frac{2}{3} \hat{T}_{e1} \left(\frac{2}{\hat{B}_0} \mathbf{b}_0 \times \hat{\kappa} \right) \cdot \left[\hat{\nabla} \left(\hat{B}_0 \hat{\phi} \right) - \frac{k_B \bar{T}_e}{e \bar{B} \bar{L} V_A} \frac{1}{\hat{n}_{e0}} \hat{\nabla} \hat{p}_{e1} - \frac{5 k_B \bar{T}_e}{2 e \bar{B} \bar{L} V_A} \hat{\nabla} \hat{T}_{e1} \right] \\ &\quad - \frac{2}{3} \hat{T}_{e1} \hat{B}_0 \hat{\nabla}_{\parallel 0} \left(\frac{\hat{V}_{\parallel e}}{\hat{B}_0} \right) \\ &\quad + \frac{2}{3} \hat{T}_{e0} \hat{B}_0 \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \left(\frac{\hat{V}_{\parallel e}}{\hat{B}_0} \right) \\ &\quad + 0.71 \frac{2 \bar{B}}{3 e \mu_0 \bar{n}_e \bar{L} V_A} \frac{\hat{T}_{e1}}{\hat{n}_{e0}} \hat{B}_0 \hat{\nabla}_{\parallel 0} \hat{J}_{\parallel 1} - 0.71 \frac{2 \bar{B}}{3 e \mu_0 \bar{n}_e \bar{L} V_A} \frac{\hat{T}_{e0}}{\hat{n}_{e0}} \hat{B}_0 \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \hat{J}_{\parallel 1} \\ &\quad - 0.71 \frac{2 \bar{B}}{3 e \mu_0 \bar{n}_e \bar{L} V_A} \frac{\hat{T}_{e1}}{\hat{n}_{e0}} \hat{B}_0 \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \hat{J}_{\parallel 0} \\ &\quad + \frac{2 \bar{B}^2}{3 \mu_0 \bar{n}_e k_B \bar{T}_e} \frac{\hat{B}_0^2}{\hat{n}_{e0}} \hat{\eta}_{\parallel} \hat{J}_{\parallel 1}^2. \end{aligned}$$

4 Normalization by \bar{B} (Recommended)

Using \bar{B} in the normalization factors in ϕ and J .

$$\begin{aligned}
 \hat{T}_j &= \frac{T_j}{\bar{T}_j}, & \hat{n} &= \frac{n_i}{\bar{n}}, & \hat{L} &= \frac{L}{\bar{L}}, \\
 \hat{t} &= \frac{t}{\bar{t}}, & \hat{B} &= \frac{B}{\bar{B}}, & \hat{J} &= \frac{\mu_0 \bar{L}}{\bar{B}} J, \\
 \hat{\psi} &= \frac{\psi}{\bar{L}}, & \hat{\phi} &= \frac{\bar{t}}{\bar{L}^2 \bar{B}} \phi, & \hat{\omega} &= \frac{\bar{t}}{m_i \bar{n}} \omega, \\
 \tau &= \frac{\bar{T}_i}{\bar{T}_e}, & \hat{V} &= \frac{V}{V_A}, & V_A &= \frac{\bar{L}}{\bar{t}} = \frac{\bar{B}}{\sqrt{\mu_0 m_i n_i}}, \\
 \hat{P}_j &= \frac{P_j}{k_B \bar{n} \bar{T}_j}, & \hat{\kappa} &= \bar{L} \kappa, & \hat{\nabla} &= \bar{L} \nabla \\
 \hat{\eta}_{\parallel} &= \frac{\eta_{\parallel}}{\mu_0 V_A \bar{L}}, & \hat{\kappa}_{\parallel, \perp} &= \frac{2\kappa_{\parallel, \perp}}{3V_A \bar{L} \bar{n}},
 \end{aligned}$$

so we have

$$\begin{aligned}
 \hat{P} = (\tau \hat{P}_i + \hat{P}_e) &= \frac{P}{(k_B \bar{n} T_e)}, \\
 &= \tau \hat{P}_i * (1 + Z_i), \text{ if } T_i = T_e
 \end{aligned} \tag{4.1}$$

$$\hat{E} = \frac{E}{V_A \bar{B}} = -\hat{\nabla} \hat{\phi} \tag{4.2}$$

$$\hat{\mathbf{V}}_{E \times B} = \frac{\mathbf{V}_{E \times B}}{V_A} = \frac{\hat{\mathbf{E}}_r \times \hat{\mathbf{B}}_0}{\hat{B}_0^2} = \boxed{\frac{\mathbf{b}_0 \times \hat{\nabla} \hat{\phi}}{\hat{B}_0}} \tag{4.3}$$

$$\hat{\nabla}_{\perp} \hat{\Phi}_0 = \left[\mathbf{b}_0 \times \hat{\nabla} \hat{\Phi}_0 \right] \times \mathbf{b}_0 = \hat{\mathbf{V}}_{E \times B} \times \mathbf{B}_0 \tag{4.4}$$

After ignoring polarization drift (2.4.1) \mathbf{V}_P and taking $\mathbf{b} \cdot \nabla = \mathbf{b}_0 \cdot \nabla - \mathbf{b}_0 \times \nabla \psi \cdot \nabla$, The linearized equations are

$$\hat{\omega} = \mathbf{b} \cdot \hat{\nabla} \times \hat{n}_i \hat{V}_i \simeq \frac{\hat{n}_{i0}}{\hat{B}_0} \left(\hat{\nabla}_\perp^2 \hat{\phi} + \frac{1}{\hat{n}_{i0}} \hat{\nabla}_\perp \hat{\phi} \cdot \hat{\nabla}_\perp \hat{n}_{i0} + \frac{k_B \bar{T}_i}{Z_i e \bar{B} \bar{L} V_A} \frac{1}{\hat{n}_{i0}} \hat{\nabla}_\perp^2 \hat{p}_{i1} \right) \quad (4.5)$$

$$U_{\text{para0}} = \frac{k_B \bar{T}_i}{Z_i e \bar{B} \bar{L} V_A}, \quad U_{\text{para1}} = \frac{k_B \bar{T}_e}{m_i V_A^2}, \quad U_{\text{para2}} = \frac{k_B \bar{T}_i}{Z_i e \bar{B} \bar{L} V_A}, \quad U_{\text{para3}} = 1.0 \quad (4.6)$$

$$\begin{aligned} \frac{\partial}{\partial t} \hat{\omega} = & \hat{B}_0^2 \hat{\nabla}_{\parallel 0} \left(\frac{\hat{J}_{\parallel 1}}{\hat{B}_0} \right) - \hat{B}_0^2 \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \left(\frac{\hat{J}_{\parallel 0}}{\hat{B}_0} \right) + 2 \frac{k_B \bar{T}_e}{m_i V_A^2} \mathbf{b}_0 \times \hat{\kappa} \cdot \hat{\nabla} \hat{p}_1 \\ & - \frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla}_\perp \hat{\Phi}_0 \cdot \hat{\nabla} \hat{\omega} - \frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla}_\perp \hat{\phi} \cdot \hat{\nabla} \hat{\omega}_0 - \hat{V}_{\parallel i} \hat{\nabla}_{\parallel 0} \hat{\omega}_0 \\ & - \frac{k_B \bar{T}_i}{2 Z_i e \bar{B} \bar{L} V_A} \frac{1}{\hat{B}_0^2} \mathbf{b}_0 \times \hat{\nabla} \hat{p}_{i1} \cdot \hat{\nabla} \left[\hat{\nabla}_\perp^2 \hat{\Phi}_0 \right] - \frac{k_B \bar{T}_i}{2 Z_i e \bar{B} \bar{L} V_A} \frac{1}{\hat{B}_0^2} \mathbf{b}_0 \times \hat{\nabla} \hat{P}_{i0} \cdot \hat{\nabla} \left[\hat{\nabla}_\perp^2 \hat{\phi} \right] \\ & + \frac{1}{2} \mathbf{b}_0 \times \hat{\nabla} \hat{n}_{i1} \cdot \hat{\nabla} \left[\frac{\hat{\nabla}_\perp \hat{\Phi}_0}{\hat{B}_0} \right]^2 + \mathbf{b}_0 \times \hat{\nabla} \hat{n}_{i0} \cdot \hat{\nabla} \left[\frac{\hat{\nabla}_\perp \hat{\Phi}_0 \cdot \hat{\nabla}_\perp \hat{\phi}}{\hat{B}_0^2} \right] \\ & + \frac{k_B \bar{T}_i}{2 Z_i e \bar{B} \bar{L} V_A} \frac{1}{\hat{B}_0^2} \mathbf{b}_0 \times \hat{\nabla} \hat{\phi} \cdot \hat{\nabla} \left(\hat{\nabla}_\perp^2 \hat{P}_{i0} \right) + \frac{k_B \bar{T}_i}{2 Z_i e \bar{B} \bar{L} V_A} \frac{1}{\hat{B}_0^2} \mathbf{b}_0 \times \hat{\nabla} \hat{\Phi}_0 \cdot \hat{\nabla} \left(\hat{\nabla}_\perp^2 \hat{p}_{i1} \right) \\ & - \frac{k_B \bar{T}_i}{2 Z_i e \bar{B} \bar{L} V_A} \frac{1}{\hat{B}_0} \hat{\nabla}_\perp^2 \left[\frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla} \hat{\Phi}_0 \cdot \hat{\nabla} \hat{p}_{i1} \right] - \frac{k_B \bar{T}_i}{2 Z_i e \bar{B} \bar{L} V_A} \frac{1}{\hat{B}_0} \hat{\nabla}_\perp^2 \left[\frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla} \hat{\phi} \cdot \hat{\nabla} \hat{P}_{i0} \right] \\ & - \hat{B}_0^2 \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \left(\frac{\hat{J}_{\parallel 1}}{\hat{B}_0} \right) \\ & - \frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla}_\perp \hat{\phi} \cdot \hat{\nabla} \hat{\omega} - \hat{V}_{\parallel i} \hat{\nabla}_{\parallel 0} \hat{\omega} + \hat{V}_{\parallel i} \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \hat{\omega}_0 \\ & - \frac{k_B \bar{T}_i}{2 Z_i e \bar{B} \bar{L} V_A} \frac{1}{\hat{B}_0^2} \mathbf{b}_0 \times \hat{\nabla} \hat{p}_{i1} \cdot \hat{\nabla} \left[\hat{\nabla}_\perp^2 \hat{\phi} \right] \\ & + \frac{1}{2} \mathbf{b}_0 \times \hat{\nabla} \hat{n}_{i0} \cdot \hat{\nabla} \left[\frac{\hat{\nabla}_\perp \hat{\phi}}{\hat{B}_0} \right]^2 + \mathbf{b}_0 \times \hat{\nabla} \hat{n}_{i1} \cdot \hat{\nabla} \left[\frac{\hat{\nabla}_\perp \hat{\Phi}_0 \cdot \hat{\nabla}_\perp \hat{\phi}}{\hat{B}_0^2} \right] \\ & + \frac{k_B \bar{T}_i}{2 Z_i e \bar{B} \bar{L} V_A} \frac{1}{\hat{B}_0^2} \mathbf{b}_0 \times \hat{\nabla} \hat{\phi} \cdot \hat{\nabla} \left(\hat{\nabla}_\perp^2 \hat{p}_{i1} \right) \\ & - \frac{k_B \bar{T}_i}{2 Z_i e \bar{B} \bar{L} V_A} \frac{1}{\hat{B}_0} \hat{\nabla}_\perp^2 \left(\frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla} \hat{\phi} \cdot \hat{\nabla} \hat{p}_{i1} \right), \end{aligned}$$

$$Ni_{\text{para}1} = \frac{k_B \bar{T}_i}{Z_i e \bar{B} \bar{L} V_A}, \quad Vi_{\text{para}} = \frac{\mu_0 k_B \bar{T}_e \bar{n}}{\bar{B}^2} = \frac{k_B \bar{T}_e}{m_i V_A^2}, \quad (4.7)$$

$$\begin{aligned} \frac{\partial}{\partial t} \hat{n}_i &= -\frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla}_\perp \hat{\phi} \cdot \hat{\nabla} \hat{n}_{i0} - \frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla}_\perp \hat{\Phi}_0 \cdot \hat{\nabla} n_i \\ &\quad - \frac{2\hat{n}_{i0}}{\hat{B}_0} \mathbf{b}_0 \times \hat{\kappa} \cdot \hat{\nabla}_\perp \hat{\phi} - \frac{2\hat{n}_i}{\hat{B}_0} \mathbf{b}_0 \times \hat{\kappa} \cdot \hat{\nabla}_\perp \hat{\Phi}_{0,net} \\ &\quad - \frac{2k_B \bar{T}_e}{Z_i e \bar{B} \bar{L} V_A} \frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\kappa} \cdot \hat{\nabla}_\perp \hat{p}_{i1} \\ &\quad - \hat{V}_{\parallel i} \hat{\nabla}_{\parallel 0} \hat{n}_{i0} - \hat{n}_{i0} \hat{B}_0 \hat{\nabla}_{\parallel 0} \left(\frac{V_{\parallel i}}{\hat{B}_0} \right) \\ &\quad - \frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla}_\perp \hat{\phi} \cdot \hat{\nabla} \hat{n}_i - \frac{2\hat{n}_i}{\hat{B}_0} \mathbf{b}_0 \times \hat{\kappa} \cdot \hat{\nabla}_\perp \hat{\phi} \\ &\quad - \hat{V}_{\parallel i} \hat{\nabla}_{\parallel 0} \hat{n}_i + \hat{V}_{\parallel i} \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \hat{n}_{i0} \\ &\quad - \hat{n}_i \hat{B}_0 \hat{\nabla}_{\parallel 0} \left(\frac{\hat{V}_{\parallel i}}{\hat{B}_0} \right) + \hat{n}_{i0} \hat{B}_0 \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \left(\frac{\hat{V}_{\parallel i}}{\hat{B}_0} \right), \\ \frac{\partial}{\partial t} \hat{V}_{\parallel i} &= -\frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla}_\perp \hat{\Phi}_0 \cdot \hat{\nabla} \hat{V}_{\parallel i} \\ &\quad - \frac{k_B \bar{T}_e}{m_i V_A^2} \frac{1}{\hat{n}_{i0}} \hat{\nabla}_{\parallel 0} \hat{p}_1 + \frac{k_B \bar{T}_e}{m_i V_A^2} \frac{1}{\hat{n}_{i0}} \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \hat{P}_0 \\ &\quad - \frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla}_\perp \hat{\phi} \cdot \hat{\nabla} \hat{V}_{\parallel i} \\ &\quad + \frac{k_B \bar{T}_e}{m_i V_A^2} \frac{1}{\hat{n}_{i0}} \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \hat{p}_1, \end{aligned}$$

$$\begin{aligned}
\psi_{\text{para1}} &= \frac{k_B \bar{T}_e}{e \bar{B} \bar{L} V_A}, \\
Ti_{\text{para1}} &= \frac{2}{3} \frac{1}{\bar{L} V_A}, \quad Ti_{\text{para2}} = \frac{k_B \bar{T}_i}{Z_i e \bar{B} \bar{L} V_A},
\end{aligned} \tag{4.8}$$

$$\begin{aligned}
\frac{\partial}{\partial t} \hat{\psi} &= -\frac{1}{\hat{B}_0} \hat{\nabla}_{\parallel 0} \hat{\phi} + \frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla}_{\perp} \hat{\Phi}_0 + \frac{\eta_{\parallel}}{\mu_0 V_A \bar{L}} \left(\frac{\hat{J}_{\parallel 1}}{\hat{B}_0} \right) \\
&+ \frac{k_B \bar{T}_e}{e \bar{B} \bar{L} V_A} \frac{1}{\hat{n}_{e0} \hat{B}_0} \hat{\nabla}_{\parallel 0} \hat{p}_{e1} - \frac{k_B \bar{T}_e}{e \bar{B} \bar{L} V_A} \frac{1}{\hat{n}_{e0} \hat{B}_0} \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \hat{P}_{e0} \\
&+ \frac{k_B \bar{T}_e}{e \bar{B} \bar{L} V_A} \frac{0.71}{\hat{B}_0} \hat{\nabla}_{\parallel 0} \hat{T}_{e1} - \frac{k_B \bar{T}_e}{e \bar{B} \bar{L} V_A} \frac{0.71}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \hat{T}_{e0} \\
&+ \frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla}_{\perp} \hat{\phi} - \frac{k_B \bar{T}_e}{e \bar{B} \bar{L} V_A} \frac{1}{\hat{n}_{e0} \hat{B}_0} \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \hat{p}_{e1} \\
&- \frac{k_B \bar{T}_e}{e \bar{B} \bar{L} V_A} \frac{0.71}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \hat{T}_{e1}, \\
\frac{\partial}{\partial t} \hat{T}_i &= -\frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla}_{\perp} \hat{\phi} \cdot \hat{\nabla} \hat{T}_{i0} - \frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla}_{\perp} \hat{\Phi}_0 \cdot \hat{\nabla} \hat{T}_{i1} - \hat{V}_{\parallel i} \hat{\nabla}_{\parallel 0} \hat{T}_{i0} \\
&- \frac{2}{3} \hat{T}_{i0} \left(\frac{2}{\hat{B}_0} \mathbf{b}_0 \times \hat{\kappa} \right) \cdot \left[\hat{\nabla} \hat{\phi} + \frac{k_B \bar{T}_i}{Z_i e \bar{B} \bar{L} V_A} \frac{1}{\hat{n}_{i0}} \hat{\nabla} \hat{p}_{i1} + \frac{5 k_B \bar{T}_i}{2 Z_i e \bar{B} \bar{L} V_A} \hat{\nabla} \hat{T}_{i1} \right] \\
&- \frac{2}{3} \hat{T}_{i0} \hat{B}_0 \hat{\nabla}_{\parallel 0} \left(\frac{\hat{V}_{\parallel i}}{\hat{B}_0} \right) \\
&- \frac{2}{3} \hat{T}_{i1} \left(\frac{2}{\hat{B}_0} \mathbf{b}_0 \times \hat{\kappa} \right) \cdot \left[\hat{\nabla} \hat{\Phi}_0 + \frac{k_B \bar{T}_i}{Z_i e \bar{B} \bar{L} V_A} \frac{1}{\hat{n}_{i0}} \hat{\nabla} \hat{P}_{i0} + \frac{5 k_B \bar{T}_i}{2 Z_i e \bar{B} \bar{L} V_A} \hat{\nabla} \hat{T}_{i0} \right] \\
&+ \frac{1}{\hat{n}_{i0}} \hat{\nabla}_{\parallel} \left(\hat{\kappa}_{\parallel i} \hat{\nabla}_{\parallel} \hat{T}_{i1} \right) + \frac{1}{\hat{n}_{i0}} \hat{\nabla}_{\perp} \left(\hat{\kappa}_{\perp i} \hat{\nabla}_{\perp} \hat{T}_{i1} \right) \\
&+ \frac{2 Z_i m_e \bar{t}}{m_i \tau_e} \left(\hat{T}_{e1} - \hat{T}_{i1} \right) \\
&- \frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla}_{\perp} \hat{\phi} \cdot \hat{\nabla} \hat{T}_{i1} - \hat{V}_{\parallel i} \hat{\nabla}_{\parallel 0} \hat{T}_{i1} + \hat{V}_{\parallel i} \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \hat{T}_{i0} \\
&- \frac{2}{3} \hat{T}_{i1} \left(\frac{2}{\hat{B}_0} \mathbf{b}_0 \times \hat{\kappa} \right) \cdot \left[\hat{\nabla} \hat{\phi} + \frac{k_B \bar{T}_i}{Z_i e \bar{B} \bar{L} V_A} \frac{1}{\hat{n}_{i0}} \hat{\nabla} \hat{p}_{i1} + \frac{5 k_B \bar{T}_i}{2 Z_i e \bar{B} \bar{L} V_A} \hat{\nabla} \hat{T}_{i1} \right] \\
&- \frac{2}{3} \hat{T}_{i1} \hat{B}_0 \hat{\nabla}_{\parallel 0} \left(\frac{\hat{V}_{\parallel i}}{\hat{B}_0} \right) + \frac{2}{3} \hat{T}_{i0} \hat{B}_0 \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \left(\frac{\hat{V}_{\parallel i}}{\hat{B}_0} \right),
\end{aligned}$$

$$T_{e\text{para}1} = \frac{2}{3} \frac{1}{Z_i \bar{L} V_A}, \quad T_{e\text{para}2} = \frac{k_B \bar{T}_e}{e \bar{B} \bar{L} V_A}, \quad T_{e\text{para}3} = \frac{\bar{B}}{e \mu_0 \bar{n}_e \bar{L} V_A}, \quad T_{e\text{para}4} = \frac{\bar{B}^2}{\mu_0 k_B \bar{n}_e \bar{T}_e}, \quad (4.9)$$

$$\begin{aligned} \frac{\partial}{\partial t} \hat{T}_e &= -\frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla}_\perp \hat{\phi} \cdot \hat{\nabla} \hat{T}_{e0} - \frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla}_\perp \hat{\Phi}_0 \cdot \hat{\nabla} \hat{T}_{e1} - \hat{V}_{\parallel e} \hat{\nabla}_{\parallel 0} \hat{T}_{e0} \\ &\quad - \frac{2}{3} \hat{T}_{e0} \left(\frac{2}{\hat{B}_0} \mathbf{b}_0 \times \hat{\kappa} \right) \cdot \left[\hat{\nabla} \hat{\phi} - \frac{k_B \bar{T}_e}{e \bar{B} \bar{L} V_A} \frac{1}{\hat{n}_{e0}} \hat{\nabla} \hat{p}_{e1} - \frac{5 k_B \bar{T}_e}{2 e \bar{B} \bar{L} V_A} \hat{\nabla} \hat{T}_{e1} \right] \\ &\quad - \frac{2}{3} \hat{T}_{e0} \hat{B}_0 \hat{\nabla}_{\parallel 0} \left(\frac{\hat{V}_{\parallel e}}{\hat{B}_0} \right) \\ &\quad - \frac{2}{3} \hat{T}_{e1} \left(\frac{2}{\hat{B}_0} \mathbf{b}_0 \times \hat{\kappa} \right) \cdot \left[\hat{\nabla} \hat{\Phi}_0 - \frac{k_B \bar{T}_e}{e \bar{B} \bar{L} V_A} \frac{1}{\hat{n}_{e0}} \hat{\nabla} \hat{p}_{e0} - \frac{5 k_B \bar{T}_e}{2 e \bar{B} \bar{L} V_A} \hat{\nabla} \hat{T}_{e0} \right] \\ &\quad + 0.71 \frac{2 \bar{B}}{3 e \mu_0 \bar{n}_e \bar{L} V_A} \frac{\hat{T}_{e0}}{\hat{n}_{e0}} \hat{B}_0 \hat{\nabla}_{\parallel 0} \left(\frac{\hat{J}_{\parallel 1}}{\hat{B}_0} \right) + 0.71 \frac{2 \bar{B}}{3 e \mu_0 \bar{n}_e \bar{L} V_A} \frac{\hat{T}_{e1}}{\hat{n}_{e0}} \hat{B}_0 \hat{\nabla}_{\parallel 0} \left(\frac{\hat{J}_{\parallel 0}}{\hat{B}_0} \right) \\ &\quad - 0.71 \frac{2 \bar{B}}{3 e \mu_0 \bar{n}_e \bar{L} V_A} \frac{\hat{T}_{e0}}{\hat{n}_{e0}} \hat{B}_0 \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla}_{\parallel 0} \left(\frac{\hat{J}_{\parallel 0}}{\hat{B}_0} \right) \\ &\quad + \frac{1}{\hat{n}_{e0}} \hat{\nabla}_{\parallel} \left(\hat{\kappa}_{\parallel e} \hat{\nabla}_{\parallel} \hat{T}_{e1} \right) + \frac{1}{\hat{n}_{e0}} \hat{\nabla}_\perp \left(\hat{\kappa}_{\perp e} \hat{\nabla}_\perp \hat{T}_{e1} \right) \\ &\quad - \frac{2 m_e \bar{t}}{m_i \tau_e} \left(\hat{T}_{e1} - \hat{T}_{i1} \right) + \frac{4 \bar{B}^2}{3 \mu_0 \bar{n} k_B \bar{T}_e} \frac{\hat{\eta}_{\parallel}}{\hat{n}_{e0}} \hat{J}_{\parallel 0} \hat{J}_{\parallel 1} \\ &\quad - \frac{1}{\hat{B}_0} \mathbf{b}_0 \times \hat{\nabla}_\perp \hat{\phi} \cdot \hat{\nabla} \hat{T}_{e1} - \hat{V}_{\parallel e} \hat{\nabla}_{\parallel 0} \hat{T}_{e1} + \hat{V}_{\parallel e} \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \hat{T}_{e0} \\ &\quad - \frac{2}{3} \hat{T}_{e1} \left(\frac{2}{\hat{B}_0} \mathbf{b}_0 \times \hat{\kappa} \right) \cdot \left[\hat{\nabla} \hat{\phi} - \frac{k_B \bar{T}_e}{e \bar{B} \bar{L} V_A} \frac{1}{\hat{n}_{e0}} \hat{\nabla} \hat{p}_{e1} - \frac{5 k_B \bar{T}_e}{2 e \bar{B} \bar{L} V_A} \hat{\nabla} \hat{T}_{e1} \right] \\ &\quad - \frac{2}{3} \hat{T}_{e1} \hat{B}_0 \hat{\nabla}_{\parallel 0} \left(\frac{\hat{V}_{\parallel e}}{\hat{B}_0} \right) + \frac{2}{3} \hat{T}_{e0} \hat{B}_0 \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \left(\frac{\hat{V}_{\parallel e}}{\hat{B}_0} \right) \\ &\quad + 0.71 \frac{2 \bar{B}}{3 e \mu_0 \bar{n}_e \bar{L} V_A} \frac{\hat{T}_{e1}}{\hat{n}_{e0}} \hat{B}_0 \hat{\nabla}_{\parallel 0} \left(\frac{\hat{J}_{\parallel 1}}{\hat{B}_0} \right) \\ &\quad - 0.71 \frac{2 \bar{B}}{3 e \mu_0 \bar{n}_e \bar{L} V_A} \frac{\hat{T}_{e0}}{\hat{n}_{e0}} \hat{B}_0 \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \left(\frac{\hat{J}_{\parallel 1}}{\hat{B}_0} \right) - 0.71 \frac{2 \bar{B}}{3 e \mu_0 \bar{n}_e \bar{L} V_A} \frac{\hat{T}_{e1}}{\hat{n}_{e0}} \hat{B}_0 \mathbf{b}_0 \times \hat{\nabla} \hat{\psi} \cdot \hat{\nabla} \left(\frac{\hat{J}_{\parallel 0}}{\hat{B}_0} \right) \\ &\quad + \frac{2 \bar{B}^2}{3 \mu_0 \bar{n} k_B \bar{T}_e} \frac{\hat{\eta}_{\parallel}}{\hat{n}_{e0}} \hat{J}_{\parallel 1}^2. \end{aligned}$$

5 Results

6 Conclusion

The equations of 6-field are listed below when we drop the polarization velocity and viscous terms

$$\begin{aligned}
\frac{\partial}{\partial t} \varpi &= - \left(\frac{1}{B_0} \mathbf{b} \times \nabla_{\perp} \Phi + V_{\parallel i} \mathbf{b} \right) \cdot \nabla \varpi \\
&\quad + B^2 \nabla_{\parallel} \left(\frac{J_{\parallel}}{B} \right) + 2 \mathbf{b} \times \boldsymbol{\kappa} \cdot \nabla p_1 \\
&\quad - \frac{1}{2\Omega_i} \left[\frac{1}{B} \mathbf{b} \times \nabla P_i \cdot \nabla (\nabla_{\perp}^2 \Phi) - Z_i e B \mathbf{b} \times \nabla n_i \cdot \nabla \left(\frac{\nabla_{\perp} \Phi}{B} \right)^2 \right] \\
&\quad + \frac{1}{2\Omega_i} \left[\frac{1}{B} \mathbf{b} \times \nabla \Phi \cdot \nabla (\nabla_{\perp}^2 P_i) - \nabla_{\perp}^2 \left(\frac{1}{B} \mathbf{b} \times \nabla \Phi \cdot \nabla P_i \right) \right], \tag{6.1}
\end{aligned}$$

$$\begin{aligned}
\frac{\partial}{\partial t} n_i &= - \left(\frac{1}{B_0} \mathbf{b} \times \nabla_{\perp} \Phi + V_{\parallel i} \mathbf{b} \right) \cdot \nabla n_i \\
&\quad - \frac{2n_i}{B} \mathbf{b} \times \boldsymbol{\kappa} \cdot \nabla \Phi - \frac{2}{Z_i e B} \mathbf{b} \times \boldsymbol{\kappa} \cdot \nabla P_i - n_i B \nabla_{\parallel} \left(\frac{V_{\parallel i}}{B} \right), \tag{6.2}
\end{aligned}$$

$$\frac{\partial}{\partial t} V_{\parallel i} = - \left(\frac{1}{B} \mathbf{b} \times \nabla_{\perp} \Phi + V_{\parallel i} \mathbf{b} \right) \cdot \nabla V_{\parallel i} - \frac{1}{m_i n_i} \mathbf{b} \cdot \nabla P, \tag{6.3}$$

$$\frac{\partial}{\partial t} \psi = - \frac{1}{B} \nabla_{\parallel} \phi + \frac{\eta}{\mu_0} \nabla_{\perp}^2 \psi + \frac{1}{e n_e B} \nabla_{\parallel} P_e + \frac{0.71}{e B} \nabla_{\parallel} T_e, \tag{6.4}$$

$$\begin{aligned}
\frac{\partial}{\partial t} T_i &= - \left(\frac{1}{B} \mathbf{b} \times \nabla_{\perp} \Phi + V_{\parallel i} \mathbf{b} \right) \cdot \nabla T_i \\
&\quad - \frac{2}{3} T_i \left[\left(\frac{2}{B} \mathbf{b} \times \boldsymbol{\kappa} \right) \cdot \left(\nabla \Phi + \frac{1}{Z_i e n_i} \nabla P_i + \frac{5}{2} \frac{1}{Z_i e} \nabla T_i \right) + B \nabla_{\parallel} \left(\frac{V_{\parallel i}}{B} \right) \right] \\
&\quad + \frac{2}{3n_i} \nabla_{\parallel} (\kappa_{\parallel i} \nabla_{\parallel} T_i) + \frac{2}{3n_i} \nabla_{\perp} (\kappa_{\perp i} \nabla_{\perp} T_i) \\
&\quad + \frac{2m_e}{m_i} \frac{Z_i}{\tau_e} (T_e - T_i), \tag{6.5}
\end{aligned}$$

$$\begin{aligned}
\frac{\partial}{\partial t} T_e &= - \left(\frac{1}{B} \mathbf{b} \times \nabla_{\perp} \Phi + V_{\parallel e} \mathbf{b} \right) \cdot \nabla T_e \\
&\quad - \frac{2}{3} T_e \left[\left(\frac{2}{B} \mathbf{b} \times \boldsymbol{\kappa} \right) \cdot \left(\nabla \Phi - \frac{1}{e n_e} \nabla P_e - \frac{5}{2} \frac{1}{e} \nabla T_e \right) + B \nabla_{\parallel} \left(\frac{V_{\parallel e}}{B} \right) \right] \\
&\quad + 0.71 \frac{2T_e}{3e n_e} B \nabla_{\parallel} \left(\frac{J_{\parallel}}{B} \right) \\
&\quad + \frac{2}{3n_e} \nabla_{\parallel} (\kappa_{\parallel e} \nabla_{\parallel} T_e) + \frac{2}{3n_e} \nabla_{\perp} (\kappa_{\perp e} \nabla_{\perp} T_e) \\
&\quad - \frac{2m_e}{m_i} \frac{1}{\tau_e} (T_e - T_i) + \frac{2}{3n_e} \eta_{\parallel} J_{\parallel}^2 \tag{6.6}
\end{aligned}$$

The definitions make the closure of the equations

$$\begin{aligned}
Z_i n_{i0} + Z_{im} n_{im} &= n_{e0} \\
n_j &= n_{j0} + n_{j1}, \\
P_j &= P_{j0} + p_{j1}, \\
P = P_i + P_e &= P_0 + p_1 = (P_{i0} + P_{e0}) + (p_{i1} + p_{e1}), \\
P_t &= P_i + P_e + P_{im}, \\
\Phi &= \Phi_0 + \phi, \\
J_{\parallel} &= J_{\parallel 0} + J_{\parallel 1}, \\
V_{\parallel e} &= \frac{Z_i n_i + Z_{im} n_{im}}{n_e} V_{\parallel i} - \frac{J_{\parallel 1}}{en_e} = V_{\parallel i} - \frac{J_{\parallel 1}}{en_e}, \\
\mathbf{b} &= \mathbf{b}_0 - \mathbf{b}_0 \times \nabla \psi, \\
\varpi &= n_{i0} \frac{m_i}{B_0} \left(\nabla_{\perp}^2 \phi + \frac{1}{n_{i0}} \nabla_{\perp} \phi \cdot \nabla_{\perp} n_{i0} + \frac{1}{n_{i0} Z_{ie}} \nabla_{\perp}^2 p_{i1} \right) \\
J_{\parallel 1} &= -\frac{1}{\mu_0} B_0 \nabla_{\perp}^2 \psi.
\end{aligned}$$

$$\begin{aligned}
\frac{\partial}{\partial t} \varpi &= -\frac{1}{B_0} \mathbf{b} \times \nabla_{\perp} \Phi \cdot \nabla \varpi + B^2 \nabla_{\parallel} \left(\frac{J_{\parallel}}{B} \right) + 2\mathbf{b} \times \boldsymbol{\kappa} \cdot \nabla p_1 \\
&\quad - \frac{1}{2\Omega_i} \left[\frac{1}{B} \mathbf{b} \times \nabla P_i \cdot \nabla (\nabla_{\perp}^2 \Phi) - Z_i e B \mathbf{b} \times \nabla n_i \cdot \nabla \left(\frac{\nabla_{\perp} \Phi}{B} \right)^2 \right] \\
&\quad + \frac{1}{2\Omega_i} \left[\frac{1}{B} \mathbf{b} \times \nabla \Phi \cdot \nabla (\nabla_{\perp}^2 P_i) - \nabla_{\perp}^2 \left(\frac{1}{B} \mathbf{b} \times \nabla \Phi \cdot \nabla P_i \right) \right] + \mu_{\parallel i} \nabla_{\parallel 0}^2 \varpi,
\end{aligned} \tag{6.7}$$

$$\frac{\partial}{\partial t} n_i = -\frac{1}{B_0} \mathbf{b} \times \nabla_{\perp} \Phi \cdot \nabla n_i - \frac{2n_i}{B} \mathbf{b} \times \boldsymbol{\kappa} \cdot \nabla \Phi - \frac{2}{Z_i e B} \mathbf{b} \times \boldsymbol{\kappa} \cdot \nabla P_i - n_i B \nabla_{\parallel} \left(\frac{V_{\parallel i}}{B} \right), \tag{6.8}$$

$$\frac{\partial}{\partial t} V_{\parallel i} = -\frac{1}{B} \mathbf{b} \times \nabla_{\perp} \Phi \cdot \nabla V_{\parallel i} - \frac{1}{m_i n_i} \mathbf{b} \cdot \nabla P, \tag{6.9}$$

$$\frac{\partial}{\partial t} \psi = -\frac{1}{B} \nabla_{\parallel} \phi + \frac{\eta}{\mu_0} \nabla_{\perp}^2 \psi + \frac{1}{e n_e B} \nabla_{\parallel} P_e + \frac{0.71}{e B} \nabla_{\parallel} T_e, \tag{6.10}$$

$$\begin{aligned}
\frac{\partial}{\partial t} T_i &= -\frac{1}{B} \mathbf{b} \times \nabla_{\perp} \Phi \cdot \nabla T_i \\
&\quad - \frac{2}{3} T_i \left[\left(\frac{2}{B} \mathbf{b} \times \boldsymbol{\kappa} \right) \cdot \left(\nabla \Phi + \frac{1}{Z_i e n_i} \nabla P_i + \frac{5}{2} \frac{1}{Z_i e} \nabla T_i \right) + B \nabla_{\parallel} \left(\frac{V_{\parallel i}}{B} \right) \right] \\
&\quad - \frac{2}{3 n_i} \nabla_{\parallel 0} q_{\parallel i} + \frac{2m_e}{m_i} \frac{Z_i}{\tau_e} (T_e - T_i),
\end{aligned} \tag{6.11}$$

$$\begin{aligned}
\frac{\partial}{\partial t} T_e &= -\frac{1}{B} \mathbf{b} \times \nabla_{\perp} \Phi \cdot \nabla T_e \\
&\quad - \frac{2}{3} T_e \left[\left(\frac{2}{B} \mathbf{b} \times \boldsymbol{\kappa} \right) \cdot \left(\nabla \Phi - \frac{1}{e n_e} \nabla P_e - \frac{5}{2} \frac{1}{e} \nabla T_e \right) + B \nabla_{\parallel} \left(\frac{V_{\parallel e}}{B} \right) \right] \\
&\quad + 0.71 \frac{2T_e}{3e n_e} B \nabla_{\parallel} \left(\frac{J_{\parallel}}{B} \right) - \frac{2}{3 n_e} \nabla_{\parallel 0} q_{\parallel e} - \frac{2m_e}{m_i} \frac{1}{\tau_e} (T_e - T_i) + \frac{2}{3 n_e} \eta_{\parallel} J_{\parallel}^2
\end{aligned} \tag{6.12}$$

BOUT++ two fluid 6 field model($\varpi, n_i, V_{\parallel i}, T_i, T_e, \psi$): based on Braginskii Equations, the density, momentum and energy of ion and electron are described in drift ordering.

Continuity Terms
Compressional Terms
Electron Hall
Gyroviscous Terms
Thermal Force
Energy Exchange
Energy Flux
Thermal Conduction

$$\nabla_{\parallel} q_{\parallel i, e} = \nabla_{\parallel 0} q_{\parallel i, e} - \mathbf{b}_0 \times \nabla \psi \cdot \nabla q_{\parallel i, e} \tag{6.13}$$

References

- [1] A. N. Simakov and P. J. Catto, Phys. Plasmas **10**, 4744 (2003). <http://dx.doi.org/10.1063/1.1703527>
- [2] X. Q. Xu, W. M. Nevins, R. H. Cohen, J. R. Myra, P. B. Snyder, New J. Phys. **4**, 53.1 (2002).
- [3] X. Q. Xu et al., Commun. Comput. Phys. **4**, 949 (2008).
- [4] T. Y. Xia, X. Q. Xu, Nucl. Fusion **55**, 113030 (2015). <http://iopscience.iop.org/0029-5515/55/11/113030>