

Optimization Techniques in Finance

Homework assignment #4

Andrew Lesniewski

Baruch College

New York

December 3, 2018

Problems

1. Consider the following *analytic centering problem* with equality constraints:

$$\min - \sum_{i=1}^n \log x_i, \quad \text{subject to } Ax = b. \quad (1)$$

- (i) Show that in the simplest case of a single constraint $\sum_{i=1}^n x_i = 1$, the solution to this problem is $x_i^* = 1/n$, $\lambda^* = n$.
- (ii) Show that the dual Lagrange function is

$$q(\lambda) = -b^T \lambda + \sum_{i=1}^n \log(A^T \lambda)_i + n. \quad (2)$$

Formulate the dual problem.

- (iii) Show that the solution to the first order conditions is

$$x_i = \frac{1}{(A^T \lambda)_i}.$$

Therefore, to solve the equality constraint analytic centering problem, we solve the unconstrained dual problem for λ^* and then obtain x^* from these formulas.

- (iv) Implement Newton's method in Python in order to solve the optimization problem (1). Assume $\alpha = 0.1$ and $\beta = 0.5$ in the backtracking line search algorithm, and make sure that you take the structure of the problem into account. Carry out the convergence analysis of the algorithm for the problem with $p = 100$ and $n = 500$ (choose your own $p \times n$ constraint matrix!) by following the trajectories of Newton iterates (until they reach the exit criterion) starting at five different initial points.
- (v) Carry out a similar analysis for solving the corresponding dual problem.

2. Consider a KKT matrix

$$\begin{pmatrix} H & A^T \\ A & 0 \end{pmatrix}, \quad (3)$$

where $H \in \text{Mat}_n(\mathbb{R})$ is positive semidefinite, $A \in \text{Mat}_{pn}(\mathbb{R})$ is of maximum rank $\text{rank}(A) = p < n$.

- (a) Show that each of the following statements is equivalent to nonsingularity of the KKT matrix:
 - (i) $\mathcal{N}(H) \cap \mathcal{N}(A) = \{0\}$, where $\mathcal{N}(P)$ denotes the null-space of the matrix P .
 - (ii) $Ax = 0$, $x \neq 0$ imply that $x^T H x > 0$.
 - (iii) $F^T H F > 0$, where $F \in \text{Mat}_{n, n-p}(\mathbb{R})$ is a matrix for which $\mathcal{R}(F) = \mathcal{N}(A)$. Here $\mathcal{R}(F)$ denotes the range of F .
 - (iv) $H + A^T Q A > 0$, for some $Q > 0$.
- (b) Show that if the KKT matrix is nonsingular, then it has exactly n positive and p negative eigenvalues.

This assignment is due on December 17