

© COPYRIGHT NOTICE CELS191 LECTURE MATERIAL

The content and delivery of all resources in this course are subject to copyright and may be used only for the University's educational purposes.

Lecture handouts and slides contain material used under copyright licenses for the purposes of teaching.

You may access this material for your private study or research but may not make further copies or distribute these materials for any purpose, or in any other form, whether with or without charge.





Professor Jon Waters Zoology & Genetics jon.waters@otago.ac.nz

Research & Interests:

Profile: https://www.otago.ac.nz/zoology/staff/otago008938.html



CELS191 2025

Molecular Biology & Genetics

Lecture 20

X-linked Traits & Recombination

Professor Jon Waters
Te Tari Mātai Kararehe | Department of Zoology

Lecture 20 Objectives

After you have revised this lecture you should be able to:

- Identify the inheritance patterns of genes on sex-linked chromosomes.
- Explain that linked genes show non-independent assortment.
- Explain that crossing over leads to recombination of linked genes.


SEX-LINKED TRAITS

- Involving genes on the sex chromosomes (X or Y) (rather than autosomes)
- As a result, their inheritance patterns differ between males (XY) and females (XX)
- Examples: haemophilia; red-green colour blindness


SEX DETERMINATION

Sex chromosomes x,y determine the sex of the individual.

Other chromosomes are called autosomes.



In bees, fertilised eggs develop into diploid females; unfertilised eggs develop into haploid males







HETEROGAMETIC

HOMOGAMETIC


BIRDS







Male birds carry two identical sex chromosomes (ZZ) and females have two different ones (ZW)

Drosophila








Drosophila females have two X chromosomes and may also have a Y chromosome; males have an X chromosome and, if they are fertile, a Y chromosome





Eye Colour is a sex-linked trait in Drosophila








Eye Colour is a sex-linked trait in Drosophila


















The majority of those with red-green deficiencies should read the number 5




Red-Green colour blindness is a sex-linked trait










A father with the trait will transmit the mutant allele to all daughters but to no sons. When the mother is a dominant homozygote, the daughters will have the normal phenotype but will be carriers of the mutation.








A carrier who mates with a normal male will pass the mutation to half her sons and half her daughters.
The sons with the mutation will have the disorder. The daughters who have the mutation in a single dose will be carriers like their mother.







A carrier mates with a male with the trait, there is a 50% chance that each child will have the trait. Daughters who do not have the trait will be carriers, whereas males without the trait will be completely free of the harmful recessive allele.

Gene Mapping

- Q. How many genes are there in humans?
- **A.** Around 20,000
- Q. How many chromosomes are there in humans?
- **A.** 23 pairs


That's about 1,000 linked genes per chromosome

- Q. How are these linked genes inherited?
- A. Linked genes show biases towards parental genotypes
- Q. How can we determine the order and location of genes within a chromosome?
- A. By using recombination frequencies

A quick lesson on the convention of how fruit fly (Drosophila) genes are symbolized


- 1. The first mutant (non-wild type) discovered provides the letter(s). e.g. 'vg' for vestigial wings
- 2. The wild type allele for that gene is given a * after the symbol. e.g. 'vg*' is wild type wings
- 3. vg is recessive to vg⁺






What ratio of phenotypes would we expect?





Expected phenotype ratio 1:1:1:1






A homozygote recessive. Looks like a test cross to me!!

These are the results expected from Mendel's 2nd law (independent assortment of alleles)


CONCLUSION

- These two genes do not sort independently
- They are linked on the same chromosome

Why don't all gametes have parental genotypes?








We do get <u>some</u> recombinants.....why?????



How does recombination occur for linked genes?



Two chromatids of a tetrad


(one from each pair)cross over at random
points and swap genetic material.

This is a simple cross where
only one chiasma has formed.

The proportion of recombinant gametes is termed the recombination frequency








In this case, 17% of the gametes contain a chromosome that has formed a chiasma between the two genes

Recombination Frequency = (206+185) / (944+965+206+185) = 0.17 or 17%




The smaller the distance between two genes - the less likely a chiasma will be formed between them




- Distant (unlinked) genes have recombination frequencies around 50% (half recombinant and half parental types).
- Close genes have recombination frequencies of 0-50%.
- ❖ There is a near linear relationship between distance and recombination frequency.







Step Towards a Genetic Map



Sturtevant was able to derive a partial map of Drosophila by using recombinant frequencies of five traits. He used an arbitrary unit of distance the: centimorgan (cM) = 100 xrecombination frequency

Example Question

A species of flowering plant has a gene (purple) that determines flower colour, and another gene (tall) that controls plant height. A scientist performs a test cross between two lines of this species: one heterozygous plant line with purple flowers and tall stems; and the other line pure-bred with white flowers and short stems. The progeny from this cross comprise: 345 purple-flowered tall plants; 340 white-flowered short plants; 255 whiteflowered tall plants; and 260 purple-flowered short plants.

1. Give the genotype for each phenotype using standard genetic symbols

Pptt PpTt X pptt → PpTt pptt ppTt purple tall white short white tall purple short purple tall white short 255 345 260 340

2. Calculate the recombination frequency of these two genes

(255 + 260)/(345 + 340 + 255 + 260) = 515/1200 = 0.429 (42.9%)

3. Why is there a bias towards parental phenotypes?

Fewer recombinants means the 2 genes must be linked (on same chromosome)(violates Mendel's 2nd Law (independent assortment))

Lecture 20 Summary

- Sex-linked genes show inheritance patterns that differ between sexes.
- X-linked recessive traits such as red-green colour blindness are more common in males than in females.
- Crossing over during meiosis can shuffle genetic material between the non-sister chromatids of two homologous chromosomes.
- The distance between 2 linked genes can be estimated from the proportion of new phenotypes/genotype combinations produced by a test cross

Objective-Based Questions

- Define the terms carrier, homozygous, recombinant and hemizygous.
- Eye colour is a sex-linked trait in Drosophila i.e., the allele for eye colour is on the X-chromosome.
 - a) If you cross a homozygous red eyed female with a hemizygous white eyed male, what are the sex, genotypes and phenotypes of the offspring?
 - b) If you cross a homozygous white eyed female with a hemizygous red eyed male, what are the sex, genotypes and phenotypes of the offspring?
- What does it mean when we talk about linked genes? How are linked genes inherited? With linked genes, is the bias towards PARENTAL or RECOMBINANT genotypes?
- Fill in the gaps:

The GRE	EATER	R/LESS	SER the	distance between	two genes, the	e more likely	a chiasma v	will be
formed	betv	veen	them.	Distant genes have	HIGHER/LOW	ER recombina	ation freque	ncies.
There	is	а	near		relationship	between	distance	and



CELS191 2025

Molecular Biology & Genetics

Lecture 20

X-linked Traits & Recombination

Professor Jon Waters
Te Tari Mātai Kararehe | Department of Zoology