


# HUBS191 Lecture Material

This pre-lecture material is to help you prepare for the lecture and to assist your note-taking within the lecture, it is NOT a substitute for the lecture !




Please note that although every effort is made to ensure this pre-lecture material corresponds to the live-lecture there may be differences / additions.

# ***HUBS 191***

## ***Lecture 32***

### **Innate Immunity II**

**A/Prof Joanna Kirman**



© The content and delivery of all resources in this course are copyrighted. This includes video and audio recordings, PowerPoints, lecture notes and handouts. You may access the materials provided for your private study or research but may not further distribute the materials for any purpose, or in any other form, whether with or without charge.

# Objectives

2

## **To be able to:**

- Describe the key components of the inflammatory response
- Describe the 5 stages of phagocytosis
- Explain the main features of the complement system

## **Readings:**

- Marieb 10e Chapter 21 second part of Section 21.2, page 794 (Section Phagocytosis and Figure 21.2)
- Marieb 10e Chapter 21 page 795-797 (Section Inflammation: Tissue Response to Injury and Figure 21.3 and 21.4)
- Marieb 10e Chapter 21 page 798-799 (Section Complement and Figure 21.6)


# 3 layers of immune defense

3

1. Physical & Chemical Barriers (skin, mucosal membranes)
2. The Innate Immune System
3. The Adaptive Immune System


# The inflammatory response

4




# The inflammatory response

5




# The inflammatory response

6



# The inflammatory response


7





# The inflammatory response


8



Neutrophils squeeze through the 'leaky' capillary wall and follow the chemical trail to the injury site

# Many *myeloid* cells are phagocytic

9




In 1886-87 Elie Metchnikoff discovered the cells in the blood could ingest and destroy microbes – a process called phagocytosis

Phagein = Ancient Greek “to devour”


# Stages of phagocytosis

10



Marieb Figure 21.2

**(b) Events of phagocytosis.**



Neutrophils phagocytosing *Staphylococcus aureus* (green)  
Dr Richard Sequeira, University of Auckland

# Killing and digestion of phagocytosed microbes


12

- Low pH - acid environment
- Reactive oxygen (hydrogen peroxide) and reactive nitrogen intermediates (nitric oxide)
- Enzymes
  - Proteases
  - Lipases
  - Nucleases

# The Complement Cascade

13

- Complement – 9 major proteins/protein complexes (C1-9) act in sequence to clear pathogens from blood and tissues
  - ▣ Label pathogens (opsonisation)
  - ▣ Recruit phagocytes (chemotaxis)
  - ▣ Destroy pathogens (lysis)



# 3 Complement Pathways

14

## CLASSICAL

**Antibody bound  
to pathogen binds  
complement**

## ALTERNATIVE

**Pathogen binds  
complement to  
surface/pathogen  
component**


## LECTIN

**Carbohydrate  
components of  
microbes bind  
complement**



# Complement pathways converge

15





# 3 outcomes from the complement cascade


16

<b>LABEL</b> <b>Opsonisation</b> (labels pathogens which bind to complement receptors on phagocytes)	<b>DESTROY</b> <b>Membrane Attack</b> <b>Complex formation:</b> pores in bacterial cells → death	<b>RECRUIT</b> <b>Complement</b> proteins act as peptide mediators of inflammation and recruit phagocytes
<b>C3b</b>	<b>C9</b>	<b>C3a and C5a</b>

# Label (Opsonisation)

17


- Opsonisation = coating of a microbe with:
  - ▣ Antibody and/or
  - ▣ Complement fragment C3b



# Recruit

18

- Phagocytes attracted into site
- Mast cells degranulated by C3a and C5a




- Inflammatory mediators released including proteins that attract phagocytes

# Destroy

19

- microbes coated with C3b are phagocytosed
- assembly of the MAC causes lysis

MACs form from activated complement components (C5b and C6–C9) that insert into the target cell membrane, creating pores that can lyse the target cell.



# Revision MCQ

20

Killing of a phagocytosed microbe requires:

- a) binding and activation of toll-like receptors.
- b) degranulation caused by C3a and C5.
- c) anti-microbial peptides, such as defensins.
- d) reactive oxygen and nitrogen intermediates.

# HUBS191

## Copyright Warning Notice

This coursepack may be used only for the University's educational purposes. It includes extracts of copyright works copied under copyright licences. You may not copy or distribute any part of this coursepack to any other person.

Where this coursepack is provided to you in electronic format you may only print from it for your own use. You may not make a further copy for any other purpose. Failure to comply with the terms of this warning may expose you to legal action for copyright infringement and/or disciplinary action by the University

