

# Lecture 5

## Effect of Substituents on Orientation in $S_EAr$

Dr Andrea Vernall

[andrea.vernall@otago.ac.nz](mailto:andrea.vernall@otago.ac.nz)

## Reference textbook

- “Organic Chemistry” by Paula Yurkanis Bruice, 8th Edn.
- This lecture – Chapter 18, in particular 18.13, 18.14, 18.15

# Learning Objectives – Lecture 5

## *Section 2– Substituted benzenes and electrophilic aromatic substitution*

Describe and account for the differences in how an electron-donating (+I or +M) or weakly deactivating substituent affects the reactivity of different positions relative to it (ortho-, meta- and para-) in the aromatic ring towards electrophilic aromatic substitution.

Describe and account for the differences in how an electron-withdrawing (-I or -M) substituent affects the reactivity of different positions relative to it (ortho-, meta- and para-) in the aromatic ring towards electrophilic aromatic substitution

Appreciate that the overall directing and reactivity effects of some substituents (e.g. halogens) may be based on some combination of opposing inductive and mesomeric effects

Understand that steric effects also play a part in determining the relative reactivity of the different sites in an aromatic ring


Predict and account for the effects of common substituents on the chemical shifts of aromatic protons in  $^1\text{H}$  NMR spectra of substituted benzene derivatives

# Lecture Outline: L5

- 1. Introduction**
- 2. Electron Donating Substituents**
- 3. Electron Withdrawing Substituents**
- 4. Steric considerations**
- 5. Mixed examples**

# 1. Introduction

When substituted benzenes react with electrophiles where does the new substituent attach?



- All activating and weakly deactivating (the halogens) substituents direct  $E^+$  to the **ortho and para positions**.
- All other deactivating groups direct  $E^+$  to the **meta position**.

# 1. Introduction – Table from Lecture 4

(refer to Table 18.1 page 923)

## Strongly Activating towards $S_EAr$

|                |            |             |
|----------------|------------|-------------|
| $-\text{NH}_2$ | Ortho/Para | $+M \gg -I$ |
| $-\text{NHR}$  | Ortho/Para | $+M \gg -I$ |
| $-\text{NR}_2$ | Ortho/Para | $+M \gg -I$ |
| $-\text{OH}$   | Ortho/Para | $+M \gg -I$ |
| $-\text{OR}$   | Ortho/Para | $+M \gg -I$ |

## Activating towards $S_EAr$

|                         |            |           |
|-------------------------|------------|-----------|
| $-\text{NHCOR}$ (amide) | Ortho/Para | $+M > -I$ |
| $-\text{OCOR}$ (ester)  | Ortho/Para | $+M > -I$ |
| $-\text{Alkyl}$         | Ortho/Para | $+I$      |

# 1. Introduction – Table from Lecture 4

(refer to Table 18.1 page 923)

## Weakly Deactivating towards $S_EAr$

|                  |            |         |
|------------------|------------|---------|
| —F, —Cl, —Br, —I | Ortho/Para | —I > +M |
|------------------|------------|---------|

## Deactivating towards $S_EAr$

|                       |      |        |
|-----------------------|------|--------|
| —COH (aldehyde)       | Meta | —I, —M |
| —COR (ketone)         | Meta | —I, —M |
| —COOR (ester)         | Meta | —I, —M |
| —COOH (acid)          | Meta | —I, —M |
| —COCl (acid chloride) | Meta | —I, —M |


## Strongly Deactivating towards $S_EAr$

|                               |      |           |
|-------------------------------|------|-----------|
| —NO <sub>2</sub>              | Meta | —I, —M    |
| —CN                           | Meta | —I, —M    |
| —SO <sub>3</sub> H            | Meta | —I, —M    |
| —NH <sub>3</sub> <sup>+</sup> | Meta | strong —I |
| —NR <sub>3</sub> <sup>+</sup> | Meta | strong —I |

## 2. Electron donating substituents

- Recall from Lecture 4, +I and +M substituents activate the Ar ring to  $S_EAr$  and speed up the  $S_EAr$  reaction (relative to 'S' being H)
- But what relative ring position(s) is more activated than others?

Methylbenzene (toluene), the  $CH_3$  group is **inductively donating (+I)**




Consider *ortho* attack




## 2. Electron donating substituents

Methylbenzene (toluene), the CH<sub>3</sub> group is **inductively donating (+I)**

Consider *meta* attack



Consider *para* attack




## 2. Electron donating substituents

Methylbenzene (toluene), the CH<sub>3</sub> group is **inductively donating (+I)**

- Conclusion – S<sub>E</sub>Ar with a **+I substituent** present **directs to *ortho* and *para*** E<sup>+</sup> reaction as these have **more stable intermediates** than that of *meta*.
- The transition states leading to their formation are **lower in energy**.
- Thus **Δ E<sub>a</sub> is smaller**
- Hence the *ortho* and *para* products are formed faster.


**Now you can fully explain your lab 1 results !**



## 2. Electron donating substituents

Methoxybenzene is **mesomerically electron donating (+M)** (+M > -I)


Consider *ortho* attack



## 2. Electron donating substituents

Methoxybenzene is **mesomerically electron donating (+M)** (+M > -I)


Consider *meta* attack



## 2. Electron donating substituents

Methoxybenzene is **mesomerically electron donating (+M)** (+M > -I)


Consider *para* attack



## 2. Electron donating substituents

Methoxybenzene is **mesomerically electron donating (+M)** (+M > -I)


- Both ***ortho*** and ***para*** substitution have an 'extra' **very stable intermediate** while *meta* does not.
- Therefore +M substituents are *ortho*, *para* directing




## 2. Electron donating substituents

Methoxybenzene is **mesomerically electron donating (+M)** (+M > -I)

- Now with an **activating substituent** a **less reactive E<sup>+</sup>** can sometimes be used...
- Recall - to brominate benzene, bromine had to be turned into a better E<sup>+</sup>



- But bromination of methoxybenzene is fast and occurs without a catalyst ....




### 3. Electron withdrawing substituents

- Recall from Lecture 4, -I and -M substituents **deactivate** the Ar ring to  $S_E\text{Ar}$  and slow down the  $S_E\text{Ar}$  reaction (relative to 'S' being H)
- But what relative ring position(s) is more activated than others?

An anilinium ion (protonated aniline) is **strongly inductively withdrawing (-I)**

Consider *ortho* attack







### 3. Electron withdrawing substituents

An anilinium ion (protonated aniline) is **strongly inductively withdrawing (-I)**

Consider *meta* attack




Consider *para* attack



### 3. Electron withdrawing substituents

All **-M substituents** are also all **-I**

Consider *ortho* attack




- S<sub>E</sub>Ar para substitution results in **similar high energy contributors**.
- However, the meta pathway does not.
- **Thus -M substituents are meta directing.**

### 3. Electron withdrawing substituents

All **-M** substituents are also all **-I**

e.g. Nitration of nitrobenzene,  $\text{NO}_2$  group is deactivating and meta directing



The rate is  $10^{-4}$  **slower** than that of benzene. Forcing conditions are required.



**Much** more forcing nitration conditions will result in possibly tri-nitro products. But since each nitro group is deactivating the rates of these further substitution reactions is even slower.

*Does this also explain why **di- $\text{NO}_2$**  products were not formed in lab 1.....?*

### 3. Electron withdrawing substituents


All **-M** substituents are also all **-I**

- Friedel Crafts acylation and alkylation are the slowest of the  $S_EAr$ 's
- If a benzene ring has a moderate or strongly deactivating group a Friedel Crafts reaction will not even occur




### 3. Electron withdrawing substituents - halogens

- Halogens are an interesting case, although **electron withdrawing (-I)** and classified as **weakly deactivating** for  $S_EAr$  are o/p directing  
... why?
- **Weakly deactivating** just means  $S_EAr$  is comparatively slower to if an 'H' was the substituent instead.
- Halogens are the only 'deactivating' group that is NOT meta directing, instead o/p




## 4. Steric considerations

- Often with small sized substituents the products are the expected statistical ratio based on activation/deactivation of o, m, p positions




- But if the substituent is LARGE the para position is favoured, **steric factors**



## 5. Mixed examples

- What about a phenyl with 2 substituents?
- This may be additive or opposing in 'directing' and 'activating' aspects.




Methyl and bromo groups direct to their o and p positions

e.g. "ortho to bromo" "ortho to methyl", specify o, m, or p in relation to something...

## 5. Mixed examples

- If two groups direct to different positions the group with the stronger activating properties wins.



The **methoxy +M effect (strongly activating)** is much greater than the **alkyl +I effect (activating)**