



# HUBS191 Lecture Material

This pre-lecture material is to help you prepare for the lecture and to assist your note-taking within the lecture, it is NOT a substitute for the lecture !



Please note that although every effort is made to ensure this pre-lecture material corresponds to the live-lecture there may be differences / additions.



# ***HUBS191***

## ***Lecture 35***

### **T cells**

Prof. Alex McLellan,  
Dept. Microbiology & Immunology




UNIVERSITY  
of  
**OTAGO**  
Te Whare Wānanga o Otago  
NEW ZEALAND




# Objectives

- Know that DC take up antigen and stimulate T cells in lymph node or spleen.
- Understand how DC use MHC-I & MHC-II to stimulate CD8 and CD4 T cells.
- Know how T cells develop (born in bone marrow → TCR rearrangement in the thymus) and form effector and memory cells.
- Understand that CD4 helper cells help both CD8 T cells and B cells (next lecture) responses.
- Pre-reading: Marieb; p802-806, 811-819


# Infection and inflammation of tissue e.g. skin



# *Endogenous antigens*




The 'proteasome' degrades proteins to peptides.



# MHC-I antigen processing

- Antigenic proteins are degraded to peptides (in the cytoplasm) by the proteasome.
- Peptide loading of MHC-I takes place in the endoplasmic reticulum (ER).


# *Exogenous antigens*






# MHC-II antigen processing

- Antigenic proteins are degraded in acidic phagolysosome.
- Peptide loading of MHC-II takes place in phagolysosome.







# T cells

- Are lymphocytes that arise in the bone marrow and fully develop in thymus
- T cells express T cell receptor (TCR) with co-receptors (either CD4 or CD8).
- Recognise MHC / peptide complexes.

Production of T cell precursors



T cell development / TCR gene rearrangement




Thymus




lymphoid organs,  
blood & tissues.

# T cells develop in the thymus



# T cells express a unique T cell receptor

Immature T cells (TCR genes in germline state) in bone marrow




Mature (naïve) T cells expressing unique antigen receptors (TCR)

# Thymic gene rearrangement



- Immature T cells (thymocytes) rearrange the 'variable' parts of their TCR genes in the thymus
- The rearrangement process is essentially random.
- This ensures that individual T cells are unique in terms of their TCR. Creates 'diversity' in T cell repertoire.





T cells express a T cell receptor (TCR)  
that recognises peptide + MHC



# CD4 and CD8 'co-receptors' on T cells



# CD4 and CD8 expression on T cells



CD4 and CD8 molecules assist with the docking of the TCR onto MHC-II or MHC-I respectively

# T cell differentiation


- T cells that have not been activated by MHC / peptide are 'naïve'.
- Activated T cells are also known as 'effector T cells'.

# CD4 T helper cell


- Recognises MHC-II / peptide
- Helps CD8 T cell become cytotoxic
- Helps B cell make antibody

# CD8 T cell

- Recognise MHC-I / peptide
- Develops into 'cytotoxic T lymphocyte' (CTL) aka 'cytotoxic T cell'.




Cytokines produced by CD4 T cells help CD8 T cells become activated






Adaptive defenses  $\rightarrow$  Humoral immunity  
 $\rightarrow$  Cellular immunity

## (a) Helper T cells help in humoral immunity



## (b) Helper T cells help in cellular immunity



# Memory T cells

In addition to the formation of 'effector' cells, T cell activation results in the formation of memory T cells

Memory CD4 or CD8 T cells reside in the body for long periods of time.

Memory T cells become effector cells much quicker than naïve T cells.

# Summary

- Dendritic cells (DC) stimulate T cells using MHC-peptide. Occurs in lymph node and spleen
- T cells express unique T cell receptors (TCR) and develop into mature T cells in the thymus
- CD4 T cells are helper cells that respond to MHC-II-peptide.
- CD8 T cells respond to MHC-I-peptide and become cytotoxic T lymphocytes (CTL).
- T cells form memory cells.

# **The T cell receptor of CD4 T cells recognises:**

- A) The CD8 co-receptor on CD8 T cells.
- B) MHC-I / peptide complexes on antigen presenting cells.
- C) MHC-II / peptide complexes on antigen presenting cells.
- D) Native antigen on MHC-II

# The T cell receptor of CD8 T cells recognises:

- A) MHC-I / peptide complexes on antigen presenting cells (e.g. dendritic cells, macrophages, B cells).
- B) MHC-I / peptide complexes on virus-infected or cancer cells.
- C) MHC-II / peptide complexes on antigen presenting cells.
- D) Both (A) and (B) are correct.

# **The following is true for presentation of viral antigens:**

- A) Viral antigens are exclusively presented on MHC-I
- B) Viral antigens are mostly presented on MHC-I, but can also be presented on MHC-II by APC (e.g. DC)
- C) Viral antigens are mostly presented on MHC-II
- D) Viral antigens are mostly presented on MHC-II, but can be presented on MHC-I by DC

# **T cells express a diverse range of T cell receptors (TCR) due to:**

- A) Somatic hypermutation
- B) Expression of germline sequences from both chromosomes
- C) Expression of rearranged TCR genes
- D) Clonal selection
- E) Thymic selection

# **The site of T cell germline TCR gene rearrangement is the:**

- A) Thyroid gland
- B) Pancreas
- C) Thymus
- D) Spleen
- E) Bone marrow



**Immature T cell-precursors ('baby T cells') arise in the:**

- A) Thyroid gland
- B) Pancreas
- C) Thymus
- D) Spleen
- E) Bone marrow

## **Antigen presenting cells include:**

- A) neutrophils, mast cells, eosinophils & basophils
- B) T cells and B cells
- C) Natural killer cells, monocytes & macrophages
- D) B cells, macrophages, monocytes and dendritic cells

**Lymphocytes that have not seen antigen are termed:**

A) guileless

B) jejune

C) provincial

D) callow

E) naïve

# **Following stimulation via MHC-I, CD8 T cells become:**

- A) natural killer cells
- B) apoptotic cells
- C) plasma cells
- D) Cytotoxic T lymphocytes (aka cytotoxic T cells)

**Long-lived T cells (or B cells) that respond vigorously to a second encounter of antigen are:**


A) Recollection cells

B) Effector cells

C) Memory cells

D) Thymocytes

E) Medullary cells



# HUBS191

## **Copyright Warning Notice**

This coursepack may be used only for the University's educational purposes. It includes extracts of copyright works copied under copyright licences. You may not copy or distribute any part of this coursepack to any other person.

Where this coursepack is provided to you in electronic format you may only print from it for your own use. You may not make a further copy for any other purpose. Failure to comply with the terms of this warning may expose you to legal action for copyright infringement and/or disciplinary action by the University

