

**Module 3, Lecture 8**

**Structure and Reactions of Organic Molecules**

**Introduction to reaction mechanisms  
Substitution and Elimination**

**James Crowley**

[jcrowley@chemistry.otago.ac.nz](mailto:jcrowley@chemistry.otago.ac.nz)

**Dr. Bill Hawkins**

[bhawkins@chemistry.otago.ac.nz](mailto:bhawkins@chemistry.otago.ac.nz)

References to Brown *et al* text shown in BLUE

53


**Learning Objectives:**

- to be able to identify what factors are required for a good leaving group
- to have an understanding of elimination reactions
- to be able to identify what factors are required fro a good leaving group
- to have an understanding of E1 elimination mechanism

**Textbook:**     [\*\*Chapter 27, sections 27.5-27.7, Brown\*\*](#)

54

## What determines whether a reaction goes via S<sub>N</sub>1 or S<sub>N</sub>2?



Require **good** leaving group -departs in rate controlling step

### S<sub>N</sub>2:

- Nu<sup>-</sup> binds as leaving group departs
- Strong Nu<sup>-</sup> can make up for poor leaving group

### S<sub>N</sub>1:

- Only change occurring in slow step is cleavage of C-X bond
- Good leaving group **essential**.

55

## What determines whether a reaction goes via S<sub>N</sub>1 or S<sub>N</sub>2?

Relative rates of C-X cleavage

Longer,  
weaker  
bonds

X	Relative Rate
I <sup>-</sup>	150
Br <sup>-</sup>	50
Cl <sup>-</sup>	1
F <sup>-</sup>	2 x 10 <sup>-2</sup>
HO <sup>-</sup>	5 x 10 <sup>-8</sup>
RO <sup>-</sup>	5 x 10 <sup>-9</sup>

56


## What determines whether a reaction goes via S<sub>N</sub>1 or S<sub>N</sub>2?

### Relative rates of C-X cleavage


<i>X</i>	<i>Relative Rate</i>
I: <sup>-</sup>	150
Br: <sup>-</sup>	50
Cl: <sup>-</sup>	1
F: <sup>-</sup>	2 x 10 <sup>-2</sup>
HO: <sup>-</sup>	5 x 10 <sup>-8</sup>
RO: <sup>-</sup>	5 x 10 <sup>-9</sup>

57

HO:<sup>-</sup> and RO:<sup>-</sup> can be converted into better leaving groups under strongly acidic conditions (protonation).





In (X), electrons in C-O bond is drawn closer to +ve charge O.



58

### Example:



59

### Nucleophiles

#### $S_N1$ :

- $\text{Nu}^-$  not involved in slow step
- Carbocation reacts readily with any nucleophile

#### $S_N2$ :

- Good nucleophile essential

**Nucleophilicity:**  
measure of affinity  
towards electrophilic  $\text{C}^{\delta+}$

-ve

$\text{HO}^-$  good



$\text{H}_2\text{O}$  weak

neutral

60

## Nucleophiles

$(\text{CH}_3)_3\text{CO}^-$  strong base but poor nucleophile




61

## Effect of Solvent on $\text{S}_\text{N}$ Reactions

The biological medium for chemical reactions is generally  $\text{H}_2\text{O}$ .


$\text{H}_2\text{O}$  is a *polar* solvent. Also, able to form H-bonds.



62

## Effect of Solvent on $S_N$ Reactions

**$S_N1$ :** fastest in polar solvents, particularly those which H-bond.




First transition state stabilized by:

- polar effect at C
- H-bonding to L

63


## Effect of Solvent on $S_N$ Reactions

**$S_N2$ :** Solvent can influence  $Nu:^-$  through H-bonding



64


## Elimination Reactions (27.6)



- Common mechanisms, E1 and E2.
- Parallel those for S<sub>N</sub>1 and S<sub>N</sub>2.
- Elimination mechanism determined by base strength:
  - strong base - E2;
  - weak base - E1.

65

## E1 Mechanism




Experiments show:  $\text{rate} = k [\text{C-Cl}]$


66

## E1 Mechanism

1st Step (same as for S<sub>N</sub>1) (rate determining step)



2nd Step A base (H<sub>2</sub>O, most abundant) removes H<sup>+</sup> to form C=C:



67

## E1 Mechanism


Same rate determining step as S<sub>N</sub>1, thus same controlling factors:

- Strong base not required
- H-bonding solvents promote reaction
- E1 most common when leaving group is on 3° carbon
- E1 and S<sub>N</sub>1 often co-occur.

68




for previous example:



- Ratio of E1 : S<sub>N</sub>1 controlled by relative rates of the 2 alternative fast steps.
- If leaving group is changed, overall rate changes but E1 : S<sub>N</sub>1 ratio unchanged.

69

## E2 Mechanism




Experiments show:

$$\text{rate} = k [\text{CH}_3\text{CH}_2\text{Br}] [\text{HO}^-]$$

70

## E2 Mechanism



**Concerted** bond breaking and bond formation

- Halide **and** base involved in single, rate determining step
- Like  $\text{S}_{\text{N}}2$  Must have strong base - e.g.  $\text{HO}^-$
- Possible for  $1^\circ$ ,  $2^\circ$  and  $3^\circ$  compounds.

71

## E2 Mechanism

**$\text{S}_{\text{N}}2$ :**

- Requires backside attack;
- 
- Difficult if reaction centre crowded;
- Rate order is  $1^\circ > 2^\circ > 3^\circ$ .


**$\text{E}2$ :**

- Easily accessible  $\text{H}^+$  is removed
- Little difference in ease of  $\text{H}^+$  removal for  $1^\circ$ ,  $2^\circ$  and  $3^\circ$


72

## Questions

1. The rate of the following reaction is dependent on [alkyl bromide] and  $[\text{OH}^-]$ . Draw an energy profile of the reaction and the transition state structure(s).



2. Account for the following reactions:



73

## \* Homework \*

*Chemistry – the central science 3<sup>rd</sup> Ed*

Brown et al.

Problems 27.40, 27.77

Answers on Blackboard

74