

MA3614 - Complex Variable Methods and Applications

1720996

March 26, 2021

Contents

1	Week 1	3
1.1	Fundamentals	3
1.1.1	Representations of z and \bar{z}	3
1.1.2	Multiplication, powers and roots of unity	3
1.1.3	Triangle inequality in \mathbb{C}	3
1.1.4	Convergence of a sequence in \mathbb{C}	3
2	Week 2	4
2.1	Foundations of complex numbers	4
2.1.1	Roots of the unity polynomial	4
2.1.2	Some definitions	4
2.1.3	Limits	4
2.1.4	Continuity	5
3	Week 3	5
3.1	Functions and the Cauchy-Riemann equations	5
3.1.1	Analytic functions	5
3.1.2	Combining differentiable functions	5
3.1.3	The Cauchy-Riemann equations	6
4	Week 4	6
4.1	Analytic functions	6
4.1.1	Gradient	6
4.1.2	Directional derivative	6
4.1.3	Analytic function definition	6
5	Week 5	6
5.1	Analytic functions	6
5.1.1	Harmonic functions	6
5.1.2	Harmonic Conjugate	7
6	Week 6	7
6.1	Elementary functions of z	7
6.1.1	Representation of polynomials and zeros	7
6.1.2	Rational functions	7
6.1.3	Partial fractions representation	7
6.1.4	Residues	8
7	Week 7	8
7.1	Elementary functions of z	8
7.1.1	General case of the residues	8
7.1.2	Complex trig functions	8
7.1.3	The real and imaginary parts of $\sin(z)$ and $\cos(z)$	9

8	Week 8	9
8.0.1	Exponential function	9
8.0.2	cot and tanh	9
8.0.3	Log z and the multi-valued $\log z$	9
8.0.4	Complex powers z^α	9
9	Week 9	10
9.1	Integrals, arcs and contours	10
9.1.1	Series and the residue more generally	10
9.2	Cauchy theorems	10
9.2.1	Integral of a complex valued function	10
9.2.2	A smooth arc	10
9.2.3	A contour	11
9.2.4	The ML inequality	11
9.2.5	Independence of path when $f = F'$	11
10	Week 10	12
10.1	Loop integrals	12
10.1.1	Closed loops and powers of z	12
10.1.2	Path independence, loop integrals and anti-derivatives	12
10.1.3	Loop integrals and rational functions	12
11	Week 17	12
11.1	Harmonic functions - further results	12
11.1.1	Creating an analytic function from a harmonic function	13
12	Week 18	13
12.1	Functions defined by loop integrals	13
12.2	Loop integrals of $f(z)/q(z)$ where q =polynomial	13
12.3	Other versions of the formulae and entire functions	13
12.3.1	The fundamental theorem of algebra	14
12.3.2	Further Results	14
12.3.3	The mean value property	14
12.3.4	The maximum modulus theorem	14
12.4	Definitions: sequences in \mathbb{C}	14
12.4.1	Result about convergence	14
12.5	Definitions: series in \mathbb{C}	14
12.5.1	Results about series in \mathbb{C}	15
12.6	Series of functions	15
12.7	Uniform convergence and analytic functions	15
13	Week 19	15
13.1	Taylor series for analytic functions	15
13.1.1	Key formula in the proof of the Taylor series	15
13.1.2	Taylor's series, comments about R	16
13.2	Maclaurin series case	16
13.3	Real coefficients, even functions, odd functions, etc.	16
13.4	Series you are expected to know	16
14	Week 20	17
14.1	The Koebe function, de Branges' theorem and a conjecture	17
14.2	Multiplying series - the Cauchy product	17
14.3	Leibnitz's formula for the n th derivative of a product	17
14.4	The generalised L'Hopital's rule	17
14.5	Power series	18

15 Week 21	18
15.1 Cauchy-Hadamard theorem	18
15.2 Properties of a function defined by a power series	18
15.3 Laurent series	18
15.3.1 Classifying zeros and poles	19
16 Week 22	19
16.1 Integrating a Laurent series - the residue	19
16.2 The Residue theorem	19
16.2.1 Techniques to calculate the residue	19
17 Week 23	20
17.1 The integrals on C_R^+ when we have a $a^{imz}term$	20
17.2 Counting zeros and poles	20

1 Week 1

1.1 Fundamentals

1.1.1 Representations of z and \bar{z}

A complex number z can be defined in both cartesian or polar form:

$$z = x + iy = re^{i\theta}, \quad (1)$$

where $x = r \cos \theta$ and $y = r \sin \theta$. Here, r is the modulus and θ is the argument.

Definition 1 *The principal argument of z is*

$$\arg z \in (-\pi, \pi], \quad (2)$$

where $\arg z$ is multi-valued.

Note, $|z|^2 = z\bar{z}$. $|z|$ = absolute value of z .

1.1.2 Multiplication, powers and roots of unity

Suppose $z = re^{i\theta}$, $z_1 = r_1e^{i\theta_1}$, $r_2e^{i\theta_2}$.

- **Multiplication:** $z_1z_2 = r_1r_2e^{i(\theta_1+\theta_2)}$.
- **Powers:** $z^n = r^n e^{in\theta}$, $n = 0, \pm 1, \pm 2, \dots$
- Observe that $e^{2\pi i} = \exp(2\pi i) = 1$.
- **Roots of unity:** Let $\omega = \exp(2\pi i/n)$. $1, \omega, \omega^2, \dots, \omega^{n-1}$ all satisfy $z^n - 1 = 0$ and are uniformly spaced on the unit circle.

1.1.3 Triangle inequality in \mathbb{C}

Definition 2

$$||z_1| - |z_2|| \leq |z_1 + z_2| \leq |z_1| + |z_2| \quad (3)$$

1.1.4 Convergence of a sequence in \mathbb{C}

Definition 3 *A sequence z_0, z_1, z_2, \dots converges to z if for every $\epsilon > 0$ there exists an $N = N(\epsilon)$ such that*

$$|z_n - z| < \epsilon, \quad \forall n \geq N. \quad (4)$$

From here on, $||$ now means the absolute value of a complex number.

2 Week 2

2.1 Foundations of complex numbers

Theorem 1 A polynomial of degree n can always be factorised in the form

$$p_n(z) = a_n z^n + a_{n-1} z^{n-1} + \cdots + a_1 z + a_0 \quad (5)$$

$$= a_n (z - \alpha_1)(z - \alpha_2) \cdots (z - \alpha_n). \quad (6)$$

where $a_0, \dots, a_n, \alpha_1, \dots, \alpha_n \in \mathbb{C}$ and $a_n \neq 0$.

2.1.1 Roots of the unity polynomial

Let $\omega = \exp(2\pi i/n)$. Let $\xi = \rho \exp(i\alpha)$ and let $z_0 = \sqrt[n]{\rho} \exp(i\alpha/n)$ be one solution. The n roots of ξ are $z_0, z_0\omega, \dots, z_0\omega^{n-1}$.

2.1.2 Some definitions

Let $A \subset \mathbb{C}$. We write

$$f : A \rightarrow \mathbb{C}$$

with A denoting the domain of definition of f .

- **Open disk:** A set of the form

$$\{z \in \mathbb{C} : |z - z_0| < \rho\}, \quad \rho > 0. \quad (7)$$

The boundary is the unit circle $|z - z_0| = \rho$ which is *not* part of the set.

- **Unit disk:** This is the set

$$\{z \in \mathbb{C} : |z| < 1\}. \quad (8)$$

- **Neighbourhood:** A neighbourhood of a point z_0 means a disk of the form $\{z \in \mathbb{C} : |z - z_0| < \rho\}$ for some $\rho > 0$.
- **Interior point:** The interior point of A is a point $z_0 \in A$ such that a neighbourhood of z_0 is also in A .
- **Open set:** A set such that every point is an interior point.
- **Boundary point:** A boundary point of A is a point z_0 such that every neighbourhood of z_0 contains points which are in A and also contain points which are not in A .
- **Boundary:** The boundary of A is the set of all its boundary points.
- **Polygonal path:** Let w_1, w_2, \dots, w_{n+1} be points in \mathbb{C} and let l_k be the straight line segment joining w_k to w_{k+1} . The successive line segments l_1, l_2, \dots, l_{n+1} is a polygonal path joining w_1 to w_{n+1} .
- **Connected:** A set A is connected if every pair of points z_1 and z_2 in A can be joined by a polygonal path which is contained in A .
- **Domain:** An open connected set.
- **Region:** A domain or a domain together with some or all of the boundary points.
- **Bounded:** A set A is bounded if there exists $R > 0$ such that the set is contained in the disk $\{z : |z| < R\}$.
- **Unbounded:** A set is unbounded if it's not bounded.
- A domain (which is thus connected) and does not have holes.

2.1.3 Limits

Definition 4 Let f be defined in a neighbourhood of z_0 and let $f_0 \in \mathbb{C}$. If for every $\epsilon > 0$ there exists a real number $\delta > 0$ such that

$$|f(z)| < \epsilon \text{ for all } z \text{ satisfying } 0 < |z - z_0| < \delta,$$

then we say that

$$\lim_{z \rightarrow z_0} f(z) = f_0. \quad (9)$$

2.1.4 Continuity

Definition 5 A function $w = f(z)$ is continuous at $z = z_0$ provided $f(z_0)$ is defined and

$$\lim_{z \rightarrow z_0} f(z) = f(z_0). \quad (10)$$

Suppose that $f(z)$ and $g(z)$ are continuous at z_0 .

- $f(z) \pm g(z)$ and $f(z)g(z)$ are continuous at z_0 .
- $f(z)/g(z)$ is continuous at z_0 provided $g(z) \neq 0$.

Suppose that $f(z)$ is continuous at z_0 and $g(z)$ is continuous at $f(z_0)$ then $g(f(z))$ is continuous at z_0 . Let $f(z) = u(x, y) + iv(x, y)$. If f is continuous at $z_0 = x_0 + iy_0$ then u and v are both continuous as functions on \mathbb{R}^2 at (x_0, y_0) . Conversely, if u and v are both continuous at (x_0, y_0) then f is continuous at $z_0 = x_0 + iy_0$.

3 Week 3

3.1 Functions and the Cauchy-Riemann equations

3.1.1 Analytic functions

Theorem 2 Let f be a complex valued function defined in a neighbourhood of z_0 . The derivative of f at z_0 is given by

$$\frac{df}{dz}(z_0) \equiv f'(z_0) := \lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h} \quad (11)$$

provided the limit exists. Note that here $h \in \mathbb{C}$.

- A function f is analytic at z_0 if f is differentiable at all points in some neighbourhood of z_0 .
- A function f is analytic in a domain if f is analytic at all points in the domain.
- A function $f : \mathbb{C} \rightarrow \mathbb{C}$ is an entire function if it is analytic on the whole complex plane \mathbb{C} .

3.1.2 Combining differentiable functions

Let f and g be differentiable at z_0 . We have the following:

1.

$$(f \pm g)'(z_0) = f'(z_0) \pm g'(z_0).$$

2.

$$(cf)'(z_0) = cf'(z_0)$$

for all constants $c \in \mathbb{C}$.

3.

$$(fg)'(z_0) = f(z_0)g'(z_0) + f'(z_0)g(z_0). \quad (12)$$

This is the product rule.

4.

$$\left(\frac{f}{g}\right)'(z_0) = \frac{g(z_0)g'(z_0) - f(z_0)g'(z_0)}{g(z_0)^2}, \quad \text{if } g(z_0) \neq 0. \quad (13)$$

This is the quotient rule.

5. Let now f be a function which is differentiable at $g(z_0)$. Then

$$\left.\frac{d}{dz}f(g(z))\right|_{z=z_0} = f'(g(z_0))g'(z_0). \quad (14)$$

This is the chain rule.

3.1.3 The Cauchy-Riemann equations

Let $f(z) = u(x, y) + iv(x, y)$. When f is analytic at z_0 the following limit exists:

$$\frac{df}{dz}(z_0) \equiv f'(z_0) := \lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h}. \quad (15)$$

By considering the case when h is real and then purely imaginary we get

$$f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}, \quad (16)$$

$$= \frac{1}{i} \left(\frac{\partial u}{\partial y} + i \frac{\partial v}{\partial y} \right) = \frac{\partial v}{\partial y} - i \frac{\partial u}{\partial y}. \quad (17)$$

Equating the real and imaginary parts gives the Cauchy-Riemann equations.

Theorem 3 *The Cauchy-Riemann equations are*

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}. \quad (18)$$

The Cauchy-Riemann equations in polar coordinates are

$$\frac{\partial \tilde{u}}{\partial r} = \frac{1}{r} \frac{\partial \tilde{v}}{\partial \theta}, \quad \frac{1}{r} \frac{\partial \tilde{u}}{\partial \theta} = -\frac{\partial \tilde{v}}{\partial r}. \quad (19)$$

4 Week 4

4.1 Analytic functions

4.1.1 Gradient

Theorem 4 *The gradient of ϕ is*

$$\nabla \phi = \frac{\partial \phi}{\partial x} \underline{i} + \frac{\partial \phi}{\partial y} \underline{j} + \frac{\partial \phi}{\partial z} \underline{k}. \quad (20)$$

4.1.2 Directional derivative

Theorem 5 *The directional derivative of ϕ in the direction of a unit vector \underline{n} is*

$$\frac{\partial \phi}{\partial n}(\underline{r}) = \left. \frac{\partial}{\partial s} \phi(\underline{r} + s\underline{n}) \right|_{s=0} \quad (21)$$

$$= \left(n_1 \frac{\partial \phi}{\partial x_1} + n_2 \frac{\partial \phi}{\partial x_2} + n_3 \frac{\partial \phi}{\partial x_3} \right)(\underline{r}) = \underline{n} \cdot \nabla \phi(\underline{r}). \quad (22)$$

When s is small

$$\phi(\underline{r} + s\underline{n}) - \phi(\underline{r}) \approx s \frac{\partial \phi}{\partial n}(\underline{r}) = (s\underline{n}) \cdot \nabla \phi(\underline{r}). \quad (23)$$

4.1.3 Analytic function definition

Definition 6 *A function that is analytic holds the Cauchy-Riemann equations true.*

5 Week 5

5.1 Analytic functions

5.1.1 Harmonic functions

Theorem 6 *$\phi(x, y)$ is harmonic if*

$$\nabla^2 \phi = \frac{\partial^2 \phi}{\partial x^2} + \frac{\partial^2 \phi}{\partial y^2} = 0. \quad (24)$$

5.1.2 Harmonic Conjugate

Theorem 7 If $f = u + iv$ is analytic then u and v are harmonic functions. v is said to be the harmonic conjugate of u .

6 Week 6

6.1 Elementary functions of z

6.1.1 Representation of polynomials and zeros

Polynomials are entire functions and can be represented in several ways.

$$\begin{aligned} p_n(z) &= \sum_{k=0}^n a_k z^k \\ &= \sum_{k=0}^n \frac{p_n^{(k)}(0)}{k!} z^k, \text{ finite Maclaurin series,} \\ &= \sum_{k=0}^n \frac{p_n^{(k)}(z_0)}{k!} (z - z_0)^k, \text{ Taylor polynomial,} \\ &= a_n(z - \alpha_1)(z - \alpha_2) \cdots (z - \alpha_n), \text{ in terms of the zeros,} \\ &= a_n(z - \zeta_1)^{r_1}(z - \zeta_2)^{r_2} \cdots (z - \zeta_m)^{r_m}, \end{aligned}$$

where ζ_1, \dots, ζ_m are the distinct zeros and $r_1 + \cdots + r_m = n$.

At the zero ζ_k of multiplicity r_k we have

$$p_n(\zeta_k) = p'(\zeta_k) = \cdots = p_n^{(r_k-1)}(\zeta_k) = 0, p_n^{(r_k)}(\zeta_k) \neq 0. \quad (25)$$

6.1.2 Rational functions

Theorem 8 A rational function is the ratio of two polynomials, p, q , such that

$$R(z) = \frac{p(z)}{q(z)}, \quad q(z) = (z - \zeta_1)(z - \zeta_2) \cdots (z - \zeta_n). \quad (26)$$

where ζ_1, \dots, ζ_n are singular points.

If the limits exists as $z \rightarrow \zeta_k$ then ζ_k is a removable singularity.

Otherwise $R(z)$ has a pole singularity at ζ_k .

A simple pole is the case when $1/R(z)$ has a simple zero at ζ_k .

The order of the pole of $R(z)$ is the multiplicity of the zero of $1/R(z)$.

6.1.3 Partial fractions representation

From eq. (26), when $\deg p(z) < \deg q(z)$ and the zeros of $q(z)$ are simple we have the partial fraction representation of the form

$$R(z) = \frac{p(z)}{q(z)} = \sum_{k=1}^n \frac{A_k}{z - \zeta_k}. \quad (27)$$

When $\deg p(z) \geq \deg q(z)$ and the zeros of $q(z)$ are simple we have a representation of the form

$$R(z) = \frac{p(z)}{q(z)} = (\text{some polynomial}) + \sum_{k=1}^n \frac{A_k}{z - \zeta_k}. \quad (28)$$

In either case, A_k is the residue at ζ_k .

6.1.4 Residues

When $R(z)$ is in the form of eq. (28), to get A_k we have

$$\begin{aligned} A_k &= \lim_{z \rightarrow \zeta_k} (z - \zeta_k) R(z) = \lim_{z \rightarrow \zeta_k} \frac{(z - \zeta_k) p(z)}{q(z)}, \\ &= p(\zeta_k) \lim_{z \rightarrow \zeta_k} \frac{(z - \zeta_k)}{q(z)} = \frac{p(\zeta_k)}{q'(\zeta_k)}. \end{aligned} \quad (29)$$

When $q(z)$ has a zero at ζ of multiplicity $r \geq 1$ we need terms involving

$$\frac{1}{z - \zeta}, \frac{1}{(z - \zeta)^2}, \dots, \frac{1}{(z - \zeta)^r}.$$

The general case is as follows.

Let

$$R(z) = \frac{p(z)}{q(z)}. \quad (30)$$

We re-label to concentrate on one of the zeros of $q(z)$ at ζ and write

$$R(z) = \dots + \frac{B_1}{z - \zeta} + \dots + \frac{B_r}{(z - \zeta)^r} + \dots. \quad (31)$$

Then

$$(z - \zeta)^r R(z) = B_r + B_{r-1}(z - \zeta) + \dots + B_1(z - \zeta)^{r-1} + (z - \zeta)^r (\text{a function at } \zeta). \quad (32)$$

To get the residue B_1 we have

$$B_1 = \frac{1}{(r-1)!} \lim_{z \rightarrow \zeta} \left(\frac{d^{r-1}}{dz^{r-1}} (z - \zeta)^r R(z) \right). \quad (33)$$

A similar type of calculation is done for the other coefficients.

7 Week 7

7.1 Elementary functions of z

7.1.1 General case of the residues

Let

$$R(z) = \frac{p(z)}{(z - \zeta_1)^{r_1} (z - \zeta)^{r_2} \dots (z - \zeta_n)^{r_n}}. \quad (34)$$

With the procedures above we can get the coefficients in the following candidate representation of $R(z)$.

$$\left(\frac{A_{1,1}}{z - \zeta_1} + \dots + \frac{A_{r_1,1}}{(z - \zeta_1)^{r_1}} \right) + \dots + \left(\frac{A_{1,n}}{z - \zeta_n} + \dots + \frac{A_{r_n,n}}{(z - \zeta_n)^{r_n}} \right). \quad (35)$$

The coefficients are

$$A_{i,j} = \frac{1}{(r_j - i)!} \lim_{z \rightarrow \zeta_j} \left(\frac{d^{r_j - i}}{dz^{r_j - i}} (z - \zeta_j)^{r_j} R(z) \right), \quad i = 1, 2, \dots, r_j. \quad (36)$$

7.1.2 Complex trig functions

We define

$$\begin{aligned} \cosh z &= \frac{1}{2}(e^z + e^{-z}), & \sinh z &= \frac{1}{2}(e^z - e^{-z}), \\ \cos z &= \frac{1}{2}(e^{iz} + e^{-iz}), & \sin z &= \frac{1}{2i}(e^{iz} - e^{-iz}). \end{aligned}$$

As in the real case

$$\begin{aligned} \frac{d}{dz} \cosh z &= \sinh z, & \frac{d}{dz} \sinh z &= \cosh z, \\ \frac{d}{dz} \cos z &= -\sin z, & \frac{d}{dz} \sin z &= \cos z. \end{aligned}$$

We also have the identities

$$\cos^2 z + \sin^2 z = \cosh^2 z - \sinh^2 z = 1. \quad (37)$$

For all $z_1, z_2 \in \mathbb{C}$ we have the addition formulas

$$\sin(z_1 \pm z_2) = \sin z_1 \cos z_2 \pm \cos z_1 \sin z_2, \quad (38)$$

$$\cos(z_1 \pm z_2) = \cos z_1 \cos z_2 \mp \sin z_1 \sin z_2. \quad (39)$$

7.1.3 The real and imaginary parts of $\sin(z)$ and $\cos(z)$

With $z = x + iy$, $x, y \in \mathbb{R}$ we have

$$\sin(x + iy) = \sin x \cosh y + i \cos x \sinh y, \quad (40)$$

$$\cos(x + iy) = \cos x \cosh y - i \sin x \sinh y. \quad (41)$$

The real and imaginary parts of these functions are hence harmonic functions.

8 Week 8

8.0.1 Exponential function

$$e^z \equiv \exp(z) := e^x e^{iy} = e^x (\cos y + i \sin y). \quad (42)$$

As in the real case we have for all $z, z_1, z_2 \in \mathbb{C}$,

$$\frac{d}{dz} e^z = e^z, \quad e^{-z} = 1/e^z, \quad e^{z_1+z_2} = e^{z_1} e^{z_2}. \quad (43)$$

The function $w = \exp(z)$ is periodic with period $2\pi i$ and is one-to-one on

$$G = \{z = x + iy : -\pi < y \leq \pi\} \quad (44)$$

with inverse

$$\text{Log } w = \ln |w| + i \text{Arg } w \quad (45)$$

which is the principal valued logarithm.

8.0.2 cot and tanh

$$\cot z = \frac{\cos z}{\sin z}, \quad \tan z = \frac{\sin z}{\cos z} = -\cot(z + \pi/2) = \frac{1}{\tan(\pi/2 - z)}. \quad (46)$$

$\cot z$ has simple poles at $k\pi$ and $\tan z$ has simple poles at $\pi/2 + k\pi$ where $k \in \mathbb{Z}$.

8.0.3 Log z and the multi-valued $\log z$

Definition 7 The principal valued logarithm is

$$\text{Log } z = \ln |z| + i \text{Arg } z. \quad (47)$$

The multi-valued version $w = \log z$ means all complex numbers such that

$$e^w = z$$

and the set of values is

$$\{\text{Log } z + 2k\pi i : k \in \mathbb{Z}\}.$$

In both cases

$$e^{\text{Log } z} = e^{\log z} = z.$$

8.0.4 Complex powers z^α

Definition 8 The principal value of z^α is defined as

$$e^{\alpha \text{Log } z}. \quad (48)$$

The possibly multi-valued version of this is

$$e^{\alpha \log z} \quad (49)$$

9 Week 9

9.1 Integrals, arcs and contours

9.1.1 Series and the residue more generally

Taylor series: If $f(z)$ is analytic in the disk $|z - z_0| < R$ then

$$f(z) = \sum_{n=0}^{\infty} \frac{f^{(n)}(z_0)}{n!} (z - z_0)^n \quad (50)$$

and the series converges uniformly in $|z - z_0| \leq R' < R$.

Laurent series: If $f(z)$ is analytic in $0 < r < |z - z_0| < R$ then

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n + \sum_{n=1}^{\infty} \frac{a_{-n}}{(z - z_0)^n}, \quad (51)$$

there is uniform convergence in $0 \leq r < r_1 \leq |z - z_0| \leq R_1 < R$.

Both series are unique once z_0 is specified.

All the coefficients can be written as loop integrals.

The coefficients a_{-1} is the residue at z_0 when $r = 0$.

9.2 Cauchy theorems

Let f be a function which is analytic in a domain D and let Γ be a positively orientated loop in D and let z be a point inside D .

Theorem 9 The Cauchy-Goursat theorem:

$$\oint_{\Gamma} f(\zeta) d\theta = 0. \quad (52)$$

Theorem 10 The Cauchy integral formula:

$$f(z) = \frac{1}{2\pi i} \oint_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta. \quad (53)$$

Theorem 11 The generalised Cauchy integral formula:

$$f^{(n)}(z) = \frac{n!}{2\pi i} \oint_{\Gamma} \frac{f(\zeta)}{(\zeta - z)^{n+1}} d\zeta, \quad n = 0, 1, 2, \dots \quad (54)$$

9.2.1 Integral of a complex valued function

If $f : [a, b] \rightarrow \mathbb{C}$ with $f = u + iv$, $u, v \in \mathbb{R}$ then

$$\int_a^b f(x) dx = \int_a^b u(x) dx + i \int_a^b v(x) dx. \quad (55)$$

9.2.2 A smooth arc

Definition 9 A set $\gamma \subset \mathbb{C}$ is a smooth arc if the set can be described in the form

$$\{z(t) : a \leq t \leq b\} \quad (56)$$

where $z'(t)$ is continuous on $[a, b]$ and $z'(t) \neq 0$ on $[a, b]$.

The arc is said to be closed if the starting point $z(a)$ and the end point $z(b)$ are the same.

If the arc is not closed then we also require that $z(t)$ is one-to-one on $[a, b]$, i.e. it does not intersect itself.

If the arc is closed then we require that $z(t)$ is *one-to-one* on $[a, b]$ with

$$z(b) = z(a), \quad z'(b) = z'(a).$$

Definition 10 A smooth arc with a specific ordering of points is known as a directed smooth arc,

9.2.3 A contour

Definition 11 A contour is one point or a finite sequence of directed smooth arcs γ_k with the end of γ_k being the start of arc γ_{k+1} .

Theorem 12 Let $\gamma = \{z(t) : a \leq t \leq b\}$ and let $a = t_0 < t_1 < \dots < t_m = b$. The length of the arc is approximately

$$\sum_{i=1}^m |z(t_i) - z(t_{i-1})|. \quad (57)$$

When $t - t_{i-1}$ is small, we have

$$l(\gamma) = \text{length of } \gamma = \int_a^b |z'(t)| dt. \quad (58)$$

Definition 12 Let $a = t_0 < t_1 < \dots < t_m = b$ and let

$$A_m = \sum_{i=1}^m h_i f(z(t_{i-1/2})), \quad h_i = z(t_i) - z(t_{i-1}).$$

$$\begin{aligned} h_i f(z(t_{i-1/2})) &= (z(t_i) - z(t_{i-1})) f(z(t_{i-1/2})) \\ &\approx f(z(t_{i-1/2})) z'(t_{i-1/2})(t_i - t_{i-1}). \end{aligned}$$

$$\int_{\gamma} f(z) dz = \lim_{\substack{m \rightarrow \infty \\ \max_i |h_i| \rightarrow 0}} A_m = \int_a^b f(z(t)) z'(t) dt. \quad (59)$$

The value here does not depend on which particular valid parameterisation $z(t)$ that we use to describe γ .

9.2.4 The ML inequality

Lemma 1 Let M and L be defined by

$$M = \max_{z \in \Gamma} |f(z)| \quad \text{and} \quad L = \text{length of } \Gamma. \quad (60)$$

From the bound on $|f(z)|$ and the triangle inequality we have

$$\left| \sum_{i=1}^m h_i f(z(t_{i-1/2})) \right| \leq \sum_{i=1}^m |h_i| |f(z(t_{i-1/2}))| \leq M \sum_{i=1}^m |h_i| \leq ML.$$

As the bound above is independent of m and as the integral is an appropriate limit of such a sum we have

$$\left| \int_{\gamma} f(z) dz \right| \leq ML. \quad (61)$$

9.2.5 Independence of path when $f = F'$

Theorem 13 If there exists an anti-derivative F along the path then

$$\frac{d}{dt} F(z(t)) = F'(z(t)) z'(t) = f(z(t)) z'(t). \quad (62)$$

This is the integrand in the expression for the contour integral.

Theorem 14 Suppose that the function $f(z)$ is continuous in a domain D and has an anti-derivative $F(z)$ throughout D . Then for any contour Γ contained in D with initial point z_I and an end point z_E we have

$$\int_{\Gamma} f(z) dz = F(z_E) - F(z_I). \quad (63)$$

10 Week 10

10.1 Loop integrals

10.1.1 Closed loops and powers of z

Let Γ denote a closed loop.

Let $n \in \mathbb{Z}$ and $z_0 \in \mathbb{C}$.

When $n \neq -1$ the anti-derivative of $(z - z_0)^n$ is $(z - z_0)^{n+1}/(n+1)$ and as a consequence

$$\oint_{\Gamma} (z - z_0)^n dz = 0. \quad (64)$$

When $n = -1$ the function $1/(z - z_0)$ has an anti-derivative $\text{Log}(z - z_0)$ but this function is discontinuous on a branch cut starting from z_0 . The value of the integral depends on whether z_0 is inside or outside the loop.

$$\oint_{\Gamma} \frac{dz}{z - z_0} = \begin{cases} 2\pi i, & \text{if } z_0 \text{ is inside } \Gamma, \\ 0, & \text{if } z_0 \text{ is outside } \Gamma. \end{cases}$$

The integral does not exist in the usual sense when z_0 is on Γ .

10.1.2 Path independence, loop integrals and anti-derivatives

The following are equivalent statements involving the integral of f :

1. All loop integrals of f are 0.
2. The value of the integral of f only depends on the end points.
3. There exists an anti-derivative F , i.e. $F' = f$.

10.1.3 Loop integrals and rational functions

Theorem 15 If z_1, \dots, z_m are points inside Γ at which $R(z)$ has poles then

$$\begin{aligned} \oint_{\Gamma} R(z) dz &= \sum_{k=1}^m A_k \oint_{\Gamma} \frac{dz}{z - z_k}, \\ &= 2\pi i \sum_{k=1}^m A_k. \end{aligned} \quad (65)$$

11 Week 17

11.1 Harmonic functions - further results

Suppose that $f(z) = u(x, y) + iv(x, y)$ is analytic in a domain D with $u, v \in \mathbb{R}$. We have the following:

- As all derivatives of f exist and are analytic it follows that all partial derivatives of u and v exist and are continuous.
- Both u and v are harmonic, i.e. $\nabla^2 u = 0$ and $\nabla^2 v = 0$, with v being the harmonic conjugate of u .
-

$$f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = \frac{\partial v}{\partial y} - i \frac{\partial u}{\partial y}. \quad (66)$$

As f' is analytic the first partial derivatives of u and v are harmonic functions. All the partial derivatives of u and v are harmonic.

11.1.1 Creating an analytic function from a harmonic function

Suppose we have a function ϕ which is harmonic in a domain D . Let

$$g(z) = \frac{\partial \phi}{\partial x} - i \frac{\partial \phi}{\partial y}.$$

Define

$$u_1 = \frac{\partial \phi}{\partial x} \quad \text{and} \quad v_1 = -\frac{\partial \phi}{\partial y}. \quad (67)$$

Check that the Cauchy-Riemann equations are satisfied.

If we now restrict D to be a simply connected domain then the property that g is analytic in D implies that all loop integrals of g are 0 and this in turn implies that g has an anti-derivative G in D , i.e. $G' = g$. If we represent G as $G = u + iv$, $u, v, \in \mathbb{R}$ then as G is analytic we now have

$$g = G' = \frac{\partial u}{\partial x} - i \frac{\partial u}{\partial y} = \frac{\partial \phi}{\partial x} - i \frac{\partial \phi}{\partial y}. \quad (68)$$

This implies that $\phi = u + C$, where C is a constant, and we know that a function v exists which is a harmonic conjugate of ϕ .

Theorem 16 *When a domain D is simply connected and u is a harmonic in D there exists a harmonic conjugate v such that $u + iv$ is analytic in D .*

This result is **not** true when D is an annulus.

12 Week 18

12.1 Functions defined by loop integrals

Let Γ denote a closed loop traversed once in the anti-clockwise direction and let $g(\zeta)$ denote any continuous function defined on Γ . If we define

$$G(z) = \oint_{\Gamma} \frac{g(\zeta)}{\zeta - z} d\zeta \quad (69)$$

then this defines an analytic function for z inside Γ and it also defines an analytic function for z outside Γ . As an example:

$$g(\zeta) = \frac{1}{2\pi i} \left(\frac{f(\zeta)}{\zeta - z_0} \right) \quad \text{gives} \quad G(z) = \begin{cases} f'(z_0), & \text{if } z = z_0, \\ \frac{f(z) - f(z_0)}{z - z_0}, & \text{if } z \neq z_0. \end{cases} \quad (70)$$

In this case $g(\zeta)$ is defined inside the loop and has a singularity at $\zeta = z_0$.

12.2 Loop integrals of $f(z)/q(z)$ where $q = \text{polynomial}$

Suppose $f(z)$ is analytic inside a loop and

$$q(z) = (z - z_1)^{r_1} (z - z_2)^{r_2} \dots (z - z_n)^{r_n},$$

with $r_k \geq 1$ for $k = 1, \dots, n$. Using partial fractions we get

$$\frac{f(z)}{q(z)} = f(z) \left(\dots + \frac{A_{1,k}}{z - z_k} + \dots + \frac{A_{r_k,k}}{(z - z_k)^{r_k}} + \dots \right). \quad (71)$$

By using the generalised Cauchy integral formula on each term for each point z_k inside Γ we can determine

$$\oint_{\Gamma} \frac{f(z)}{q(z)} dz. \quad (72)$$

12.3 Other versions of the formulae and entire functions

The Cauchy integral formula when $\Gamma = \text{circle of radius } R$:

$$f^{(n)}(z_0) = \frac{n!}{2\pi R^n} \int_0^{2\pi} f(-R + Re^{it}) e^{-int} dt. \quad (73)$$

This is used to show that a bounded entire function is a constant (Liouville's theorem) by considering what happens when $n = 1$ and $R \rightarrow \infty$.

12.3.1 The fundamental theorem of algebra

Theorem 17 *Every non-constant polynomial with complex coefficients has at least one zero.*

12.3.2 Further Results

Lemma 2 *If $f(z)$ is analytic in a domain D and $|f(z)|$ is constant in D then $f(z)$ is a constant.*

12.3.3 The mean value property

$$f(z_0) = \frac{1}{2\pi} \int_0^{2\pi} f(z_0 + Re^{it}) dt, \quad |f(z_0)| \leq \frac{1}{2\pi} \int_0^{2\pi} |f(z_0 + Re^{it})| dt. \quad (74)$$

Lemma 3 *Suppose $f(z)$ is analytic in a disk centered at z_0 . If the maximum value of $|f(z)|$ over the disk is the value at the centre, i.e. the value $|f(z_0)|$, then $f(z)$ is constant on the disk.*

12.3.4 The maximum modulus theorem

Theorem 18 *If f is analytic in a domain D and $|f(z)|$ achieves its maximum value at a point $z_0 \in D$ then f is a constant in D .*

12.4 Definitions: sequences in \mathbb{C}

- A sequence z_0, z_1, z_2, \dots converges to z if for every $\epsilon > 0$ there exists an $N = N(\epsilon)$ such that

$$|z_n - z| < \epsilon \quad \text{for all } n \geq N. \quad (75)$$

- A sequence z_0, z_1, z_2, \dots is a Cauchy sequence if for every $\epsilon > 0$ there exists an $N = N(\epsilon)$ such that

$$|z_n - z_m| < \epsilon \quad \text{for all } n \geq N \text{ and } m \geq M. \quad (76)$$

12.4.1 Result about convergence

A sequence in \mathbb{C} converges if and only if it is a Cauchy sequence.

12.5 Definitions: series in \mathbb{C}

- Let c_0, c_1, c_2, \dots denote a sequence. A series is an expression of the form

$$c_0 + c_1 + c_2 + \dots \quad \text{and we write as } \sum_{k=0}^{\infty} c_k. \quad (77)$$

The sequence of partial sums are given by

$$s_n = \sum_{k=0}^n c_k, \quad n = 0, 1, 2, \dots \quad (78)$$

- The series converges if the sequence of partial sums converges and it diverges if the sequence of partial sums diverges. When we have convergence we say that

$$s = \sum_{k=0}^{\infty} c_k \quad (79)$$

is the sum of the series.

- If $\sum |c_k|$ converges then $\sum c_k$ is absolutely convergent.

12.5.1 Results about series in \mathbb{C}

- If a series $\sum c_k$ converges then $c_n \rightarrow 0$ as $n \rightarrow \infty$.
- If the series $\sum |c_k|$ converges then $\sum c_k$ converges.
- **Comparison test:** If there exists K such that $|c_k| \leq M_k$ for all $k \geq K$ and $\sum M_k$ converges then $\sum c_k$ converges.
- From the identity

$$(1 - c)(1 + c + c^2 + \cdots + c^n) = 1 - c^{n+1} \quad (80)$$

we have that the geometric series

$$\sum_{k=0}^{\infty} c^k = \frac{1}{1 - c}, \quad \text{when } |c| < 1. \quad (81)$$

- **Ratio test:** If $|c_{k+1}/c_k| \rightarrow L$ as $k \rightarrow \infty$ then the series converges if $L < 1$ and it diverges if $L > 1$.
- **Root test:** If $|c_k|^{1/k} \rightarrow L$ as $k \rightarrow \infty$ then the series converges if $L < 1$ and it diverges if $L > 1$.

12.6 Series of functions

Suppose that $f_0(z), f_1(z), \dots$ are all defined on D and let

$$F_n(z) = \sum_{k=0}^n f_k(z), \quad n = 0, 1, 2, \dots \quad (82)$$

$\sum f_k(z)$ converges pointwise on D if $(F_n(z))$ converges $\forall z \in D$. The sequence converges uniformly to $F(z)$ on D if

$$\sup_{z \in D} |F_n(z) - F(z)| \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (83)$$

A sufficient condition for a series to converge uniformly is the Weierstrass M-test: If $|f_k(z)| \leq M_k$ for all $z \in D$ and $\sum M_k$ converges then the series converges uniformly in D .

Uniform convergence preserves continuity: If $F_n(z)$, $n = 0, 1, 2, \dots$ are continuous in D and $F_n \rightarrow F$ uniformly on D then the limit function $F(z)$ is also continuous in D .

12.7 Uniform convergence and analytic functions

Theorem 19 Let $F_n(z)$ be a sequence of analytic functions in a simply connected domain D and converging uniformly to $F(z)$ in D . Then $F(z)$ is analytic in D .

13 Week 19

13.1 Taylor series for analytic functions

If $f(z)$ is analytic at z_0 then the series

$$f(z_0) + f'(z_0)(z - z_0) + \frac{f''(z_0)}{2!}(z - z_0)^2 + \cdots = \sum_{k=0}^{\infty} \frac{f^{(k)}(z_0)}{k!}(z - z_0)^k \quad (84)$$

is called the Taylor series for $f(z)$ around z_0 .

Theorem 20 If $f(z)$ is analytic in the disk $|z - z_0| < R$ then the Taylor series converges to $f(z)$ for all z in this disk and in any closed disk $|z - z_0| \leq R' < R$ the convergence is uniform.

13.1.1 Key formula in the proof of the Taylor series

$$f(z) = \sum_{k=0}^n \frac{f^{(k)}(z_0)}{k!}(z - z_0)^k + T_n(z) \quad (85)$$

where

$$T_n(z) = \frac{(z - z_0)^{n+1}}{2\pi i} \oint_C \frac{f(\zeta)}{(\zeta - z_0)^{n+1}(\zeta - z)} d\zeta. \quad (86)$$

It can be shown that $\max\{|T_n(z)| : |z - z_0| \leq R'\} \rightarrow 0$ as $n \rightarrow \infty$.

13.1.2 Taylor's series, comments about R

If $f(z)$ is analytic at z_0 then the Taylor series is

$$f(z) = \sum_{k=0}^{\infty} \frac{f^{(k)}(z_0)}{k!} (z - z_0)^k. \quad (87)$$

If $f(z)$ is analytic in $|z - z_0| < R$ then the series converges to $f(z)$ in this disk with uniform convergence in $|z - z_0| \leq R' < R$ for all $R' < R$.

If $f(z)$ is not an entire function then the largest R is such that $f(z)$ has a non-analytic point on $|z - z_0| = R$.

13.2 Maclaurin series case

Maclaurin series is the case of Taylor series when $z_0 = 0$.

$$f(z) = \sum_{k=0}^{\infty} \frac{f^{(k)}(0)}{k!} z^k. \quad (88)$$

If $f(z)$ is analytic in $|z| < R$ then the series converges to $f(z)$ in this disk with uniform convergence in $|z| \leq R' < R$ for all $R' < R$.

13.3 Real coefficients, even functions, odd functions, etc.

If $f(z) = u(x, y) + iv(x, y)$ is real when z is real then

$$v(x, 0) = 0 \quad \text{and} \quad f^{(n)}(0) = \left. \frac{\partial^n u(x, 0)}{\partial x^n} \right|_{x=0} \text{ is real.} \quad (89)$$

If $R =$ radius of convergence and $0 < r < R$ then we have

$$\frac{f^{(n)}(0)}{n!} = \frac{1}{2\pi r^n} \int_{-\pi}^{\pi} f(re^{it}) e^{-int} dt \quad (90)$$

$$= \frac{1}{2\pi r^n} \int_0^{\pi} (f(re^{it}) + (-1)^n f(-re^{it})) e^{-int} dt. \quad (91)$$

If $f(-z) = f(z)$ then the Maclaurin series only has even powers.

If $f(-z) = -f(z)$ then the Maclaurin series only has odd powers.

13.4 Series you are expected to know

$$\frac{1}{1-z} = 1 + z + z^2 + \cdots + z^n + \cdots, \quad \text{valid for } |z| < 1 \quad (92)$$

$$e^z = 1 + z + \frac{z^2}{2!} + \cdots + \frac{z^n}{n!} + \cdots \quad (93)$$

$$e^{-z} = 1 - z + \frac{z^2}{2!} + \cdots + \frac{-z^n}{n!} + \cdots \quad (94)$$

$$\cos(z) = 1 - \frac{z^2}{2!} + \frac{z^4}{4!} + \cdots \quad (95)$$

$$\sin(z) = z - \frac{z^3}{3!} + \frac{z^5}{5!} + \cdots \quad (96)$$

$$\cosh(z) = 1 + \frac{z^2}{2!} + \frac{z^4}{4!} + \cdots \quad (97)$$

$$\sinh(z) = z + \frac{z^3}{3!} + \frac{z^5}{5!} + \cdots \quad (98)$$

$$(99)$$

14 Week 20

14.1 The Koebe function, de Branges' theorem and a conjecture

The Koebe function says

$$f(z) = \frac{z}{(1-z)^2} = z + 2z^2 + 3z^3 + \cdots + nz^n + \cdots. \quad (100)$$

Suppose that you consider all functions $g(z)$ which are analytic in the unit disk, are one-to-one and satisfy $g(0) = 0$ and $g'(0) = 1$. Such functions have Maclaurin series of the form

$$g(z) = z + a_2z^2 + a_3z^3 + \cdots + a_nz^n + \cdots \quad (101)$$

de Branges proved that $|a_n| \leq n$.

Bieberbach proved that $|a_2| \leq 2$.

14.2 Multiplying series - the Cauchy product

If $f(z)$ and $g(z)$ are both analytic in $|z - z_0| < R$ then $h(z) = f(z)g(z)$ is also analytic in $|z - z_0| < R$.
Let $z_0 = 0$.

$$f(z) = a_0 + a_1z + a_2z^2 + \cdots, \quad (102)$$

$$g(z) = b_0 + b_1z + b_2z^2 + \cdots, \quad (103)$$

$$h(z) = c_0 + c_1z + c_2z^2 + \cdots. \quad (104)$$

We get

$$c_n = a_0b_n + a_1b_{n-1} + \cdots + a_nb_0. \quad (105)$$

This expression for c_n is known as the Cauchy product.

14.3 Leibnitz's formula for the n th derivative of a product

Repeatedly using the product rule gets

$$h^{(n)} = \sum_{k=0}^n \binom{n}{k} f^{(k)} f^{(n-k)}. \quad (106)$$

This is known as Leibnitz's rule for the n th derivative of a product.

14.4 The generalised L'Hopital's rule

If

$$g(z_0) = g'(z_0) = \cdots = g^{(m-1)}(z_0) = 0 \quad \text{and} \quad g^{(m)}(z_0) \neq 0 \quad (107)$$

and if

$$f(z_0) = f'(z_0) = \cdots = f^{(m-1)}(z_0) = 0 \quad (108)$$

then for z near z_0 we have

$$f(z) = a_m(z - z_0)^m + a_{m+1}(z - z_0)^{m+1} + \cdots, \quad (109)$$

$$g(z) = b_m(z - z_0)^m + b_{m+1}(z - z_0)^{m+1} + \cdots, \quad (110)$$

$$\frac{f(z)}{g(z)} \rightarrow \frac{a_m}{b_m} = \frac{f^{(m)}(z_0)}{g^{(m)}(z_0)} \quad \text{as } z \rightarrow z_0. \quad (111)$$

If the multiplicity of the zero of $g(z)$ at z_0 is greater than the multiplicity of the zero of $f(z)$ then there is no limit and $f(z)/g(z)$ has a singularity at z_0 .

14.5 Power series

A series of the form

$$\sum_{n=0}^{\infty} a_n(z - z_0)^n. \quad (112)$$

The series always converges at $z = z_0$. When it converges at other points the region where it converges is a disk $\{z : |z - z_0| < R\}$ and it is analytic in the disk.

The largest R is the radius of convergence. When $R < \infty$, $\{z : |z - z_0| = R\}$ is the circle of convergence. In all cases

$$R = \frac{1}{\limsup |a_n|^{1/n}}. \quad (113)$$

$R = 0$ when we only have convergence at $z = z_0$.

$R = \infty$ when we have convergence for all z .

15 Week 21

15.1 Cauchy-Hadamard theorem

Theorem 21

$$\sum_{n=0}^{\infty} a_n(z - z_0)^n \quad (114)$$

has radius of convergence $R = \frac{1}{\alpha}$, where α is the limit of

$$b_n = \sup\{|a_m|^{1/m} : m \geq n\} \geq 0 \quad (115)$$

for the sequence $|a_n|^{1/n}$.

15.2 Properties of a function defined by a power series

Let

$$f(z) = \sum_{n=0}^{\infty} a_n(z - z_0)^n, \quad R = \frac{1}{\limsup |a_n|^{1/n}}. \quad (116)$$

When $R > 0$ this defines an analytic function in $|z - z_0| < R$.

One way to relate the coefficients a_n to the derivatives of $f(z)$ is to use the generalized Cauchy integral formula. We take a loop Γ in the disk with z_0 inside the loop.

$$\frac{f^{(m)}(z_0)}{m!} = \frac{1}{2\pi i} \oint_{\Gamma} \frac{f(z)}{(z - z_0)^{m+1}} dz \quad (117)$$

$$= \frac{1}{2\pi i} \sum_{n=0}^{\infty} \oint_{\Gamma} \frac{f(z)}{(z - z_0)^{n-(m+1)}} dz. \quad (118)$$

The only integral in the last line which is non-zero is when $n - (m + 1) = -1$, i.e when $n = m$ we get

$$\frac{f^{(m)}(z_0)}{m!} = a_m. \quad (119)$$

15.3 Laurent series

A Laurent series is of the form

$$\sum_{n=-\infty}^{\infty} a_n(z - z_0)^n. \quad (120)$$

When it converges the region is an annulus $\{z : r < |z - z_0| < R\}$.

$$\sum_{n=-\infty}^{-1} a_n(z - z_0)^n, \quad \text{converges in } |z - z_0| > r. \quad (121)$$

$$\sum_{n=0}^{\infty} a_n(z - z_0)^n, \quad \text{converges in } |z - z_0| < R. \quad (122)$$

To have a function defined at some points we need the coefficients a_n to be such that $r < R$.

15.3.1 Classifying zeros and poles

When $f(z)$ has a zero of multiplicity $m \geq 1$ at z_0 we have

$$f(z) = a_m(z - z_0)^m + a_{m+1}(z - z_0)^{m+1} + \cdots = (z - z_0)^m g(z) \quad (123)$$

with $g(z)$ being analytic at z_0 and $g(z) = a_m \neq 0$.

If $f(z)$ has a removable singularity at z_0 then it has a Laurent series with no negative powers valid in $0 < |z - z_0| < R$, i.e.

$$f(z) = \sum_{n=0}^{\infty} a_n(z - z_0)^n \quad \text{and} \quad \lim_{z \rightarrow z_0} f(z) = a_0. \quad (124)$$

If $f(z)$ has a pole of order m then in $0 < |z - z_0| < R$ we have

$$f(z) = \sum_{n=-m}^{\infty} a_n(z - z_0)^n = \frac{\phi(z)}{(z - z_0)^m} \quad (125)$$

with $\phi(z)$ being analytic at z_0 and $\phi(z_0) = a_{-m} \neq 0$.

An essential singularity at z_0 has infinitely many negative powers

$$f(z) = \sum_{n=-\infty}^{\infty} a_n(z - z_0)^n, \quad 0 < |z - z_0| < R. \quad (126)$$

16 Week 22

16.1 Integrating a Laurent series - the residue

If we have the Laurent series valid for $0 < |z - z_0| < r$ of the form

$$f(z) = \sum_{n=-\infty}^{\infty} a_n(z - z_0)^n \quad (127)$$

and Γ is any loop in this region with z_0 as an interior point then

$$\oint_{\Gamma} f(z) dz = \sum_{n=-\infty}^{\infty} a_n \oint_{\Gamma} (z - z_0)^n dz = 2\pi i a_{-1}, \quad (128)$$

$$\text{Res}(f, z_0) = a_{-1} = \text{Residue of } f(z) \text{ at } z = z_0. \quad (129)$$

16.2 The Residue theorem

If z_1, z_2, \dots, z_n are isolated singularities inside Γ and C_1, C_2, \dots, C_n are non-intersecting circles traversed once in the anti-clockwise direction then $\Gamma \cup (-C_1) \cup \cdots \cup (-C_n)$ is the boundary of a region in which $f(z)$ is analytic and

$$\oint_{\Gamma} f(z) dz = \sum_{k=1}^n \oint_{C_k} f(z) dz \quad (130)$$

$$= 2\pi i \sum_{k=1}^n \text{Res}(f, z_k). \quad (131)$$

16.2.1 Techniques to calculate the residue

In the case of a simple pole of $f(z)$ at z_0 most examples for calculating the residue have involved calculating the limit

$$\text{Res}(f, z_0) = \lim_{z \rightarrow z_0} (z - z_0) f(z). \quad (132)$$

More generally, when we have a pole of order $m \geq 1$ we can calculate the residue by using

$$\text{Res}(f, z_0) = \frac{1}{(m-1)!} \lim_{z \rightarrow z_0} \frac{d^{m-1}}{dz^{m-1}} ((z - z_0)^m f(z)). \quad (133)$$

17 Week 23

17.1 The integrals on C_R^+ when we have a a^{imz} term

With $z = x + iy$, $miz = -my + imx$, $e^{imz} = e^{-my}e^{imx}$. When $m > 0$, $|e^{imz}| = e^{-my} < 1$ when $y \geq 0$.
When $\deg(Q) \geq \deg(P) + 2$ we have

$$\int_{C_R^+} \frac{P(z)}{Q(z)} dz \rightarrow 0 \quad \text{and} \quad \int_{C_R^+} \frac{P(z)}{Q(z)} e^{imz} dz \rightarrow 0 \quad (134)$$

as $R \rightarrow \infty$ by using the *ML* inequality.

When $\deg(Q) = \deg(P) + 1$ Jordan's lemma also gives

$$\int_{C_R^+} \frac{P(z)}{Q(z)} e^{imz} dz \rightarrow 0 \quad (135)$$

as $R \rightarrow \infty$.

When $\deg(Q) = \deg(P) + 1$ there is also a constant $A \geq 0$ such that

$$\left| \frac{Q(Re^{i\theta})iRe^{i\theta}}{Q(Re^{i\theta})} \right| \leq A, \quad \text{for sufficiently large } R. \quad (136)$$

17.2 Counting zeros and poles

Suppose that $f(z)$ is analytic in a domain except for a finite number of poles. Let

$$G(z) = \frac{f'(z)}{f(z)}. \quad (137)$$

Let z_0 be a zero of multiplicity m and let z_p be a pole of $f(z)$ of order n .

$$\text{Res}(G, z_0) = m, \quad \text{and} \quad \text{Res}(G, z_p) = -n. \quad (138)$$

Let $f(z)$ be analytic inside a simple loop C and let $N_0(f)$ be the number of zeros of $f(z)$ inside C .

$$N_0(f) = \frac{1}{2\pi i} \oint_C \frac{f'(z)}{f(z)} dz. \quad (139)$$

If $g(z)$ is also analytic inside C and $|g(z)| < |f(z)|$ on C then

$$N_0(f + g) = N_0(f). \quad (140)$$

This is Rouché's called theorem.