

Global exponential stability of periodic neural networks with time-varying delays

Haijun Jiang^{a,b,*}, Jinde Cao^a

^a*Department of Mathematics, Southeast University, Nanjing 210096, China*

^b*College of Mathematics and System Sciences, Xinjiang University, Urumqi 830046, China*

Received 14 April 2005; received in revised form 22 October 2005; accepted 12 January 2006

Communicated by T. Heskes

Available online 27 June 2006

Abstract

In this paper, we consider the periodic neural networks with variable coefficients and time-varying delays, and investigate the existence of periodic solution and its global exponential stability. Various sufficient conditions ensuring the existence of periodic and its global exponential stability are given. The results obtained in this paper extend and generalize those given in previous literature.

© 2006 Elsevier B.V. All rights reserved.

Keywords: Neural networks; Variable coefficients; Time-varying delays; Global exponential stability; Periodic solution

1. Introduction

Recurrently connected neural networks, sometimes called Grossberg–Hopfield neural networks, have attracted increasing interest in both theoretical studies and engineering applications. Many important results on the existence and uniqueness of equilibrium point, global asymptotic stability and global exponential stability for autonomous neural networks have been established and successfully applied to signal processing system, especially in static image treatment, and to solve nonlinear algebraic equations, such application rely on the qualitative properties of stability.

Among the most popular models in the literature of autonomous neural networks is the following continuous-time delayed neural networks.

$$\begin{aligned} \frac{du_i(t)}{dt} = & -d_i u_i(t) + \sum_{j=1}^n a_{ij} g_j(u_j(t)) \\ & + \sum_{j=1}^n b_{ij} f_j(u_j(t - \tau_{ij})) + I_i, \quad i = 1, 2, \dots, n, \end{aligned} \quad (1)$$

where activation functions g_j and f_j satisfy certain defining conditions, and d_i, a_{ij}, b_{ij}, I_i ($i, j = 1, 2, \dots, n$) are constants.

All these neural networks have been extensively studied. In [1,2,4–6,8–11,14,17–21,23,26,27,29], the authors have obtained many a criterion for checking the existence, uniqueness, global asymptotical stability, global exponential stability and robust stability of neural networks. Several techniques have been introduced, and the combination of Lyapunov function (or functional) method with inequality technique and the combination of Lyapunov function (or functional) method with LMI technique have been widely adopted. But all results are focused upon autonomous neural networks.

However, as we well know, the nonautonomous phenomenon often occurs in many realistic systems. Particularly, when we consider a long-time dynamical behavior of a system, the parameters of the system usually will raise change along with time. In addition, in many

*Corresponding author. College of Mathematics and System Sciences, Xinjiang University, Urumqi 830046, China.

E-mail addresses: jianghai@xju.edu.cn (H. Jiang), jdcao@seu.edu.cn (J. Cao).

applications, the property of periodic oscillatory solutions of a neural networks also is of great interest. Therefore, the research on the nonautonomous neural networks with delays is very important in like manner.

In this paper, we will consider the following dynamical systems and their periodic limits:

$$\begin{aligned} \frac{du_i(t)}{dt} = & -d_i(t)u_i(t) + \sum_{j=1}^n a_{ij}(t)g_j(u_j(t)) \\ & + \sum_{j=1}^n b_{ij}(t)f_j(u_j(t - \tau_{ij}(t))) + I_i(t), \end{aligned} \quad (2)$$

where $i = 1, 2, \dots, n$, $a_{ij}(t), b_{ij}(t), I_i(t) : \mathbb{R}^+ \rightarrow \mathbb{R}$ are continuously periodic functions with period $\omega > 0$, i.e., $d_i(t) = d_i(t + \omega)$, $a_{ij}(t) = a_{ij}(t + \omega)$, $b_{ij}(t) = b_{ij}(t + \omega)$, $I_i(t) = I_i(t + \omega)$, $\tau_{ij}(t) = \tau_{ij}(t + \omega)$ for all $t > 0$ and $i, j = 1, 2, \dots, n$.

From the mathematical point of view, system (2) is completely different from system (1), and a known mathematical method do not directly apply. Here, we develop some methods from functional differential equations (see [3,13,25]) to study the existence of periodic solution and its global exponential stability of system (2).

In fact, there have been considerable research on the nonautonomous neural networks (see [7,12,15,16,22,24,28,30]). In [22], when $n = 1$, the authors investigate the periodic solutions and almost periodic solution of the following first-order nonautonomous differential equation

$$\dot{x}(t) = -d(t)u(t) + a(t)f(u(t)) + I(t), \quad t > 0 \quad (3)$$

when $f(u) = \tanh(x)$ is a sigmoid-type function, $d(t) > 0$, $a(t)$ and $I(t)$ are continuous ω -periodic function with respect to the time variable t . Under assumptions $d(t) \geq a(t)$ and $1/\omega \int_0^\omega [d(s) - |a(s)|] ds = \mu > 0$, the authors obtain a sufficient condition to ensure the existence, uniqueness and globally exponential stability of periodic solution for periodic system (3). In [7], by combining the theory of the exponential dichotomy and Lyapunov function, the authors study the existence and attractivity of almost periodic solutions for cellular neural networks with distributed delays and variable coefficients. In [15,16,24], the cellular neural networks with variable coefficients and time-varying delays are studied, by using the Lyapunov functional method, the techniques of matrix analysis and inequality analysis, the authors established a series of the criteria on the boundedness, global exponential stability and the existence of periodic solutions. In [12,30], the recurrent neural networks and BAM neural networks with periodic coefficients and delays are studied. Some new sufficient conditions ensuring the existence, uniqueness and global exponential stability of the periodic solution were obtained by using the continuation theorem based on coincidence degree and the Lyapunov functional method. In [28], the authors discuss the dynamical system with variable coefficients and constant delays, obtained the existence of periodic solution and its globally exponential stability.

In this paper, we will discuss the existence of periodic solution and its global exponential stability for nonautonomous system (2). We do not assume that the activations functions are bounded. We also do not use existing complicated theory to prove the existence of periodic solution. Utilizing the method given in [28], we obtain the existence of periodic solution and its globally exponential stability for dynamical system (2) by constructing a new Lyapunov function. We will see that the results obtained in this paper will extend and generalize the corresponding results existing in [12,28,30].

This paper is organized as follows. In Section 2, we will give some preliminaries. In Section 3, we will establish new criteria on the existence of periodic solution and its global exponential stability for periodic nonautonomous system (2). In Section 4, an example is given to illustrate the results obtained in this paper. In Section 5, we will give some concluding remarks of the results.

2. Some preliminaries

Definition 1. Class $H\{G_1, G_2, \dots, G_n\}$ of functions: let $G = \text{diag}[G_1, G_2, \dots, G_n]$, where $G_i > 0$ ($i = 1, 2, \dots, n$). $g(x) = (g_1(x), g_2(x), \dots, g_n(x))^T$ is said to belong to $H\{G_1, G_2, \dots, G_n\}$, if the functions $g_i(x)$ ($i = 1, 2, \dots, n$) satisfy

$$\frac{|g_i(x+u) - g_i(x)|}{|u|} \leq G_i.$$

Let $\tau = \sup\{\tau_{ij}(t) : t \in [0, +\infty), i, j = 1, 2, \dots, n\}$. We denote by $C[-\tau, 0]$ the Banach space of continuous functions $u(\cdot) = (u_1(\cdot), u_2(\cdot), \dots, u_n(\cdot))^T : [-\tau, 0] \rightarrow \mathbb{R}^n$ with the following norm:

$$\|u(\cdot)\|_{(\xi, p)} = \left[\frac{1}{n} \sum_{i=1}^n \xi_i |u_i(\cdot)|^p \right]^{1/p}.$$

In this paper we always assume that all solutions of system (2) satisfy the following initial conditions:

$$u_i(\theta) = \phi_i(\theta), \quad \theta \in [-\tau, 0], \quad i = 1, 2, \dots, n, \quad (4)$$

where $\phi = (\phi_1, \phi_2, \dots, \phi_n) \in C[-\tau, 0]$. It is well known that by the fundamental theory of functional differential equations (see [3,13,25]), system (2) has a unique solution $u(t) = (u_1(t), u_2(t), \dots, u_n(t))$ satisfying the initial condition (4).

In the following, the Young inequality and Dini derivative will be used to prove the main results of this paper. On the Young inequality we have the following lemma.

Lemma 1 (Cao [4] and Zheng and Chen [28]). Let $a > 0, b > 0, p > 1, q > 1$, and $1/p + 1/q = 1$. Then we have the inequality

$$ab \leq \frac{1}{p}(ae)^p + \frac{1}{q}(be^{-1})^q,$$

where $\varepsilon > 0$. The equality holds if and only if

$$(a\varepsilon)^p = (b\varepsilon^{-1})^q.$$

On the Dini derivative we have the following lemma.

Lemma 2 (Chen et al. [8]). Assume that $u(t)$ is a differentiable function defined on R_+ . Then for any $t \in R_+$ the Dini right derivative $D^+|u(t)|$ of function $|u(t)|$ exists and has the expression as follows:

$$D^+|u(t)| = \limsup_{h \rightarrow 0^+} \frac{1}{h} [|u(t+h)| - |u(t)|] = \sigma(u(t))\dot{u}(t),$$

where

$$\sigma(u(t))\dot{u}(t) = \begin{cases} \dot{u}(t) & \text{if } u(t) > 0 \text{ or } u(t) = 0 \text{ and } \dot{u}(t) > 0, \\ -\dot{u}(t) & \text{if } u(t) < 0 \text{ or } u(t) = 0 \text{ and } \dot{u}(t) < 0, \\ 0 & \text{if } u(t) = 0 \text{ and } \dot{u}(t) = 0 \end{cases}$$

and $\dot{u}(t) = du(t)/dt$.

In this paper, for system (2) we introduce the following assumptions.

(H₁) Functions $d_i(t)$, $a_{ij}(t)$, $b_{ij}(t)$ and $I_i(t)$ ($i, j = 1, 2, \dots, n$) are bounded and continuous functions defined on $t \in R_+ = [0, \infty)$.

(H₂) Functions $f(x) = (f_1(u), f_2(u), \dots, f_n(u))^T \in H\{F_1, F_2, \dots, F_n\}$ and $g(u) = (g_1(u), g_2(u), \dots, g_n(u))^T \in H\{G_1, G_2, \dots, G_n\}$.

(H₃) $\inf_{t \in R_+} \{1 - \dot{\tau}_{ij}(t)\} > 0$, in which $\dot{\tau}_{ij}(t)$ ($i, j = 1, 2, \dots, n$) denotes the derivative of $\tau_{ij}(t)$.

3. Main results

In this section, we discuss the existence of periodic solution and its global stability. We propose a general and concise approach and give theorems on the existence of periodic solution and its global exponential stability.

Theorem 1. Suppose that (H₁) – (H₃) hold and there are constants $\xi_i > 0$, α_{ij} , $\beta_{ij} \in R$ ($i, j = 1, 2, \dots, n$) such that

$$\begin{aligned} & d_i(t)\xi_i - \sum_{j=1}^n \frac{1}{p} G_i \xi_j |a_{ji}(t)|^{\alpha_{ij}p} - \sum_{j=1}^n \frac{1}{q} G_j \xi_i |a_{ij}(t)|^{(1-\alpha_{ij})q} \\ & - \sum_{j=1}^n \frac{1}{q} F_j \xi_i |b_{ij}(t)|^{(1-\beta_{ij})p} - \sum_{j=1}^n \frac{1}{p} F_i \xi_j \frac{|b_{ji}(\psi_{ji}^{-1}(t))|^{\beta_{ji}p}}{1 - \dot{\tau}_{ji}(\psi_{ji}^{-1}(t))} > 0 \end{aligned} \quad (5)$$

for all $t \geq 0$ and $i = 1, 2, \dots, n$, where $\psi_{ij}^{-1}(t)$ is the inverse function of $\psi_{ij}(t) = t - \tau_{ij}(t)$ ($i, j = 1, 2, \dots, n$) and $1 \leq p < \infty$ is a constant. Then the dynamical system (2) has a unique periodic solution $v(t) = [v_1(t), v_2(t), \dots, v_n(t)]^T$ and, for any solution $u(t) = [u_1(t), u_2(t), \dots, u_n(t)]^T$ of system (2), there exists a scalar $\varepsilon > 0$ such that

$$|u_i(t + j\omega) - v_i(t)| = O(e^{-j\varepsilon\omega}) \quad i = 1, 2, \dots, n.$$

Proof. Define a function δ :

$$\begin{aligned} \delta(\varepsilon) = \min_{0 \leq i \leq n} \left\{ & d_i(t)\xi_i - \sum_{j=1}^n \frac{1}{p} G_i \xi_j |a_{ji}(t)|^{\alpha_{ij}p} \right. \\ & - \sum_{j=1}^n \frac{1}{q} G_j \xi_i |a_{ij}(t)|^{(1-\alpha_{ij})q} - \sum_{j=1}^n \frac{1}{q} F_j \xi_i e^{\varepsilon\tau} |b_{ij}(t)|^{(1-\beta_{ij})p} \\ & \left. - \sum_{j=1}^n \frac{1}{p} F_i \xi_j e^{\varepsilon\tau} \frac{|b_{ji}(\psi_{ji}^{-1}(t))|^{\beta_{ji}p}}{1 - \dot{\tau}_{ji}(\psi_{ji}^{-1}(t))} - \varepsilon \xi_i \right\}. \end{aligned}$$

Obviously, δ is continuous on R with respect to ε . From the inequality (5), we can see that $\delta(0) > 0$, so there exists a positive number $\varepsilon > 0$ such that $\delta(\varepsilon) > 0$.

Further, let $\bar{u}_i(t) = u_i(t + \omega) - u_i(t)$, $\bar{g}_i(u_i(t)) = g_i(u_i(t + \omega)) - g_i(u_i(t))$, $\bar{f}_i(u_i(t)) = f_i(u_i(t + \omega)) - f_i(u_i(t))$ and $w_i(t) = e^{\varepsilon t} \bar{u}_i(t)$, $i = 1, 2, \dots, n$. Defining a Lyapunov function as follows:

$$\begin{aligned} L(t) = & \sum_{i=1}^n \xi_i |w_i(t)|^p + \sum_{i=1}^n \sum_{j=1}^n \xi_i F_j e^{\varepsilon\tau} \\ & \times \int_{t-\tau_{ij}(t)}^t \frac{|b_{ij}(\psi_{ij}^{-1}(s))|^{\beta_{ij}p}}{1 - \dot{\tau}_{ij}(\psi_{ij}^{-1}(s))} |w_j(s)|^p ds. \end{aligned}$$

Calculating the Dini upper right derivative of $L(t)$ with respect to t , we have

$$\begin{aligned} D^+L(t) = & \sum_{i=1}^n \xi_i p |w_i(t)|^{p-1} e^{\varepsilon t} \text{sign}(\bar{u}_i(t)) \left[-d_i(t)\bar{u}_i(t) + \varepsilon \bar{u}_i(t) \right. \\ & + \sum_{j=1}^n a_{ij}(t)\bar{g}_j(u_j(t)) + \sum_{j=1}^n b_{ij}(t)\bar{f}_j(u_j(t - \tau_{ij}(t))) \left. \right] \\ & + \sum_{i=1}^n \sum_{j=1}^n e^{\varepsilon\tau} \xi_i F_j \frac{|b_{ij}(\psi_{ij}^{-1}(t))|^{\beta_{ij}p}}{1 - \dot{\tau}_{ij}(\psi_{ij}^{-1}(t))} |w_j(t)|^p \\ & - \sum_{i=1}^n \sum_{j=1}^n e^{\varepsilon\tau} \xi_i F_j |b_{ij}(t)|^{\beta_{ij}p} |w_j(t - \tau_{ij}(t))|^p \\ \leq & p \sum_{i=1}^n \xi_i \left[-(d_i(t) - \varepsilon) |w_i(t)|^p \right. \\ & + \sum_{j=1}^n |a_{ij}(t)| e^{\varepsilon t} |\bar{g}_j(u_j(t))| |w_i(t)|^{p-1} \\ & \times \sum_{j=1}^n |b_{ij}(t)| e^{\varepsilon t} |\bar{f}_j(u_j(t - \tau_{ij}(t)))| |w_i(t)|^{p-1} \left. \right] \\ & + \sum_{i=1}^n \sum_{j=1}^n e^{\varepsilon\tau} \xi_i F_j \frac{|b_{ij}(\psi_{ij}^{-1}(t))|^{\beta_{ij}p}}{1 - \dot{\tau}_{ij}(\psi_{ij}^{-1}(t))} |w_j(t)|^p \\ & - \sum_{i=1}^n \sum_{j=1}^n e^{\varepsilon\tau} \xi_i F_j |b_{ij}(t)|^{\beta_{ij}p} |w_j(t - \tau_{ij}(t))|^p. \end{aligned} \quad (6)$$

Since

$$e^{\varepsilon t} |\bar{f}_j(u_j(t - \tau_{ij}(t)))| \leq F_j |w_j(t - \tau_{ij}(t))| e^{\varepsilon\tau}, \quad (7)$$

$$e^{\varepsilon t} |\bar{g}_j(u_j(t))| \leq G_j |w_j(t)|. \quad (8)$$

Further, from (6)–(8), we have

$$\begin{aligned}
 D^+L(t) \leq & p \sum_{i=1}^n \xi_i \left[-(d_i(t) - \varepsilon)|w_i(t)|^p \right. \\
 & + \sum_{j=1}^n |a_{ij}(t)|G_j|w_j(t)||w_i(t)|^{p-1} \\
 & + \sum_{j=1}^n |b_{ij}(t)|e^{\varepsilon\tau}F_j|w_j(t - \tau_{ij}(t))||w_i(t)|^{p-1} \left. \right] \\
 & + \sum_{i=1}^n \sum_{j=1}^n e^{\varepsilon\tau} \xi_i F_j \frac{|b_{ij}(\psi_{ij}^{-1}(t))|^{\beta_{ij}p}}{1 - \tau_{ij}(\psi_{ij}^{-1}(t))} |w_j(t)|^p \\
 & - \sum_{i=1}^n \sum_{j=1}^n e^{\varepsilon\tau} \xi_i F_j |b_{ij}(t)|^{\beta_{ij}p} |w_j(t - \tau_{ij}(t))|^p. \quad (9)
 \end{aligned}$$

Estimating the right-hand side of (9) by using Young inequality, we have that

$$\begin{aligned}
 & \sum_{j=1}^n |a_{ij}(t)|G_j|w_j(t)||w_i(t)|^{p-1} \\
 & = \sum_{j=1}^n G_j \{ |a_{ij}(t)|^{\alpha_{ij}p} |w_j(t)|^p \}^{1/p} \\
 & \quad \times \{ |a_{ij}(t)|^{(1-\alpha_{ij})q} |w_i(t)|^p \}^{1/q} \\
 & \leq \sum_{j=1}^n G_j \left[\frac{1}{p} |a_{ij}(t)|^{\alpha_{ij}p} |w_j(t)|^p \right. \\
 & \quad \left. + \frac{1}{q} |a_{ij}(t)|^{(1-\alpha_{ij})q} |w_i(t)|^p \right] \quad (10)
 \end{aligned}$$

and

$$\begin{aligned}
 & \sum_{j=1}^n |b_{ij}(t)|e^{\varepsilon\tau}F_j|w_j(t - \tau_{ij}(t))||w_i(t)|^{p-1} \\
 & = \sum_{j=1}^n F_j e^{\varepsilon\tau} \{ |b_{ij}(t)|^{\beta_{ij}p} |w_j(t - \tau_{ij}(t))|^p \}^{1/p} \\
 & \quad \times \{ |b_{ij}(t)|^{(1-\beta_{ij})q} |w_i(t)|^p \}^{1/q} \\
 & \leq \sum_{j=1}^n F_j e^{\varepsilon\tau} \left[\frac{1}{p} |b_{ij}(t)|^{\beta_{ij}p} |w_j(t - \tau_{ij}(t))|^p \right. \\
 & \quad \left. + \frac{1}{q} |b_{ij}(t)|^{(1-\beta_{ij})q} |w_i(t)|^p \right]. \quad (11)
 \end{aligned}$$

From (9)–(11), we finally have

$$\begin{aligned}
 D^+L(t) \leq & p \sum_{i=1}^n \xi_i \left\{ -(d_i(t) - \varepsilon)|w_i(t)|^p \right. \\
 & + \sum_{j=1}^n G_j \left[\frac{1}{p} |a_{ij}(t)|^{\alpha_{ij}p} |w_j(t)|^p \right. \\
 & \left. \left. + \frac{1}{q} |a_{ij}(t)|^{(1-\alpha_{ij})q} |w_i(t)|^q \right] \right\}
 \end{aligned}$$

$$\begin{aligned}
 & + \sum_{j=1}^n F_j e^{\varepsilon\tau} \left[\frac{1}{p} |b_{ij}(t)|^{\beta_{ij}p} |w_j(t - \tau_{ij}(t))|^p \right. \\
 & \left. + \frac{1}{q} |b_{ij}(t)|^{(1-\beta_{ij})q} |w_i(t)|^q \right] \left. \right\} \\
 & + \sum_{i=1}^n \sum_{j=1}^n \xi_i F_j \frac{|b_{ij}(\psi_{ij}^{-1}(t))|^{\beta_{ij}p}}{1 - \tau_{ij}(\psi_{ij}^{-1}(t))} e^{\varepsilon\tau} |w_j(t)|^p \\
 & - \sum_{i=1}^n \sum_{j=1}^n \xi_i F_j e^{\varepsilon\tau} |b_{ij}(t)|^{\beta_{ij}p} |w_j(t - \tau_{ij}(t))|^p \\
 & \leq p \sum_{i=1}^n \{ -(d_i(t) - \varepsilon)\xi_i + \sum_{j=1}^n \frac{1}{p} G_i \xi_j |a_{ji}(t)|^{\alpha_{ji}p} \\
 & + \sum_{j=1}^n \frac{1}{q} G_j \xi_i |a_{ij}(t)|^{(1-\alpha_{ij})q} \\
 & + \sum_{j=1}^n \frac{1}{q} F_j \xi_i e^{\varepsilon\tau} |b_{ij}(t)|^{(1-\beta_{ij})q} \\
 & + \sum_{j=1}^n \frac{1}{p} F_i \xi_j e^{\varepsilon\tau} \frac{|b_{ji}(\psi_{ji}^{-1}(t))|^{\beta_{ji}p}}{1 - \tau_{ji}(\psi_{ji}^{-1}(t))} \} |w_i(t)|^p \\
 & = -p \sum_{i=1}^n \delta_i |w_i(t)|^p \leq 0.
 \end{aligned}$$

Therefore, $L(t)$ is bounded, which implies

$$\sum_{i=1}^n \xi_i |e^{\varepsilon t} \tilde{u}_i(t)|^p \quad (12)$$

is bounded and

$$|u_i(t + \omega) - u_i(t)| = O(e^{-\varepsilon t}), \quad i = 1, 2, \dots, n. \quad (13)$$

Now, define a function $v(t) = [v_1(t), v_2(t), \dots, v_n(t)]^T$ by

$$v_i(t) = \lim_{j \rightarrow \infty} u_i(t + j\omega).$$

Because of

$$u_i(t + j\omega) = u_i(t) + \sum_{k=1}^n \{u_i(t + k\omega) - u_i(t + (k-1)\omega)\}$$

and (13), $v(t)$ is well defined and is a periodic solution with period ω . Moreover,

$$|u_i(t + j\omega) - v_i(t)| = O(e^{-j\varepsilon\omega}) \quad \text{when } j \rightarrow \infty. \quad (14)$$

Next, we prove the uniqueness of periodic solution of system (2). If $u(t), v(t)$ are two solutions with period ω of system (2). By similar method used before, it is easy to prove

$$|u_i(t + j\omega) - v_i(t + j\omega)| = O(e^{-j\varepsilon\omega}) \quad \text{when } j \rightarrow \infty. \quad (15)$$

which means that the limit solution is unique. Theorem 1 is proved completely. \square

As direct consequences of Theorem 1, we have the following corollaries.

Corollary 1. Suppose that (H₁)–(H₃) hold and there are $\xi_i > 0$ ($i = 1, 2, \dots, n$) such that

$$d_i(t)\xi_i - \sum_{j=1}^n G_i \xi_j |a_{ji}(t)| - \sum_{j=1}^n F_i \xi_j \frac{|b_{ji}(\psi_{ji}^{-1}(t))|}{1 - \tau_{ji}(\psi_{ji}^{-1}(t))} > 0 \quad (16)$$

for all $t \geq 0$. Then the dynamical system (2) has a unique periodic solution $v(t) = [v_1(t), v_2(t), \dots, v_n(t)]^T$ and, for any solution $u(t) = [u_1(t), u_2(t), \dots, u_n(t)]^T$ of system (2), there exists a scalar $\varepsilon > 0$ such that

$$|u_i(t + j\omega) - v_i(t)| = O(e^{-j\varepsilon\omega}) \quad i = 1, 2, \dots, n.$$

In fact, in condition (5) taking $p = 1$ and $\alpha_{ij} = \beta_{ij} = 1$ ($i, j = 1, 2, \dots, n$), by Theorem 1, we easily obtain the result of Corollary 1.

Remark 1. In [30], the authors obtained the existence and globally exponential stability of periodic solutions for periodic delayed neural networks by utilizing the Mawhin coincidence degree theory and the Lyapunov functional methods. In [15], the authors obtained the existence and global exponential stability of periodic solutions for periodic neural networks with variable coefficients and time-varying delays by using the existence theory of periodic solutions for general functional differential equations. However, in this paper, we obtain the existence and global exponential stability by utilizing the general and very concise approach. Furthermore, the criteria obtained in this paper are different from ones of [12,16,24].

When $\tau_{ij}(t) = \tau_{ij}$, where τ_{ij} ($i, j = 1, 2, \dots, n$) are constants, then system (2) degenerates into the following form:

$$\begin{aligned} \frac{du_i(t)}{dt} = & -d_i(t)u_i(t) + \sum_{j=1}^n a_{ij}(t)g_j(u_j(t)) \\ & + \sum_{j=1}^n b_{ij}(t)f_j(u_j(t - \tau_{ij})) + I_i(t), \end{aligned} \quad (17)$$

where $i = 1, 2, \dots, n$. For system (17), we have the following result.

Theorem 2. Suppose that (H₁), (H₂) hold and there are constants $\xi_i > 0$, $\alpha_{ij}, \beta_{ij} \in R$ ($i, j = 1, 2, \dots, n$) such that

$$\begin{aligned} d_i(t)\xi_i - \sum_{j=1}^n \frac{1}{p} G_i \xi_j |a_{ji}(t)|^{\alpha_{ji}p} - \sum_{j=1}^n \frac{1}{q} G_j \xi_i |a_{ij}(t)|^{(1-\alpha_{ij})q} \\ - \sum_{j=1}^n \frac{1}{q} F_j \xi_i |b_{ij}(t)|^{(1-\beta_{ij})p} - \sum_{j=1}^n \frac{1}{p} F_i \xi_j |b_{ji}(t)|^{\beta_{ji}p} > 0 \end{aligned} \quad (18)$$

for all $t \geq 0$ and $i = 1, 2, \dots, n$. Then the dynamical system (17) has a unique periodic solution $v(t) = [v_1(t), v_2(t), \dots, v_n(t)]^T$ and, for any solution $u(t) = [u_1(t), u_2(t), \dots, u_n(t)]^T$ of system (17), there exists a scalar $\varepsilon > 0$ such that

$$|u_i(t + j\omega) - v_i(t)| = O(e^{-j\varepsilon\omega}) \quad i = 1, 2, \dots, n.$$

In order to complete the proof of Theorem 2, we only need define the following Lyapunov function similar to the

proof of Theorem 1.

$$\begin{aligned} L_1(t) = & \sum_{i=1}^n \xi_i |w_i(t)|^p + \sum_{i=1}^n \sum_{j=1}^n \xi_i F_j e^{\varepsilon t} \\ & \times \int_{t-\tau_{ij}}^t |b_{ij}(t)| |w_j(s)|^p ds. \end{aligned}$$

Further, as consequences of Theorem 2, we also have the results similar to Corollary 1. Here, we omit it.

Remark 2. From Theorem 2, we can see that the results obtained in this paper improve and extend the main results given in [28].

For autonomous system (1), as consequences of Theorems 1 and 2, we have the following result.

Corollary 2. Suppose that (H₂) and (H₃) hold and there are constants $\xi_i > 0$, $\alpha_{ij}, \beta_{ij} \in R$ ($i, j = 1, 2, \dots, n$) such that

$$\begin{aligned} d_i \xi_i - \sum_{j=1}^n \frac{1}{p} G_i \xi_j |a_{ji}|^{\alpha_{ji}p} - \sum_{j=1}^n \frac{1}{q} G_j \xi_i |a_{ij}|^{(1-\alpha_{ij})q} \\ - \sum_{j=1}^n \frac{1}{q} F_j \xi_i |b_{ij}|^{(1-\beta_{ij})p} - \sum_{j=1}^n \frac{1}{p} F_i \xi_j |b_{ji}|^{\beta_{ji}p} > 0 \end{aligned} \quad (19)$$

for $i = 1, 2, \dots, n$ and $1 \leq p < \infty$ is a constant. Then the dynamical system (1) has a unique periodic equilibrium $v^* = [v_1^*, v_2^*, \dots, v_n^*]^T$ and, for any solution $u(t) = [u_1(t), u_2(t), \dots, u_n(t)]^T$ of system (1), there exists a scalar $\varepsilon > 0$ such that $|u_i(t) - v_i^*| = O(e^{-\varepsilon t})$ $i = 1, 2, \dots, n$.

In order to complete the proof of Corollary 2, we only need construct the following Lyapunov function:

$$\begin{aligned} L_2(t) = & \sum_{i=1}^n \xi_i |w_i(t)|^p + \sum_{i=1}^n \sum_{j=1}^n \xi_i F_j e^{\varepsilon t} \\ & \times \int_{t-\tau_{ij}}^t |b_{ij}(t)| |w_j(s)|^p ds. \end{aligned}$$

Similar to the proof of Theorem 1, we easily obtain the result of Corollary 2. Here, we omit it.

For autonomous system (1), we also have the following result.

Corollary 3. Suppose that (H₂) holds and there are constants $\xi_i > 0$ ($i = 1, 2, \dots, n$) such that

$$d_i \xi_i - \sum_{j=1}^n G_i \xi_j |a_{ji}| - \sum_{j=1}^n F_i \xi_j |b_{ji}| > 0. \quad (20)$$

Then the dynamical system (1) has a unique periodic solution $v(t) = [v_1(t), v_2(t), \dots, v_n(t)]^T$ and, for any solution $u(t) = [u_1(t), u_2(t), \dots, u_n(t)]^T$ of system (1), there exists a scalar $\varepsilon > 0$ such that

$$|u_i(t + j\omega) - v_i(t)| = O(e^{-j\varepsilon\omega}) \quad i = 1, 2, \dots, n.$$

Remark 3. Let the matrices $D = \text{diag}(d_1, d_2, \dots, d_n)$, $F = \text{diag}(F_1, F_2, \dots, F_n)$, $G = \text{diag}(G_1, G_2, \dots, G_n)$, $A = (a_{ij})_{n \times n}$, $|A| = (|a_{ij}|)_{n \times n}$, $B = (b_{ij})_{n \times n}$ and $|B| = (|b_{ij}|)_{n \times n}$.

We easily find that the condition (20) is equivalent to the matrix $D - (|A|^T G + |B|^T F)$ be a non-singular M -matrix. Thus, under this condition, we can obtain the result similar to Corollary 3.

Remark 4. It should be emphasized that parameters α_{ij}, β_{ij} play key role in Theorems 1–2 and Corollary 2. By appropriately choosing the parameters α_{ij}, β_{ij} (see Remark 3), we can see that it include many known assumptions as its special cases.

4. Illustrative example

Example. Let $n = 2$, $f_j(u) = g_j(u) = \frac{1}{2}(|x+1| - |x-1|)$, $c_i(t) = 5 + \sin t$, $a_{ij}(t) = \sin t$, $b_{ij}(t) = \cos t$, $\tau_{ij}(t) = \frac{1}{2}\sin t + 1$ ($i, j = 1, 2$). We consider the following 2-dimensional neural networks:

$$\begin{aligned} \frac{du_1(t)}{dt} &= -(5 + \sin t)u_1(t) + \sum_{j=1}^2 \sin t g_j(u_j(t)) \\ &\quad + \sum_{j=1}^2 \cos t f_j(u_j(t - \tau_{1j}(t))) + \sin t, \\ \frac{du_2(t)}{dt} &= -(5 + \sin t)u_2(t) + \sum_{j=1}^2 \sin t g_j(u_j(t)) \\ &\quad + \sum_{j=1}^2 \cos t f_j(u_j(t - \tau_{2j}(t))) + \cos t. \end{aligned} \quad (21)$$

Obviously, $\psi_{ij}(t) = t - \frac{1}{2}\sin t - 1$, $\dot{\psi}_{ij}(t) = 1 - \frac{1}{2}\cos t > 0$. Thus the inverse of $\psi_{ij}(t)$ ($i, j = 1, 2$) exist, we denote it by $\bar{\psi}(t)$. By taking $\xi_i = 1$, $F = (1, 1)$, $G = (1, 1)$, $p = 1$, $\alpha_{ij} = \beta_{ij} = 1$ ($i, j = 1, 2$) then the condition (5) transform into

$$\begin{aligned} d_i(t)\xi_i - \sum_{j=1}^n G_i \xi_j |a_{ji}(t)| - \sum_{j=1}^n F_i \xi_j \frac{|b_{ji}(\bar{\psi}_{ji}^{-1}(t))|}{1 - \bar{\tau}_{ji}(\bar{\psi}_{ji}^{-1}(t))} \\ = (5 + \sin t) - 2 \times \sin t - 2 \times \frac{\cos(\bar{\psi}(t))}{1 - \frac{1}{2}\cos(\bar{\psi}(t))} > 0 \end{aligned}$$

for all $t \geq 0$ and $i = 1, 2$. Hence system (21) has a unique globally exponentially stable 2π -periodic solution.

For numerical simulation, the following four cases are given: case 1 with the initial state $(\phi_1(t), \phi_2(t)) = (-0.01 \sin t + 0.1, 0.01 \cos t - 0.1)$ for $t \in [-2.5, 0]$; case 2 with the initial state $(\phi_1(t), \phi_2(t)) = (-0.025 \sin t + 0.1, 0.025 \cos t - 0.1)$ for $t \in [-2.5, 0]$; case 3 with the initial state $(\phi_1(t), \phi_2(t)) = (-0.04 \sin t + 0.1, 0.04 \cos t - 0.1)$ for $t \in [-2.5, 0]$; case 4 with the initial state $(\phi_1(t), \phi_2(t)) = (-0.0625 \sin t + 0.1, 0.0625 \cos t - 0.1)$ for $t \in [-2.5, 0]$; The following Fig. 1 depicts the time responses of state variables of $x_1(t)$ and $x_2(t)$ with step $h = 0.01$, and Fig. 2 depicts the phase plots of state variable $x_1(t)$ and $x_2(t)$. It confirms that the proposed condition leads to the unique and globally exponentially stable 2π -periodic solution for this model.




Fig. 1. Transient response of state variables $x_1(t)$ and $x_2(t)$ for Example 1.




Fig. 2. Phase plots of state variables $x_1(t)$ and $x_2(t)$ for Example 1.

5. Conclusions

In this paper, different from [12,15,16,24,30], we introduce the new research method, that is, by introducing ingeniously many real parameters α_{ij}, β_{ij} and ω_i , applying Young inequality technique and Dini derivative. We obtain new criteria on the existence, uniqueness and global exponential stability of the periodic solutions for the general periodic nonautonomous neural networks with variable coefficients and time-varying delays. The results obtained in this paper improve and extend many previous works, and they are easy to check and apply in practice.

Acknowledgements

The authors would like to thank the reviewers and the editor for their valuable suggestions and comments. This

work was supported by the National Natural Science Foundation of China under Grants 60574043, 60373067 and 10361004, the Natural Science Foundation of Jiangsu Province of China under Grant No. BK2003053, the Postdoctoral Science Foundation of Jiangsu Province, China under Grant 1660631111, and the Doctoral Foundation of Xinjiang University.

References

- [1] S. Arik, An analysis of global asymptotic stability of delayed cellular neural networks, *IEEE Trans. Neural Networks* 13 (2) (2002) 1239–1242.
- [2] S. Arik, Global asymptotic stability of a larger class of neural networks with constant time delay, *Phys. Lett. A* 311 (2003) 504–511.
- [3] T. Burton, *Stability and Periodic Solutions of Ordinary and Functional Differential Equations*, Academic Press, Orlando, 1985.
- [4] J. Cao, New results concerning exponential stability and periodic solutions of delayed cellular neural networks, *Phys. Lett. A* 307 (2003) 136–147.
- [5] J. Cao, D.W.C. Ho, A general framework for global asymptotic stability analysis of delayed neural networks based on LMI approach, *Chaos, Solitons & Fractals* 24 (2005) 1317–1329.
- [6] J. Cao, J. Wang, Global asymptotic and robust stability of recurrent neural networks with time delays, *IEEE Trans. Circuits Syst. I* 52 (2005) 417–426.
- [7] A. Chen, J. Cao, Existence and attractivity of almost periodic solutions for cellular neural networks with distributed delays and variable coefficients, *Appl. Math. Comput.* 134 (2003) 125–140.
- [8] A. Chen, J. Cao, L. Huang, An estimation of upper bound of delays for global asymptotic stability of delayed Hopfield neural networks, *IEEE Trans. Circuits Syst. I* 49 (7) (2002) 1028–1032.
- [9] A. Chen, J. Cao, L. Huang, Exponential stability of BAM neural networks with transmission delays, *Neurocomputing* 57 (2004) 435–454.
- [10] A. Chen, J. Cao, L. Huang, Global robust stability of interval cellular neural networks with time-varying delays, *Chaos, Solitons & Fractals* 23 (2005) 787–799.
- [11] M. Forti, On global asymptotic stability of a class of nonlinear system arising in neural networks theory, *J. Differential Equations* 113 (1994) 246–264.
- [12] S. Guo, L. Huang, B. Dai, Z. Zhang, Global existence of periodic solutions of BAM neural networks with variable coefficients, *Phys. Lett. A* 317 (2003) 97–106.
- [13] J. Hale, *Theory of Functional Differential Equations*, Springer, Heidelberg, 1977.
- [14] J.J. Hopfield, Neurons with graded response have collective computational properties like those of two-stage neurons, *Proc. Natl. Acad. Sci. USA (Biophys.)* 81 (1984) 3088–3092.
- [15] H. Jiang, Z. Li, Z. Teng, Boundedness and stability for nonautonomous cellular neural networks with delay, *Phys. Lett. A* 306 (2003) 313–325.
- [16] H. Jiang, Z. Teng, Global exponential stability of cellular neural networks with time-varying coefficients and delays, *Neural Networks* 17 (2004) 1415–1425.
- [17] X. Li, L. Huang, J. Wu, Further results on the stability of delayed cellular neural networks, *IEEE Trans. Circuits Syst. I* 50 (9) (2003) 1239–1242.
- [18] X. Li, L. Huang, H. Zhu, Global stability of cellular neural networks with constant and variable delays, *Nonlinear Anal.* 53 (2003) 319–333.
- [19] C. Li, X. Liao, New algebraic conditions for global exponential stability of delayed recurrent neural networks, *Neurocomputing* 64 (2005) 319–333.
- [20] H. Lu, Stability criteria for delayed neural networks, *Phys. Rev. E* 64 (2001) 1–13.
- [21] S. Mohamad, K. Gopalsamy, Exponential stability of continuous-time and discrete-time cellular neural networks with delays, *Appl. Math. Comput.* 135 (2003) 17–38.
- [22] S. Mohamad, K. Gopalsamy, Neuronal dynamica in time varying environments: continuous and discrete time models, *Discrete Continuous Dyn. Syst.* 6 (2000) 841–860.
- [23] J. Peng, H. Qiao, Z. Xu, A new approach to stability of neural networks with time-varying delays, *Neural Networks* 15 (2002) 95–103.
- [24] M. Rehim, H. Jiang, Z. Teng, Boundedness and stability for nonautonomous cellular neural networks with delay, *Neural Networks* 17 (2004) 1017–1025.
- [25] T. Yoshizawa, *Stability Theory and the Existence of Periodic Solutions and Almost Periodic Solutions*, Springer, Berlin, 1975.
- [26] G. Yu, C. Lu, J. Tsai, T. Su, B. Liu, Stability of cellular neural networks with time-varying delay, *IEEE Trans. Circuits Syst. I* 50 (5) (2003) 677–679.
- [27] Y. Zhang, Absolute periodicity and absolute stability of delayed neural networks, *IEEE Trans. Circuits Syst. I* 49 (2002) 256–261.
- [28] Y. Zheng, T. Chen, Global exponential stability of delayed periodic dynamical system, *Phys. Lett. A* 322 (2004) 344–355.
- [29] D. Zhou, J. Cao, Globally exponential stability conditions for cellular neural networks with time-varying delays, *Appl. Math. Comput.* 131 (2002) 487–496.
- [30] J. Zhou, Z. Liu, G. Chen, Dynamic of periodic delayed neural networks, *Neural Networks* 17 (2004) 87–101.



Haijun Jiang was born in Hunan, China, in 1968. He received the B.S. degree from the Mathematics Department, Yili Teacher College, Yili, Xinjiang, China; the M.S. degree from the Mathematics Department, East China Normal University, Shanghai, China, and the Ph.D. degree from the College of Mathematics and System Sciences, Xinjiang University, China, in 1990, 1994, and 2004, respectively. At present, he is a Post doctoral Research Fellow in the Department of Southeast University, Nanjing, China. He is an Associate Professor of College of Mathematics and System Sciences at Xinjiang University, Xinjiang, China. His present research interests include nonlinear systems, mathematical biology, empdemiology, and dynamics of neural networks.



Jinde Cao received the B.S. degree from Anhui Normal University, Wuhu, China, the M.S. degree from Yunnan University, Kunming, China, and the Ph.D. degree from Sichuan University, Chengdu, China, all in mathematics/applied mathematics, in 1986, 1989, and 1998, respectively. From March 1989 to May 2000, he was with Yunnan University. In May 2000, he joined the Department of Mathematics, Southeast University, Nanjing, China. From July 2001 to June 2002, he was a Post doctoral Research Fellow in the Department of Automation and Computer-aided Engineering, Chinese University of Hong Kong, Hong Kong. From August 2002 to October 2002, he was a Senior Visiting Scholar at the Institute of Mathematics, Fudan University, Shanghai, China. From February 2003 to May 2003, he was a Senior Research Associate in the Department of Mathematics, City University of Hong Kong, Hong Kong. From July 2003 to September 2003, he was a Senior Visiting Scholar in the Institute of Intelligent Machines, Chinese Academy of Sciences, Hefei, China. From January 2004 to April 2004, he was a Research Fellow in the Department of Manufacturing Engineering and Engineering Management, City University of Hong Kong, Hong Kong. From January 2005 to April 2005, he was a Research Fellow in the Department of Mathematics, City University of Hong Kong, Hong Kong.

From January 2006 to April 2006, he was a Research Fellow in the Department of Electronics Engineering, City University of Hong Kong, Hong Kong.

He is currently a Professor and Doctoral Advisor at the Southeast University. Prior to this, he was a Professor at Yunnan University from

1996 to 2000. He is the author or coauthor of more than 100 journal papers and five edited books and a reviewer of *Mathematical Reviews* and *Zentralblatt-MATH*. His research interests include nonlinear systems, neural networks, complex systems and complex networks, stability theory, and applied mathematics.