

# Dynamical behaviors of Cohen–Grossberg neural networks with delays and reaction–diffusion terms<sup>☆</sup>

Hongyong Zhao<sup>a,\*</sup>, Kunlun Wang<sup>b</sup>

<sup>a</sup>*Department of Mathematics, Nanjing University of Aeronautics and Astronautics, Nanjing 210016, PR China*

<sup>b</sup>*Department of Computer Science and Technology, Hefei University, Hefei 230022, PR China*

Received 28 June 2005; received in revised form 12 November 2005; accepted 23 November 2005

Communicated by T. Heskes

Available online 13 June 2006

## Abstract

In this paper, we study Cohen–Grossberg neural networks with delays and reaction–diffusion terms. By employing homotopic mapping theory and constructing suitable Lyapunov functional method we present some sufficient conditions ensuring the existence, uniqueness and globally exponential stability (GES) of the equilibrium point. These conditions obtained have important leading significance in the designs and applications of GES for reaction–diffusion neural circuit systems with delays. Finally, we show a numerical example to verify the theoretical analysis.

© 2006 Elsevier B.V. All rights reserved.

**Keywords:** Homotopic mapping theory; Cohen–Grossberg neural networks; Reaction–diffusion terms; Globally exponential stability; Equilibrium point; Lyapunov functional

## 1. Introduction

Cohen–Grossberg neural networks (CGNNs) were first introduced by Cohen–Grossberg [4]. CGNNs include Hopfield neural networks as special cases [1,15,17,18,3,20,21]. The class of networks has good application in parallel computation, associative memory and optimization problems, which has been one of the most active areas of research and has received much attention. One can refer to the articles [19,2,16,14,9] for detailed discussion on these aspects. Thus, from the viewpoint of application, the dynamical study for CGNNs is quite important and significant, and cannot be replaced with the dynamical study for the traditional neural networks such as Hopfield neural networks and cellular neural networks. Furthermore, in the analysis of dynamical CGNNs for parallel

computation and optimization, to increase the rate of convergence to the equilibrium point of the networks, it is necessary to ensure a desired exponential convergence rate of the networks trajectories, starting from arbitrary initial states to the equilibrium point which corresponds to the optimal solution. From the viewpoint of mathematics and engineering, it is required that the equilibrium point of CGNNs is globally exponential stability (GES). As a consequence, many authors have paid much effort to the research on GES of equilibrium point for CGNNs in Refs. [19,2,16,14,9]. However, strictly speaking, diffusion effects cannot be avoided in neural network models when electrons are moving in asymmetric electromagnetic field, thus we must consider the space is varying with the time. Refs. [11–13,7] have considered the stability of neural networks with diffusion terms, which are expressed by partial differential equations. It is also common to consider the diffusion in biological systems (such as immigration) [8,6,10]. To the best our knowledge, few authors study GES of Cohen–Grossberg neural networks with delays and reaction–diffusion terms.

Motivated by the above discussions, in this paper we analyze further problem of GES of Cohen–Grossberg

<sup>☆</sup>This research was supported by the National Natural Science Foundation of Nanjing University of Aeronautics and Astronautics, and also supported by the National Natural Science Foundation of the Education Department of Anhui, China.

\*Corresponding author. Tel.: +86025 84489977.

E-mail addresses: [hongyongz@126.com](mailto:hongyongz@126.com), [zhaohym@yahoo.com.cn](mailto:zhaohym@yahoo.com.cn) (H. Zhao).

neural networks with delays and reaction–diffusion terms, and give a set of sufficient conditions ensuring the existence, uniqueness and GES of the equilibrium point by using homotopic mapping theory and employing Lyapunov functional method.

## 2. Preliminary

Let  $C \triangleq C([- \tau, 0] \times R^m, R^n)$  be the Banach space of continuous functions which map  $[- \tau, 0] \times R^m$  into  $R^n$  with the topology of uniform convergence.  $\Omega$  be an open bounded domain in  $R^m$  with smooth boundary  $\partial\Omega$ , and  $mes\Omega > 0$  denotes the measure of  $\Omega$ .  $L^2(\Omega)$  is the space of real functions on  $\Omega$  which are  $L^2$  for the Lebesgue measure. It is a Banach space for the norm

$$\|u(t)\| = \sum_{i=1}^n \|u_i(t)\|_2,$$

where

$$u(t) = (u_1(t), \dots, u_n(t))^T, \quad \|u_i(t)\|_2 = \left( \int_{\Omega} |u_i(t, x)|^2 dx \right)^{1/2}.$$

For any  $\varphi(t, x) \in C([- \tau, 0] \times \Omega, R^n)$ , we define

$$\|\varphi\| = \sum_{i=1}^n \|\varphi_i\|_2,$$

where

$$\varphi(t, x) = (\varphi_1(t, x), \dots, \varphi_n(t, x))^T,$$

$$\|\varphi_i\|_2 = \left( \int_{\Omega} |\varphi_i(x)|_{\tau}^2 dx \right)^{1/2},$$

$$|\varphi_i(x)|_{\tau} = \sup_{-\tau \leq s \leq 0} |\varphi_i(s, x)|.$$

Consider the following delayed Cohen–Grossberg neural networks with reaction–diffusion terms

$$\begin{cases} \frac{\partial u_i(t, x)}{\partial t} = \sum_{l=1}^m \frac{\partial}{\partial x_l} \left( D_{il} \frac{\partial u_i(t, x)}{\partial x_l} \right) - a_i(u_i(t, x)) \\ \quad \times \left[ b_i(u_i(t, x)) - \sum_{j=1}^n a_{ij} s_j(u_j(t, x)) \right. \\ \quad \left. - \sum_{k=0}^K \sum_{j=1}^n t_{ij}^{(k)} s_j(u_j(t - \tau_k, x)) + I_i \right], \quad t \geq 0, \quad x \in \Omega \\ u_i(t, x) = \varphi_i(t, x), \quad -\tau \leq t \leq 0, \quad x \in \Omega \\ \frac{\partial u_i(t, x)}{\partial n} = 0, \quad x \in \partial\Omega, \end{cases} \quad (1)$$

where  $i = 1, \dots, n$ ;  $n$  is the number of neurons in the network;  $u_i(t, x)$  denotes the state variable associated with the  $i$ th neurons at time  $t$  and in space  $x$ ;  $a_i(t, x)$  represents

an amplification function;  $b_i(t, x)$  represents an appropriately behaved function;  $a_{ij}$  represents the strength of the neuron interconnections within the network;  $s_j(t, x)$  shows how the  $j$ th neuron reacts to the input;  $t_{ij}^{(k)}$  represents the interconnection with delay  $\tau_k$ ;  $\tau_k$  is delay and satisfies  $0 \leq \tau_k \leq \tau$ ;  $I_i$  is the constant input from outside the system;  $D_{ik} \geq 0$  corresponds to the transmission diffusion operator along the  $i$ th neuron.  $u(t, x) = (u_1(t, x), \dots, u_n(t, x))^T$ ,  $x = (x_1, \dots, x_m)^T$ .

Throughout the paper, we always assume that system (1) has a continuous solution denoted by  $u(t, 0, \varphi; x)$  or simply  $u(t, x)$  if no confusion should occur.

**Definition 1.**  $u(t, x) \equiv u^* \in R^n$  is said to be an equilibrium point of system (1), if the constant vector  $u^* = (u_1^*, \dots, u_n^*)^T$  satisfies

$$b_i(u_i^*) - \sum_{j=1}^n a_{ij} s_j(u_j^*) - \sum_{k=0}^K \sum_{j=1}^n t_{ij}^{(k)} s_j(u_j^*) + I_i = 0, \quad i = 1, \dots, n. \quad (2)$$

**Lemma 1** (Cronin [5] Kronecker's theorem). Assume that  $f: \overline{\Omega} \rightarrow R^n$  is a continuous function,  $\deg(f, \Omega, p) \neq 0$ , then there exists  $x_0 \in \Omega$ , such that  $f(x_0) = p$ .

**Lemma 2** (Cronin [5] Homotopy invariance theorem). Assume that  $H(x, \lambda): \overline{\Omega} \times [0, 1] \rightarrow R^n$  is a continuous function, denote  $h_{\lambda}(x) = H(x, \lambda)$ . When  $\lambda \in [0, 1]$  and  $p \notin h_{\lambda}(\partial\Omega)$ ,  $\deg(h_{\lambda}, \Omega, p)$  is independent of  $\lambda$ .

## 3. Existence and GES of the equilibrium point

In the paper, we always assume that

(H1) There exist  $m_i$  and  $M_i$ , such that

$$0 < m_i \leq a_i(u_i) \leq M_i, \quad i = 1, \dots, n.$$

(H2)  $b_i(\cdot)$  is differentiable,  $\alpha_i \triangleq \inf_{u_i \in R} \{ \dot{b}_i(u_i) \} > 0$ , where  $\dot{b}_i(\cdot)$  is the derivative of  $b_i(\cdot)$ ,  $b_i(0) = 0$ ,  $i = 1, \dots, n$ .

(H3) There exist constants  $\beta_j$  ( $j = 1, \dots, n$ ), such that

$$|s_j(x) - s_j(y)| \leq \beta_j |x - y|, \quad \forall x, y \in R.$$

(H4) There exist constants  $p_{ij}$ ,  $q_{ij}$ ,  $p_{ij}^*$ ,  $q_{ij}^*$ , such that

$$\begin{aligned} & -2m_i \alpha_i + \sum_{j=1}^n M_i |a_{ij}|^{2p_{ij}} \beta_j^{2q_{ij}} + \sum_{j=1}^n M_j |a_{ji}|^{2-2p_{ji}} \beta_i^{2-2q_{ji}} \\ & + \sum_{k=0}^K \sum_{j=1}^n M_i |t_{ij}^{(k)}|^{2p_{ij}^*} \beta_j^{2q_{ij}^*} \\ & + \sum_{k=0}^K \sum_{j=1}^n M_j |t_{ji}^{(k)}|^{2-2p_{ji}^*} \beta_i^{2-2q_{ji}^*} < 0. \end{aligned}$$

In order to study the existence and uniqueness of the equilibrium point, we consider the following algebraic

equations associated with system (1)

$$b_i(u_i(t, x)) - \sum_{j=1}^n a_{ij}s_j(u_j(t, x)) - \sum_{k=0}^K \sum_{j=1}^n t_{ij}^{(k)} s_j(u_j(t, x)) + I_i = 0, \quad i = 1, \dots, n. \quad (3)$$

System (3) can be rewritten in the following vector form:

$$B(u) - AS(u) - TS(u) + I = 0, \quad (4)$$

where  $u = (u_1, \dots, u_n)^T$ ,  $B(u) = (b_1(u_1), \dots, b_n(u_n))^T$ ,  $A = (a_{ij})_{n \times n}$ ,  $T = (\sum_{k=0}^K t_{ij}^{(k)})_{n \times n}$ ,  $S(u) = (s_1(u_1), \dots, s_n(u_n))^T$ ,  $I = (I_1, \dots, I_n)^T$ .

**Theorem 1.** If (H1)–(H4) holds, then system (1) has a unique equilibrium point  $u^*$ .

**Proof.** Let  $h(u) = B(u) - AS(u) - TS(u) + I$ . Define the homotopic mapping  $H(u, \lambda) : \bar{\Omega} \times [0, 1] \rightarrow R^n$  as follows:

$$H(u, \lambda) \triangleq (H_1(u, \lambda), \dots, H_n(u, \lambda))^T = \lambda h(u) + (1 - \lambda)B(u), \quad \lambda \in [0, 1],$$

in which  $\alpha = \text{diag}(\alpha_i)$ .

It follows from (H2) and (H4) that

$$|b_i(u_i)| \geq \alpha_i |u_i|, \quad i = 1, \dots, n, \quad (5)$$

$$|s_j(u_j)| \leq \beta_j |u_j| + |s_j(0)|, \quad j = 1, \dots, n. \quad (6)$$

Thus we have

$$\begin{aligned} |H_i(u, \lambda)| &= \left| \lambda \left( b_i(u_i) - \sum_{j=1}^n a_{ij}s_j(u_j) - \sum_{k=0}^K \sum_{j=1}^n t_{ij}^{(k)} s_j(u_j) + I_i \right) \right. \\ &\quad \left. + (1 - \lambda)b_i(u_i) \right| \geq \alpha_i |u_i| - \sum_{j=1}^n |a_{ij}| \beta_j |u_j| \\ &\quad - \sum_{j=1}^n |a_{ij}| |s_j(0)| - \sum_{k=0}^K \sum_{j=1}^n |t_{ij}^{(k)}| \beta_j |u_j| \\ &\quad - \sum_{k=0}^K \sum_{j=1}^n |t_{ij}^{(k)}| |s_j(0)| - |I_i|. \end{aligned} \quad (7)$$

By (7), we obtain

$$\begin{aligned} \sum_{i=1}^n M_i |u_i| |H_i(u, \lambda)| &\geq \sum_{i=1}^n M_i |u_i| \left( \alpha_i |u_i| - \sum_{j=1}^n |a_{ij}| \beta_j |u_j| - \sum_{j=1}^n |a_{ij}| |s_j(0)| \right. \\ &\quad \left. - \sum_{k=0}^K \sum_{j=1}^n |t_{ij}^{(k)}| \beta_j |u_j| - \sum_{k=0}^K \sum_{j=1}^n |t_{ij}^{(k)}| |s_j(0)| - |I_i| \right) \\ &\geq \sum_{i=1}^n M_i \left( \alpha_i |u_i|^2 - \sum_{j=1}^n |a_{ij}|^{p_{ij}} \beta_j^{q_{ij}} |u_i| |a_{ij}|^{1-p_{ij}} \beta_j^{1-q_{ij}} |u_j| \right. \\ &\quad \left. - \sum_{j=1}^n |a_{ij}| |s_j(0)| |u_i| \right) \end{aligned}$$

$$\begin{aligned} &- \sum_{k=0}^K \sum_{j=1}^n |t_{ij}^{(k)}|^{p_{ij}^*} \beta_j^{q_{ij}^*} |u_i| |t_{ij}^{(k)}|^{1-p_{ij}^*} \beta_j^{1-q_{ij}^*} |u_j| \\ &- \sum_{k=0}^K \sum_{j=1}^n |t_{ij}^{(k)}| |s_j(0)| |u_i| - |I_i| |u_i| \Big) \\ &\geq \sum_{i=1}^n M_i \left( \alpha_i |u_i|^2 - \frac{1}{2} \sum_{j=1}^n |a_{ij}|^{2p_{ij}} \beta_j^{2q_{ij}} |u_i|^2 \right. \\ &\quad \left. - \frac{1}{2} \sum_{j=1}^n |a_{ij}|^{2-2p_{ij}} \beta_j^{2-2q_{ij}} |u_j|^2 \right. \\ &\quad \left. - \sum_{j=1}^n |a_{ij}| |s_j(0)| |u_i| - \frac{1}{2} \sum_{k=0}^K \sum_{j=1}^n |t_{ij}^{(k)}|^{2p_{ij}^*} \beta_j^{2q_{ij}^*} |u_i|^2 \right. \\ &\quad \left. - \frac{1}{2} \sum_{k=0}^K \sum_{j=1}^n |t_{ij}^{(k)}|^{2-2p_{ij}^*} \beta_j^{2-2q_{ij}^*} |u_j|^2 \right. \\ &\quad \left. - \sum_{k=0}^K \sum_{j=1}^n |t_{ij}^{(k)}| |s_j(0)| |u_i| - |I_i| |u_i| \right) \\ &= \sum_{i=1}^n \left( M_i \alpha_i - \frac{1}{2} \sum_{j=1}^n M_i |a_{ij}|^{2p_{ij}} \beta_j^{2q_{ij}} \right. \\ &\quad \left. - \frac{1}{2} \sum_{j=1}^n M_j |a_{ji}|^{2-2p_{ji}} \beta_i^{2-2q_{ji}} - \frac{1}{2} \sum_{k=0}^K \sum_{j=1}^n M_i |t_{ij}^{(k)}|^{2p_{ij}^*} \beta_j^{2q_{ij}^*} \right. \\ &\quad \left. - \frac{1}{2} \sum_{k=0}^K \sum_{j=1}^n M_j |t_{ji}^{(k)}|^{2-2p_{ji}^*} \beta_i^{2-2q_{ji}^*} \right) |u_i|^2 \\ &\quad - \sum_{i=1}^n M_i \left( \sum_{j=1}^n |a_{ij}| |s_j(0)| \right. \\ &\quad \left. + \sum_{k=0}^K \sum_{j=1}^n |t_{ij}^{(k)}| |s_j(0)| + |I_i| \right) |u_i|. \end{aligned}$$

So, we obtain

$$\begin{aligned} &\int_{\Omega} \left( \sum_{i=1}^n M_i |u_i| |H_i(u, \lambda)| \right) dx \\ &\geq \int_{\Omega} \left[ \sum_{i=1}^n \left( M_i \alpha_i - \frac{1}{2} \sum_{j=1}^n M_i |a_{ij}|^{2p_{ij}} \beta_j^{2q_{ij}} \right. \right. \\ &\quad \left. - \frac{1}{2} \sum_{j=1}^n M_j |a_{ji}|^{2-2p_{ji}} \beta_i^{2-2q_{ji}} - \frac{1}{2} \sum_{k=0}^K \sum_{j=1}^n M_i |t_{ij}^{(k)}|^{2p_{ij}^*} \beta_j^{2q_{ij}^*} \right. \\ &\quad \left. - \frac{1}{2} \sum_{k=0}^K \sum_{j=1}^n M_j |t_{ji}^{(k)}|^{2-2p_{ji}^*} \beta_i^{2-2q_{ji}^*} \right) |u_i|^2 \\ &\quad \left. - \sum_{i=1}^n M_i \left( \sum_{j=1}^n |a_{ij}| |s_j(0)| + \sum_{k=0}^K \sum_{j=1}^n |t_{ij}^{(k)}| |s_j(0)| \right. \right. \\ &\quad \left. \left. + |I_i| \right) |u_i| \right] dx \\ &\geq \delta_0 \|u(t)\|_2^2 - L \|u(t)\|_2, \end{aligned}$$

where

$$\delta_0 = \min_i \left\{ M_i \alpha_i - \frac{1}{2} \sum_{j=1}^n M_i |a_{ij}|^{2p_{ij}} \beta_j^{2q_{ij}} - \frac{1}{2} \sum_{j=1}^n M_j |a_{ji}|^{2-2p_{ji}} \beta_i^{2-2q_{ji}} - \frac{1}{2} \sum_{k=0}^K \sum_{j=1}^n M_i |t_{ij}^{(k)}|^{2p_{ij}^*} \beta_j^{2q_{ij}^*} - \frac{1}{2} \sum_{k=0}^K \sum_{j=1}^n M_j |t_{ji}^{(k)}|^{2-2p_{ji}^*} \beta_i^{2-2q_{ji}^*} \right\} > 0,$$

$$L = \max_i \{ \int_{\Omega} M_i (\sum_{j=1}^n |a_{ij}| |s_j(0)| + \sum_{k=0}^K \sum_{j=1}^n |t_{ij}^{(k)}| |s_j(0)| + |I_i|) dx \}.$$

Take  $U(R_0) = \{U \in R^n \mid \|u(t)\|_2 < R_0 = (L+1)/\delta_0\}$ . For any  $u \in \partial U(R_0)$ , we have

$$\int_{\Omega} \left( \sum_{i=1}^n M_i |u_i| |H_i(u, \lambda)| \right) dx \geq \delta_0 \|u(t)\|_2 \left( \|u(t)\|_2 - \frac{L}{\delta_0} \right) > 0, \quad \forall \lambda \in [0, 1],$$

i.e.,

$$H(u, \lambda) \neq 0, \quad \forall u \in \partial U(R_0), \quad \lambda \in [0, 1].$$

Following from Lemma 2, we obtain

$$\begin{aligned} \deg(h(u), U(R_0), 0) &= \deg(H(u, 1), U(R_0), 0) \\ &= \deg(H(u, 0), U(R_0), 0) \\ &= 1. \end{aligned}$$

It follows from Lemma 1 that there exist at least an  $u^* \in U(R^0)$ , such that  $h(u^*) = 0$ , i.e., system (1) has at least an equilibrium point  $u^*$ .

Next, we prove the uniqueness of the equilibrium point. Suppose that  $\bar{u}^*$  is also an equilibrium point of system (1), then

$$\begin{aligned} b_i(\bar{u}_i^*) - \sum_{j=1}^n a_{ij} s_j(\bar{u}_j^*) - \sum_{k=0}^K \sum_{j=1}^n t_{ij}^{(k)} s_j(\bar{u}_j^*) + I_i &= 0, \\ i &= 1, \dots, n. \end{aligned} \quad (8)$$

Thus

$$\begin{aligned} b_i(u_i^*) - b_i(\bar{u}_i^*) &= \left( \sum_{j=1}^n a_{ij} s_j(u_j^*) - \sum_{j=1}^n a_{ij} s_j(\bar{u}_j^*) \right) \\ &\quad + \left( \sum_{k=0}^K \sum_{j=1}^n t_{ij}^{(k)} s_j(u_j^*) - \sum_{k=0}^K \sum_{j=1}^n t_{ij}^{(k)} s_j(\bar{u}_j^*) \right). \end{aligned} \quad (9)$$

From (H2) and (H3), we obtain

$$\sum_{i=1}^n \alpha_i M_i |u_i^* - \bar{u}_i^*|^2 \leq \sum_{i=1}^n \left( \frac{1}{2} \sum_{j=1}^n M_i |a_{ij}|^{2p_{ij}} \beta_j^{2q_{ij}} \right.$$

$$\left. + \frac{1}{2} \sum_{j=1}^n M_j |a_{ji}|^{2-2p_{ji}} \beta_i^{2-2q_{ji}} + \frac{1}{2} \sum_{k=0}^K \sum_{j=1}^n M_i |t_{ij}^{(k)}|^{2p_{ij}^*} \beta_j^{2q_{ij}^*} + \frac{1}{2} \sum_{k=0}^K \sum_{j=1}^n M_j |t_{ji}^{(k)}|^{2-2p_{ji}^*} \beta_i^{2-2q_{ji}^*} \right) |u_i^* - \bar{u}_i^*|^2.$$

That is

$$\begin{aligned} \sum_{i=1}^n \left( -M_i \alpha_i + \frac{1}{2} \sum_{j=1}^n M_i |a_{ij}|^{2p_{ij}} \beta_j^{2q_{ij}} + \frac{1}{2} \sum_{j=1}^n M_j |a_{ji}|^{2-2p_{ji}} \beta_i^{2-2q_{ji}} + \frac{1}{2} \sum_{k=0}^K \sum_{j=1}^n M_i |t_{ij}^{(k)}|^{2p_{ij}^*} \beta_j^{2q_{ij}^*} + \frac{1}{2} \sum_{k=0}^K \sum_{j=1}^n M_j |t_{ji}^{(k)}|^{2-2p_{ji}^*} \beta_i^{2-2q_{ji}^*} \right) |u_i^* - \bar{u}_i^*|^2 \leq 0, \end{aligned}$$

which implies  $u_i^* = \bar{u}_i^*$  ( $i = 1, \dots, n$ ). Hence system (1) has a unique equilibrium point  $u^*$ . This completes the proof.  $\square$

Denoting  $y_i(t, x) = u_i(t, x) - u_i^*$ ,  $i = 1, \dots, n$ , system (1) can be rewritten as follows:

$$\begin{cases} \frac{\partial y_i(t, x)}{\partial t} = \sum_{l=1}^m \frac{\partial}{\partial x_l} \left( D_{il} \frac{\partial y_i(t, x)}{\partial x_l} \right) - a_i(y_i(t, x)) \\ \quad \times \left[ b_i(u_i(t, x)) - b_i(u_i^*) - \sum_{j=1}^n a_{ij} s_j(u_j(t, x)) \right. \\ \quad + \sum_{j=1}^n a_{ij} s_j(u_j^*) - \sum_{k=0}^K \sum_{j=1}^n t_{ij}^{(k)} s_j(u_j(t - \tau_k, x)) \\ \quad \left. + \sum_{k=0}^K \sum_{j=1}^n t_{ij}^{(k)} s_j(u_j^*) \right], \quad t \geq 0, \quad x \in \Omega \\ y_i(t, x) = \psi_i(t, x), \quad -\tau \leq t \leq 0, \quad x \in \Omega \\ \frac{\partial y_i(t, x)}{\partial n} = 0, \quad x \in \partial \Omega, \end{cases} \quad (10)$$

in which  $i = 1, \dots, n$ ,  $\psi_i(t, x) = \varphi_i(t, x) - u_i^*$ .  $y(t, x) = (y_1(t, x), \dots, y_n(t, x))^T$ ,  $\psi(t, x) = (\psi_1(t, x), \dots, \psi_n(t, x))^T$ . Obviously,  $u^*$  for system (1) is GES if and only if the equilibrium point  $O$  of system (10) is GES. Thus in the following, we only consider GES of the equilibrium point  $O$  for system (10).

**Theorem 2.** If (H1)–(H4) holds, then the equilibrium point  $O$  of system (10) is GES.

**Proof.** From (H4), there exists a sufficiently small constant  $0 < \lambda < \min_i \{m_i \alpha_i\}$ , such that

$$\begin{aligned} 2\lambda - 2m_i \alpha_i + \sum_{j=1}^n M_i |a_{ij}|^{2p_{ij}} \beta_j^{2q_{ij}} + \sum_{j=1}^n M_j |a_{ji}|^{2-2p_{ji}} \beta_i^{2-2q_{ji}} \\ + \sum_{k=0}^K \sum_{j=1}^n M_i |t_{ij}^{(k)}|^{2p_{ij}^*} \beta_j^{2q_{ij}^*} + \sum_{k=0}^K \sum_{j=1}^n M_j |t_{ji}^{(k)}|^{2-2p_{ji}^*} \beta_i^{2-2q_{ji}^*} e^{2\lambda \tau} \\ \geq 0. \end{aligned}$$

Taking Lyapunov functional as follows:

$$V(t) = \int_{\Omega} \sum_{i=1}^n \left( |y_i(t, x)|^2 e^{2\lambda t} + \sum_{k=0}^K \sum_{j=1}^n M_i |t_{ij}^{(k)}|^{2-2p_{ij}^*} \beta_j^{2-2q_{ij}^*} \times \int_{t-\tau_k}^t |y_j(s, x)|^2 e^{2\lambda(s+\tau_k)} ds \right) dx. \quad (11)$$

Calculating the rate of change of  $V(t)$  along (10), we have

$$\begin{aligned} D^+ V(t) &= \int_{\Omega} \sum_{i=1}^n \left( 2\lambda |y_i(t, x)|^2 e^{2\lambda t} + 2|y_i(t, x)| e^{2\lambda t} \text{sign}(y_i(t, x)) \dot{y}_i(t, x) \right. \\ &\quad + \sum_{k=0}^K \sum_{j=1}^n M_i |t_{ij}^{(k)}|^{2-2p_{ij}^*} \beta_j^{2-2q_{ij}^*} |y_j(t, x)|^2 e^{2\lambda(t+\tau_k)} \\ &\quad - \sum_{k=0}^K \sum_{j=1}^n M_i |t_{ij}^{(k)}|^{2-2p_{ij}^*} \beta_j^{2-2q_{ij}^*} \\ &\quad \times |y_j(t - \tau_k, x)|^2 e^{2\lambda t} \Big) dx \\ &= \int_{\Omega} \sum_{i=1}^n \left[ 2\lambda |y_i(t, x)|^2 e^{2\lambda t} + 2|y_i(t, x)| e^{2\lambda t} \text{sign}(y_i) \right. \\ &\quad \times \left( \sum_{l=1}^m \frac{\partial}{\partial x_l} \left( D_{il} \frac{\partial y_i(t, x)}{\partial x_l} \right) - a_i(y_i(t, x)) \right) \\ &\quad \times \left[ b_i(u_i(t, x)) - b_i(u_i^*) - a_{ij} s_j(u_j(t, x)) + a_{ij} s_j(u_j^*) \right. \\ &\quad - \sum_{k=0}^K \sum_{j=1}^n t_{ij}^{(k)} s_j(u_j(t - \tau_k, x)) + \sum_{k=0}^K \sum_{j=1}^n t_{ij}^{(k)} s_j(u_j^*) \Big] \\ &\quad + \sum_{k=0}^K \sum_{j=1}^n M_i |t_{ij}^{(k)}|^{2-2p_{ij}^*} \beta_j^{2-2q_{ij}^*} |y_j(t, x)|^2 e^{2\lambda(t+\tau_k)} \\ &\quad - \sum_{k=0}^K \sum_{j=1}^n M_i |t_{ij}^{(k)}|^{2-2p_{ij}^*} \beta_j^{2-2q_{ij}^*} \\ &\quad \times |y_j(t - \tau_k, x)|^2 e^{2\lambda t} \Big] dx. \end{aligned}$$

From the boundary condition, we have

$$\begin{aligned} &\int_{\Omega} \sum_{l=1}^m \frac{\partial}{\partial x_l} \left( D_{il} \frac{\partial y_i(t, x)}{\partial x_l} \right) |y_i(t, x)| dx \\ &= \sum_{l=1}^m \int_{\Omega} |y_i(t, x)| \frac{\partial}{\partial x_l} \left( D_{il} \frac{\partial y_i(t, x)}{\partial x_l} \right) dx \\ &= \sum_{l=1}^m \left[ \int_{\Omega} \frac{\partial}{\partial x_l} \left( |y_i(t, x)| D_{il} \frac{\partial y_i(t, x)}{\partial x_l} \right) dx \right. \\ &\quad \left. - \int_{\Omega} \frac{\partial y_i(t, x)}{\partial x_l} D_{il} \frac{\partial y_i(t, x)}{\partial x_l} dx \right] \end{aligned}$$

$$= - \sum_{l=1}^m \int_{\Omega} D_{il} \left( \frac{\partial y_i(t, x)}{\partial x_l} \right)^2 |y_i(t, x)| dx.$$

Combining (H1)–(H4) with  $2ab \leq a^2 + b^2$ , we obtain

$$\begin{aligned} D^+ V(t) &\leq \int_{\Omega} \sum_{i=1}^n \left[ 2\lambda |y_i(t, x)|^2 e^{2\lambda t} + 2e^{2\lambda t} \left( -m_i \alpha_i |y_i(t, x)|^2 \right. \right. \\ &\quad + \sum_{j=1}^n |a_{ij}| \beta_j |y_i(t, x)| |y_j(t, x)| \\ &\quad + \sum_{k=0}^K \sum_{j=1}^n M_i |t_{ij}^{(k)}| \beta_j |y_i(t, x)| |y_j(t - \tau_k, x)| \Big) \\ &\quad + \sum_{k=0}^K \sum_{j=1}^n M_i |t_{ij}^{(k)}|^{2-2p_{ij}^*} \beta_j^{2-2q_{ij}^*} |y_j(t, x)|^2 e^{2\lambda(t+\tau_k)} \\ &\quad - \sum_{k=0}^K \sum_{j=1}^n M_i |t_{ij}^{(k)}|^{2-2p_{ij}^*} \beta_j^{2-2q_{ij}^*} |y_j(t - \tau_k, x)|^2 e^{2\lambda t} \Big] dx \\ &\leq \int_{\Omega} \sum_{i=1}^n \left[ 2\lambda |y_i(t, x)|^2 e^{2\lambda t} + 2e^{2\lambda t} \left( -m_i \alpha_i |y_i(t, x)|^2 \right. \right. \\ &\quad + \sum_{j=1}^n M_i |a_{ij}|^{p_{ij}} \beta_j^{q_{ij}} |y_i(t, x)| |a_{ij}|^{1-p_{ij}} \beta_j^{1-q_{ij}} |y_j(t, x)| \\ &\quad + \sum_{k=0}^K \sum_{j=1}^n M_i |t_{ij}^{(k)}|^{p_{ij}} \beta_j^{q_{ij}} |y_i(t, x)| |t_{ij}^{(k)}|^{1-p_{ij}} \\ &\quad \times \beta_j^{1-q_{ij}} |y_j(t - \tau_k, x)| \Big) \\ &\quad + \sum_{k=0}^K \sum_{j=1}^n M_i |t_{ij}^{(k)}|^{2-2p_{ij}^*} \beta_j^{2-2q_{ij}^*} |y_j(t, x)|^2 e^{2\lambda(t+\tau_k)} \\ &\quad - \sum_{k=0}^K \sum_{j=1}^n M_i |t_{ij}^{(k)}|^{2-2p_{ij}^*} \beta_j^{2-2q_{ij}^*} |y_j(t - \tau_k, x)|^2 e^{2\lambda t} \Big] dx \\ &\leq \int_{\Omega} \sum_{i=1}^n \left[ 2\lambda |y_i(t, x)|^2 e^{2\lambda t} + e^{2\lambda t} (-2m_i \alpha_i |y_i(t, x)|^2 \right. \\ &\quad + \sum_{j=1}^n M_i |a_{ij}|^{2p_{ij}} \beta_j^{2q_{ij}} |y_i(t, x)|^2 \\ &\quad + \sum_{j=1}^n M_i |a_{ij}|^{2-2p_{ij}} \beta_j^{2-2q_{ij}} |y_j(t, x)|^2 \\ &\quad + \sum_{k=0}^K \sum_{j=1}^n M_i |t_{ij}^{(k)}|^{2p_{ij}} \beta_j^{2q_{ij}} |y_i(t, x)|^2 \\ &\quad + \sum_{k=0}^K \sum_{j=1}^n M_i |t_{ij}^{(k)}|^{2-2p_{ij}^*} \beta_j^{2-2q_{ij}^*} |y_j(t - \tau_k, x)|^2 \Big) \\ &\quad + \sum_{k=0}^K \sum_{j=1}^n M_i |t_{ij}^{(k)}|^{2-2p_{ij}^*} \beta_j^{2-2q_{ij}^*} |y_j(t, x)|^2 e^{2\lambda(t+\tau_k)} \\ &\quad - \sum_{k=0}^K \sum_{j=1}^n M_i |t_{ij}^{(k)}|^{2-2p_{ij}^*} \beta_j^{2-2q_{ij}^*} |y_j(t - \tau_k, x)|^2 e^{2\lambda t} \Big] dx \end{aligned}$$

$$\begin{aligned}
&\leq \sum_{i=1}^n \left[ 2\lambda - 2m_i\alpha_i + \sum_{j=1}^n M_i |a_{ij}|^{2p_{ij}} \beta_j^{2q_{ij}} \right. \\
&\quad + \sum_{j=1}^n M_j |a_{ji}|^{2-2p_{ji}} \beta_i^{2-2q_{ji}} + \sum_{k=0}^K \sum_{j=1}^n M_i |t_{ij}^{(k)}|^{2p_{ij}^*} \beta_j^{2q_{ij}^*} \\
&\quad \left. + \sum_{k=0}^K \sum_{j=1}^n M_j |t_{ji}^{(k)}|^{2-2p_{ji}^*} \beta_i^{2-2q_{ji}^*} e^{2\lambda\tau} \right] \|y_i(t)\|_2^2 e^{2\lambda t} \\
&\leq 0,
\end{aligned} \tag{12}$$

which implies

$$V(t) \leq V(0).$$

By Eq.(11) we have

$$\begin{aligned}
V(0) &= \int_{\Omega} \sum_{i=1}^n \left( |\psi_i(t, x)|^2 + \sum_{k=0}^K \sum_{j=1}^n M_i |t_{ij}^{(k)}|^{2-2p_{ij}^*} \beta_j^{2-2q_{ij}^*} \right. \\
&\quad \times \int_{-\tau_k}^0 |y_j(s, x)|^2 e^{2\lambda(s+\tau_k)} ds \Big) dx \\
&\leq \sum_{i=1}^n \left( \|\psi_i(t)\|_2^2 + \sum_{k=0}^K \sum_{j=1}^n M_i |t_{ij}^{(k)}|^{2-2p_{ij}^*} \beta_j^{2-2q_{ij}^*} \right. \\
&\quad \times \int_{-\tau_k}^0 \|y_j(s)\|_2^2 e^{2\lambda(s+\tau_k)} ds \Big) \\
&\leq \max_i \left\{ 1 + \sum_{k=0}^K \sum_{j=1}^n |t_{ij}^{(k)}|^{2-2p_{ij}^*} \beta_j^{2-2q_{ij}^*} (e^{\tau} - 1) \right\} \|\psi\|_2^2,
\end{aligned}$$

and

$$V(t) \geq \|y(t)\|_2^2 e^{2\lambda t}.$$

Hence

$$\|y(t)\|_2^2 e^{2\lambda t} \leq \max_i \left\{ 1 + \sum_{k=0}^K \sum_{j=1}^n |t_{ij}^{(k)}|^{2-2p_{ij}^*} \beta_j^{2-2q_{ij}^*} (e^{\tau} - 1) \right\} \|\psi\|_2^2,$$

which leads to

$$\|y(t)\|_2 \leq \gamma \|\psi\|_2 e^{-\lambda t},$$

where

$$\gamma = \left( \max_i \left\{ 1 + \sum_{k=0}^K \sum_{j=1}^n |t_{ij}^{(k)}|^{2-2p_{ij}^*} \beta_j^{2-2q_{ij}^*} (e^{\tau} - 1) \right\} \right)^{1/2}.$$

This completes the proof.  $\square$

**Corollary 1.** System (1) has a unique equilibrium point which is GES, if the conditions (H1)–(H3). Assume furthermore that one of the following conditions hold:

(H5)

$$-2m_i\alpha_i + \sum_{j=1}^n M_i |a_{ij}|^2 \beta_j^2 + \sum_{k=0}^K \sum_{j=1}^n M_i |t_{ij}^{(k)}|^2 \beta_j^2 < 0.$$

(H6)

$$\begin{aligned}
&-2m_i\alpha_i + \sum_{j=1}^n M_i |a_{ij}| \beta_j + \sum_{j=1}^n M_j |a_{ji}| \beta_i \\
&\quad + \sum_{k=0}^K \sum_{j=1}^n M_i |t_{ij}^{(k)}| \beta_j + \sum_{k=0}^K \sum_{j=1}^n M_j |t_{ji}^{(k)}| \beta_i < 0.
\end{aligned}$$

**Proof.** In (H4), let  $p_{ij} = q_{ij} = p_{ij}^* = q_{ij}^* = 1$ , then (H4) turns to (H5). Furthermore, we suppose that  $p_{ij} = q_{ij} = p_{ij}^* = q_{ij}^* = \frac{1}{2}$ , then (H4) turns to (H6). By Theorems 1 and 2, system (1) has a unique equilibrium point which is GES.

**Remark 1.** When  $D_{ik} \equiv 0$ , then system (1) becomes the system analyzed in [19,2,16,14,9]. It is worth noting that, in the paper, we did not need  $s_j$  is bounded. Thus, we improve the results in Refs. [19,2,16,14,9].

**Remark 2.** From Theorems 1 and 2, we see if reaction–diffusion terms satisfy a weaker condition  $D_{il} \geq 0$ , then the effects for the existence and GES of the equilibrium point just come from the networks parameters, the stability is completely expressed by the relations of these parameters.

**Remark 3.** Although the assertions of exponential stability in Theorems 1 and 2 are independent of the delays, the convergence rate  $\lambda$  do depend on the delays  $\tau_k$ .

#### 4. Example

**Example.** Consider Cohen–Grossberg neural networks with delays and reaction–diffusion terms

$$\begin{cases} \frac{\partial u_1(t, x)}{\partial t} = \frac{\partial}{\partial x} \left( D_{11} \frac{\partial u_1(t, x)}{\partial x} \right) - a_1(u_1(t, x)) \\ \quad \times \left[ b_1(u_1(t, x)) - \sum_{j=1}^2 a_{1j} s_j(u_j) \right. \\ \quad \left. - \sum_{j=1}^2 \sum_{k=0}^1 t_{1j}^{(k)} s_j(u_j(t - \tau_k, x)) \right] \\ \frac{\partial u_2(t, x)}{\partial t} = \frac{\partial}{\partial x} \left( D_{21} \frac{\partial u_2(t, x)}{\partial x} \right) - a_2(u_2(t, x)) \\ \quad \times \left[ b_2(u_2(t, x)) - \sum_{j=1}^2 a_{2j} s_j(u_j) \right. \\ \quad \left. - \sum_{j=1}^2 \sum_{k=0}^1 t_{2j}^{(k)} s_j(u_j(t - \tau_k, x)) \right]. \end{cases} \tag{13}$$

Let  $D_{11} > 0, D_{21} > 0$ ,  $a_i = 4 + \sin u_i$ ,  $b_i(u_i) = 4u_i$ ,  $s_1(u_1) = \arctan u_1$ ,  $s_2 = u_2$ . Clearly,  $a_i$  satisfies (H1) with  $m_i = 3$ ,  $M_i = 5$ ,  $s_j$  satisfies (H2) with  $\beta_j = 1$ ,  $b_i$  satisfies (H3) with  $\alpha_i = 4$ ,  $i, j = 1, 2$ . Moreover, we choose  $a_{11} = \frac{1}{12}$ ,  $a_{12} = \frac{1}{12}$ ,  $a_{21} = \frac{1}{6}$ ,  $a_{22} = \frac{1}{24}$ ,  $t_{11}^{(0)} = \frac{1}{12}$ ,  $t_{12}^{(0)} = \frac{1}{6}$ ,  $t_{11}^{(1)} = \frac{1}{12}$ ,  $t_{12}^{(1)} = \frac{1}{4}$ ,  $t_{21}^{(0)} = \frac{1}{6}$ ,  $t_{22}^{(0)} = \frac{1}{24}$ ,  $t_{21}^{(1)} = \frac{1}{12}$ ,  $t_{22}^{(1)} = \frac{1}{24}$ . By simple calculation, we show that (H6) holds. It follows from Theorems 1 and 2

that system (13) has a unique equilibrium point  $(0,0)^T$  which is GES (Figs. 1–6).

## 5. Conclusions

In this paper, the dynamics of Cohen–Grossberg neural networks model with delays and reaction–diffusion is studied. By employing homotopic mapping theory and constructing Lyapunov functional method, some sufficient conditions have been obtained which guarantee the model to be GES. The given algebra conditions are useful in design and applications of reaction–diffusion Cohen–Grossberg neural networks. Moreover, our methods in the paper may be extended for more complex networks.




Fig. 3. The cube graph of  $u_1$  and  $x$  about time in the system (13) with  $\tau_k = 5$ .




Fig. 1. The trajectory of  $u_1$  versus time in the system (13) with  $\tau_k = 5$ .




Fig. 4. The cube graph of  $u_2$ ,  $x$  and time in the system (13) with  $\tau_k = 5$ .




Fig. 2. The trajectory of  $u_2$  versus time in the system (13) with  $\tau_k = 5$ .




Fig. 5. The cube graph of  $u_1$  and  $u_2$  about time in the system (13) with  $\tau_k = 5$ .






Fig. 6. The cube graph of  $u_1$  and  $u_2$  about  $x$  in the system (13) with  $\tau_k = 5$ .

### Acknowledgment

The authors would like to thank the Editor and the reviewers for their helpful comments and constructive suggestions, which have been very useful for improving the presentation of this paper.

### References

- [1] J. Cao, An estimation of the domain of attraction and convergence rate for Hopfield continuous feedback neural networks, *Phys. Lett. A* 325 (2004) 370–374.
- [2] J. Cao, L. Liang, Boundedness and stability for Cohen–Grossberg neural network with time-varying delays, *J. Math. Anal. Appl.* 296 (2004) 665–685.
- [3] J. Cao, J. Wang, Globally exponentially stability and periodicity of recurrent neural networks with time delays, *IEEE Trans. Circuits Syst.—I* 52 (2005) 920–931.
- [4] M. Cohen, S. Grossberg, *IEEE Trans. Syst. Man Cybernet. SMC-13* (1983) 15.
- [5] J. Cronin, *Fixed Point and Topological Degree in Nonlinear Analysis*, American Mathematical Society, Providence, RI, 1964.
- [6] A. Hasting, Global stability in Lotka–Volterra systems with diffusion, *J. Math. Biol.* 6 (1978) 163–168.
- [7] J. Liang, J. Cao, Global exponential stability of reaction–diffusion recurrent neural networks with time-varying delays, *Phys. Lett. A* 314 (2003) 434–442.
- [8] X. Liao, C. Li, Stability in Gilpin–Ayala competition models with diffusion, *Nonlinear Anal. TMA* 18 (1997) 1751–1758.
- [9] X. Liao, C. Li, K. Wong, Criteria for exponential stability of Cohen–Grossberg neural networks, *Neural Networks* 17 (2004) 1401–1414.
- [10] F. Rothe, Convergence to the equilibrium state in the Volterra–Lotka diffusion equations, *J. Math. Biol.* 3 (1976) 319–324.
- [11] Q. Song, J. Cao, Global exponential stability and existence of periodic solutions in BAM networks with delays and reaction–diffusion terms, *Chaos Solitons Fractals* 23 (2005) 421–430.
- [12] Q. Song, Z. Zhao, Y. Li, Global exponential stability of BAM neural networks with distributed delays and reaction–diffusion terms, *Phys. Lett. A* 335 (2005) 213–225.
- [13] L. Wang, D. Xu, Global exponential stability of reaction–diffusion Hopfield neural networks with variable delays, *Sci. China Ser. E* 33 (2003) 488–495 (in Chinese).
- [14] W. Xiong, J. Cao, Global exponential stability of discrete-time Cohen–Grossberg neural networks, *Neurocomputing* 64 (2005) 433–446.
- [15] D. Xu, H. Zhao, Global dynamics of Hopfield neural networks involving variable delays, *Comput. Math. Appl.* 42 (2001) 39–45.
- [16] K. Yuan, J. Cao, Global exponential stability of Cohen–Grossberg neural networks with multiple time-varying delays, *Lecture Notes in Computer Science*, ISSN 2004, vol. 3173, 2004, pp. 78–83.
- [17] J. Zhang, Global stability analysis in delayed Hopfield neural network models, *Neural Networks* 13 (2000) 745–753.
- [18] J. Zhang, Global stability analysis in Hopfield neural networks, *Appl. Math. Lett.* 16 (2003) 925–931.
- [19] J. Zhang, Y. Suda, H. Komine, Global exponential stability of Cohen–Grossberg neural networks with variable delays, *Phys. Lett. A* 338 (2005) 44–50.
- [20] H. Zhao, Existence of periodic oscillatory solution of reaction–diffusion neural networks with delays, *Phys. Lett. A* 343 (2005) 372–383.
- [21] H. Zhao, G. Wang, Global exponential stability and periodicity of cellular neural networks with variable delays, *Phys. Lett. A* 336 (2005) 331–341.



**Hongyong Zhao** received the Ph.D. degree from Sichuan University, Chengdu, China, and the Post-Doctoral Fellow in the Department of Mathematics at Nanjing University, Nanjing, China.

He was with Department of Mathematics at Nanjing University of Aeronautics and Astronautics, Nanjing, China. He is currently a Professor of Nanjing University of Aeronautics and Astronautics, Nanjing, China. He also is the author or coauthor of more than 40 journal papers. His research interests include nonlinear dynamic systems, neural networks, control theory, and applied mathematics.



**Kunlun Wang** is with Department of Computer Science and Technology in Hefei University, Hefei, China. He is currently a professor of Department of Computer Science and Technology in Hefei University, Hefei, China. He also is the author or coauthor of more than 30 journal papers. His research interests include Artificial neural networks, Intelligent control and Speech Recognition.