

Mathematical Foundations of Data Sciences



Gabriel Peyré
CNRS & DMA
École Normale Supérieure
gabriel.peyre@ens.fr
www.gpeyre.com
www.numerical-tours.com

November 9, 2017

Chapter 14

Theory of Sparse Regularization

We now apply the basics elements of convex analysis from the previous chapter to perform a theoretical analysis of the properties of the Lasso, in particular its performances to recover sparse vectors.

14.1 Existence and Uniqueness

14.1.1 Existence

We consider problems (11.10) and (11.11), that we rewrite here as

$$\min_{x \in \mathbb{R}^N} f_\lambda(x) \stackrel{\text{def.}}{=} \frac{1}{2\lambda} \|y - Ax\|^2 + \lambda \|x\|_1 \quad (\mathcal{P}_\lambda(y))$$

and its limit as $\lambda \rightarrow 0$

$$\min_{Ax=y} \|x\|_1 = \min_x f_0(x) \stackrel{\text{def.}}{=} \iota_{\mathcal{L}_y}(x) + \|x\|_1. \quad (\mathcal{P}_0(y))$$

where $A \in \mathbb{R}^{P \times N}$, and $\mathcal{L}_y \stackrel{\text{def.}}{=} \{x \in \mathbb{R}^N ; Ax = y\}$.

We recall that the setup is that one observe noise measures

$$y = Ax_0 + w$$

and we would like conditions to ensure for x_0 to solution to $(\mathcal{P}_0(Ax_0))$ (i.e. when $w = 0$) and to be close (in some sense to be defined, and in some proportion to the noise level $\|w\|$) to the solutions of $(\mathcal{P}_0(y = Ax_0 + w))$ when λ is wisely chosen as a function of $\|w\|$.

First let us note that since $(\mathcal{P}_\lambda(y))$ is unconstrained and coercive (because $\|\cdot\|_1$ is), this problem always has solutions. Since A might have a kernel and $\|\cdot\|_1$ is not strongly convex, it might have non-unique solutions. If $y \in \text{Im}(A)$, the constraint set of $(\mathcal{P}_0(y))$ is non-empty, and it also has solutions, which might fail to be unique.

Figure 14.1 gives the intuition of the theory that will be developed in this chapter, regarding the exact or approximated recovery of sparse vectors x_0 , and the need for a careful selection of the λ parameter.

14.1.2 Polytope Projection for the Constraint Problem

The following proposition gives a geometric description of those vectors which are recovered by ℓ^1 minimization when there is no noise.

Proposition 43. *We denote $B \stackrel{\text{def.}}{=} \{x \in \mathbb{R}^N ; \|x\|_1 \leq 1\}$. Then*

$$x_0 \text{ is a solution to } \mathcal{P}_0(Ax_0) \iff A \frac{x_0}{\|x_0\|_1} \in \partial(AB) \quad (14.1)$$

where “ ∂ ” denoted the boundary and $AB = \{Ax ; x \in B\}$.




Figure 14.1: Display of the evolution $\lambda \mapsto x_\lambda$ of the solutions of $(\mathcal{P}_\lambda(y))$.

Proof. We first prove “ \Rightarrow ”. We suppose that x_0 is not a solution, and aim at showing that $A \frac{x_0}{\|x_0\|_1} \in \text{int}(AB_\rho)$. Since it is not a solution, there exists z such that $Ax_0 = Az$ and $\|z\|_1 = (1 - \delta)\|x_0\|_1$ with $\delta > 0$. Then for any displacement $h = A\varepsilon \in \text{Im}(A)$, one has $Ax_0 + h = A(z + \varepsilon)$ and

$$\|z + \varepsilon\|_1 \leq \|z\|_1 + \|\Phi^+ h\| \leq (1 - \delta)\|x_0\|_1 + \|\Phi^+\|_{1,1} \|h\|_1 < \frac{\delta}{\|A^+\|_{1,1}} \|x_0\|_1.$$

This means that choosing $\|h\|_1 < \frac{\delta}{\|A^+\|_{1,1}} \|x_0\|_1$ implies that $A \frac{x_0}{\|x_0\|_1} \in \text{int}(AB)$.

We now prove “ \Leftarrow ”. We suppose that $A \frac{x_0}{\|x_0\|_1} \in \text{int}(AB)$. Then there exists z such that $Ax_0 = (1 - \delta)Az$ and $\|z\|_1 < \|x_0\|_1$. This implies $\|(1 - \delta)z\|_1 < \|x_0\|_1$ so that $(1 - \delta)z$ is better than x_0 which is thus not a solution. \square

This results state that “friendly” identifiable vectors (those recovered by ℓ^1) are those who gets projected by A on the boundary of the polytope $\|x_0\|_1 AB$. Intuitively, if P is small in comparison to N , then this projected polytope is small, and most vector will failed to be reconstructed by solving ℓ^1 minimization. This also suggests why using random projections as in Chapter 15, because somehow they results in a low distortion embedding of the ℓ^1 ball from \mathbb{R}^N to \mathbb{R}^P .

Note that if x_0 is identifiable, so is λx_0 for ρx_0 for $\rho > 0$, and in fact, since the recovery condition only depends on the geometry of the faces of B , the obtained condition (14.1) only depends on $\text{sign}(x_0)$. We denote $\mathcal{A} : y \mapsto x^*$ the map from y to a solution of $(\mathcal{P}_0(y))$, which we assume is unique for simplicity of exposition. Condition (14.1) thus shows that A and \mathcal{A} are inverse bijection on a family of cones $\mathcal{C}_s = \{x ; \text{sign}(x) = s\}$ and $A\mathcal{C}_s$ for certain “friendly” sign patterns s . These cones $A\mathcal{C}_s$ form a partition of the image space \mathbb{R}^P . Assuming for simplicity that the columns $(a_j)_j$ of A have unit norm, for $P = 3$, the interaction of these $A\mathcal{C}_s$ with the unit sphere of \mathbb{R}^3 for a so-called Delaunay triangulation of the sphere (this construction extends to higher dimension by replacing triangle by simplexes). Such Delaunay triangulation is characterized by the




Figure 14.2: Display of the action of the linear map A on the ℓ^1 ball B , and of the inverse non-linear map \mathcal{A} defined by the solution of $(\mathcal{P}_0(y))$.

empty spherical cap property (each circumcircle associated to a triangle should not contains any columns vector a_j of the matrix). Figure 14.2 illustrate these conclusions in \mathbb{R}^2 and \mathbb{R}^3 .

14.1.3 Optimality Conditions

In the following, given an index set $I \subset \{1, \dots, N\}$, denoting $A = (a_i)_{i=1}^N$ the columns of A , we denote $A_I \stackrel{\text{def.}}{=} (a_i)_{i \in I} \in \mathbb{R}^{P \times |I|}$ the extracted sub-matrix. Similarly, for $x \in \mathbb{R}^N$, we denote $x_I \stackrel{\text{def.}}{=} (x_i)_{i \in I} \in \mathbb{R}^{|I|}$.

The following proposition rephrases the first order optimality conditions in a handy way.

Proposition 44. x_λ is a solution to $(\mathcal{P}_\lambda(y))$ for $\lambda > 0$ if and only if

$$\eta_{\lambda, I} = \text{sign}(x_{\lambda, I}) \quad \text{and} \quad \|\eta_{\lambda, I^c}\| \leq \lambda$$

where we define

$$I \stackrel{\text{def.}}{=} \text{supp}(x_\lambda) \stackrel{\text{def.}}{=} \{i ; x_{\lambda, i} \neq 0\}, \quad \text{and} \quad \eta_\lambda \stackrel{\text{def.}}{=} \frac{1}{\lambda} A^*(y - Ax_\lambda). \quad (14.2)$$

Proof. Since $(\mathcal{P}_\lambda(y))$ involves a sum of a smooth and a continuous function, its sub-differential reads

$$\partial f_\lambda(x) = \frac{1}{\lambda} A^*(Ax - y) + \lambda \partial \|\cdot\|_1(x).$$

Thus x_λ is solution to $(\mathcal{P}_\lambda(y))$ if and only if $0 \in \partial f_\lambda(x_\lambda)$, which gives the desired result. \square

The following proposition studies the limit case $\lambda = 0$ and introduces the crucial concept of “dual certificates”, which are the Lagrange multipliers of the constraint \mathcal{L}_y .

Proposition 45. x^* being a solution to $(\mathcal{P}_0(y))$ is equivalent to having $Ax^* = y$ and that

$$\exists \eta \in \mathcal{D}_0(y, x^*) \stackrel{\text{def.}}{=} \text{Im}(A^*) \cap \partial \|\cdot\|_1(x^*). \quad (14.3)$$

Proof. Since $(\mathcal{P}_0(y))$ involves a sum with a continuous function, one can also computes its sub-differential as

$$\partial f_0(x) = \partial \iota_{\mathcal{L}_y}(x) + \partial \|\cdot\|_1(x).$$

If $x \in \mathcal{L}_y$, then $\partial \iota_{\mathcal{L}_y}(x)$ is the linear space orthogonal to \mathcal{L}_y , i.e. $\ker(A)^\perp = \text{Im}(A^*)$. \square

Writing $I = \text{supp}(x^*)$, one thus has

$$\mathcal{D}_0(y, x^*) = \{\eta = A^*p ; \eta_I = \text{sign}(x_I^*), \|\eta\|_\infty \leq 1\}.$$

Although it looks like the definition of $\mathcal{D}_0(y, x^*)$ depends on the choice of a solution x^* , convex duality (studied in the next chapter) shows that it is not the case (it is the same set for all solutions).

14.1.4 Uniqueness

The following proposition shows that the Lasso selects a set of linearly independent regressors (columns of A). This is why this method is also often called “basis pursuit”.

Proposition 46. *For $\lambda \geq 0$, there is always a solution x^* to $(\mathcal{P}_\lambda(y))$ with $I = \text{supp}(x^*)$ such that $\ker(A_I) = \{0\}$*

Proof. Let x be a solution and denote $I = \text{supp}(x)$. If $\ker(A_I) \neq \{0\}$, one selects $h_I \in \ker(A_I)$ and define for $t \in \mathbb{R}$ the vector $x_t \stackrel{\text{def.}}{=} x + th$. We denote t_0 the smallest $|t|$ such that $\text{sign}(x_t) \neq \text{sign}(x)$, i.e. $\text{supp}(x_t)$ is strictly included in I . For $t < t_0$, since $Ax_t = Ax$ and $\text{sign}(x_t) = \text{sign}(x)$, x_t still satisfies the same first order condition as x_0 , and one can apply either Proposition 45 (for $\lambda = 0$) or Proposition 44 (for $\lambda > 0$), so that x_t is a solution of $(\mathcal{P}_\lambda(y))$. Since the minimized function are lower semi continuous, $x_t \rightarrow x_{t_0}$ is still a solution. If $\ker(A_J) \neq \{0\}$ with $J = \text{supp}(x_{t_0})$, one is over, otherwise one can iterate this argument on x_{t_0} in place of x and have a sequence of supports which is strictly decaying in size, so it must terminate. \square

This results in particular that if columns of A_I are not independent, then the solution of $(\mathcal{P}_\lambda(y))$ is necessarily non-unique.

Assuming that x_λ is a solution such that $\ker(A_I) = \{0\}$, then from $(\mathcal{P}_\lambda(y))$, one obtains the following implicit expression for the solution

$$x_{\lambda,I} = A_I^+ y - \lambda(A_I^* A_I)^{-1} \text{sign}(x_{\lambda,I}). \quad (14.4)$$

This expression can be understood as a form of generalized soft thresholding (one retrieve the soft thresholding when $A = \text{Id}_N$).

The following useful lemma shows that while solutions x_λ to $(\mathcal{P}_\lambda(y))$ are not necessarily unique, the associated “predictor” (i.e. denoised version of y) Ax_λ is however always uniquely defined. Note that according to (14.5), one has

$$\Phi x_\lambda = \text{Proj}_{\text{Im}(A_I)} y - \lambda A_I (A_I^* A_I)^{-1} \text{sign}(x_{\lambda,I}). \quad (14.5)$$

so up to a $O(\lambda)$ bias, this predictor is an orthogonal projection on a low dimensional subspace indexed by I .

Lemma 3. *For $\lambda \geq 0$, if (x_1, x_2) are solution to $(\mathcal{P}_\lambda(y))$, then $Ax_1 = Ax_2$.*

Proof. For $\lambda = 0$, this is trivial because $Ax_1 = Ax_2 = y$. Otherwise, let us assume $Ax_1 \neq Ax_2$. Then for $x = (x_1 + x_2)/2$, one has

$$\|x\|_1 \leq \frac{\|x_1\|_1 + \|x_2\|_2}{2} \quad \text{and} \quad \|Ax - y\|^2 < \frac{\|Ax_1 - y\|^2 + \|Ax_2 - y\|^2}{2}$$

where the second inequality follows from the strict convexity of the square. This shows that

$$\frac{1}{2\lambda} \|Ax - y\|^2 + \|x\|_1 < \frac{1}{2\lambda} \|Ax_1 - y\|^2 + \|x_1\|_1,$$

which is a contradiction to the optimality of x_1 . \square

Proposition 47. *Let x_λ be a solution to $(\mathcal{P}_\lambda(y))$ and denote $\eta_\lambda \stackrel{\text{def.}}{=} \frac{1}{\lambda} A^*(y - Ax_\lambda)$. We define the “extended support” as*

$$J \stackrel{\text{def.}}{=} \text{sat}(\eta_\lambda) \stackrel{\text{def.}}{=} \{i ; |\eta_{\lambda,i}| = 1\}.$$

If $\ker(A_J) = \{0\}$ then x_λ is the unique solution of $(\mathcal{P}_\lambda(y))$.

Proof. If \tilde{x}_λ is also a minimizer, then by Lemma 3, $Ax_\lambda = A\tilde{x}_\lambda$, so that in particular they share the same dual certificate

$$\eta_\lambda = \frac{1}{\lambda} A^*(y - Ax_\lambda) = \frac{1}{\lambda} A^*(y - A\tilde{x}_\lambda).$$

The first order condition, Proposition 44, shows that necessarily $\text{supp}(x_\lambda) \subset J$ and $\text{supp}(\tilde{x}_\lambda) \subset J$. Since $A_J x_{\lambda,J} = A_J \tilde{x}_{\lambda,J}$, and since $\ker(A_J) = \{0\}$, one has $x_{\lambda,J} = \tilde{x}_{\lambda,J}$, and thus $x_\lambda = \tilde{x}_\lambda$ because of their supports are included in J . \square

Proposition 48. Let x^* be a solution to $(\mathcal{P}_0(y))$. If there exists $\eta \in \mathcal{D}_0(y, x^*)$ such that $\ker(A_J) = \{0\}$ where $J \stackrel{\text{def.}}{=} \text{sat}(\eta)$ then x^* is the unique solution of $(\mathcal{P}_0(y))$.

Proof. The proof is the same as for Proposition 47, replacing η_λ by η . \square

These propositions can be used to show that if A is drawn according to a distribution having a density over $\mathbb{R}^{p \times n}$, then with probability 1 on the matrix A , the solution to $(\mathcal{P}_\lambda(y))$ is unique. Note that this results is not true if A is non random but y is.

14.1.5 Duality

We now related the first order conditions and “dual certificate” introduced above to the duality theory detailed in Section 12.2. This is not strictly needed to derive the theory of sparse regularization, but this offers an alternative point of view and allows to better grasp the role played by the certificates.

Theorem 41. For any $\lambda \geq 0$ (i.e. including $\lambda = 0$), one has strong duality between $(\mathcal{P}_\lambda(y))$ and

$$\sup_{p \in \mathbb{R}^P} \left\{ \langle y, p \rangle - \frac{\lambda}{2} \|p\|^2 ; \|A^* p\|_\infty \leq 1 \right\}. \quad (14.6)$$

One has for any $\lambda \geq 0$ that (x^*, p^*) are primal and dual solutions if and only if

$$A^* p^* \in \partial \|\cdot\|_1(x^*) \Leftrightarrow (I \subset \text{sat}(A^* p^*) \text{ and } \text{sign}(x_I^*) = A_I^* p^*), \quad (14.7)$$

where we denoted $I = \text{supp}(x^*)$, and furthermore, for $\lambda > 0$,

$$p^* = \frac{y - Ax^*}{\lambda}.$$

while for $\lambda = 0$, $Ax^* = y$.

Proof. There are several ways to derive the same dual. One can for instance directly use the Fenchel-Rockafeller formula (??). But it is instructive to do the computations using Lagrange duality. One can first consider the following re-writing of the primal problem

$$\min_{x \in \mathbb{R}^N} \{f(z) + \|x\|_1 ; Ax = z\} = \min_{x \in \mathbb{R}^N} \sup_{p \in \mathbb{R}^P} \mathcal{L}(x, z, p) \stackrel{\text{def.}}{=} f_\lambda(z) + \|x\|_1 + \langle z - Ax, p \rangle$$

where $f_\lambda(z) \stackrel{\text{def.}}{=} \frac{1}{2\lambda} \|z - y\|^2$ if $\lambda > 0$ and $f(z) = \iota_{\{y\}}(z)$ if $\lambda = 0$. For $\lambda > 0$ since f_λ and $\|\cdot\|_1$ are continuous, strong duality holds. For $\lambda = 0$, since the constraint appearing in f_0 is linear (actually a singleton), strong duality holds also. Thus using Theorem 32, one can exchange the min and the max and obtains

$$\max_{p \in \mathbb{R}^P} (\min_z \langle z, p \rangle + f_\lambda(z)) + (\min_x \|x\|_1 - \langle x, A^* p \rangle) = \max_{p \in \mathbb{R}^P} -f_\lambda^*(-p) - (\|\cdot\|_1)^*(A^* p).$$

Using (30), one has that $(\|\cdot\|_1^* = \iota_{\|\cdot\|_\infty \leq 1})$. For $\lambda > 0$, one has using Proposition 31 that

$$f_\lambda^* = \left(\frac{1}{2\lambda} \|\cdot - y\|^2\right)^* = \frac{1}{\lambda} \left(\frac{1}{2} \|\cdot - y\|^2\right)^*(\lambda \cdot) = \frac{1}{2\lambda} \|\lambda \cdot\|^2 + \langle \cdot, y \rangle$$

which gives the desired dual problem. The first order optimality conditions read $Ax^* = z^*$ and

$$0 \in \partial \|\cdot\|_1(x^*) - A^* p^* \text{ and } 0 \in \partial f_\lambda(z^*) + p^*.$$

The first condition is equivalent to (14.7). For $\lambda > 0$, f_λ is smooth, and the second condition is equivalent to

$$p^* = \frac{y - A^* x^*}{\lambda} \text{ and } A^* p^* \in \partial \|\cdot\|_1(x^*)$$

which are the desired formula. For $\lambda = 0$, the second condition holds as soon as $z^* = Ax^* = y$. \square

Note that in the case $\lambda > 0$, (14.6) is strongly convex, and in fact the optimal solution p_λ is computed as an orthogonal projection

$$p_\lambda \in \operatorname{argmin}_{p \in \mathbb{R}^P} \{ \|p - y/\lambda\| ; \|A^*p\|_\infty \leq 1 \}.$$

The sup in (14.6) is thus actually a max if $\lambda > 0$. If $\lambda > 0$, in case $\ker(A^*) = \operatorname{Im}(A)^\perp = \{0\}$, the constraint set of the dual is bounded, so that the sup is also a max.

14.2 Consistency and Sparsitency

14.2.1 Bregman Divergence Rates for General Regularizations

Here we consider the case of a general regularization of the form

$$\min_{x \in \mathbb{R}^N} \frac{1}{2\lambda} \|Ax - y\|^2 + J(x) \quad (14.8)$$

for a convex regularizer J .

For any $\eta \in \partial J(x_0)$, we define the associated Bregman divergence as

$$D_\eta(x|x_0) \stackrel{\text{def.}}{=} J(x) - J(x_0) - \langle \eta, x - x_0 \rangle.$$

One has $D_\eta(x_0|x_0) = 0$, and since J is convex, one has $D_\eta(x|x_0) \geq 0$ [ToDo: put here drawings].

In the case where J is differentiable, since $\partial J(x_0) = \{\nabla J(x_0)\}$, this divergence simply reads

$$D(x|x_0) \stackrel{\text{def.}}{=} J(x) - J(x_0) - \langle \nabla J(x_0), x - x_0 \rangle.$$

If furthermore J is strictly convex, then $D(x|x_0) = 0$ if and only if $x = x_0$, so that $D(\cdot|\cdot)$ is similar to a distance function (but it does not necessarily satisfies the triangular inequality).

If $J = \|\cdot\|^2$, then $D(x|x_0) = \|x - x_0\|^2$ is the Euclidean norm. If $J(x) = \sum_i x_i (\log(x_i) - 1) + \iota_{\mathbb{R}^+}(x_i)$ is the entropy, then

$$D(x|x_0) = \sum_i x_i \log \left(\frac{x_i}{x_{0,i}} \right) + x_{0,i} - x_i$$

is the so-called Kulback-Leibler divergence on \mathbb{R}_+^N .

The following proposition, which is due to Burger-Osher, state a linear rate in term of this Bregman divergence.

Proposition 49. *If there exists*

$$\eta = A^*p \in \operatorname{Im}(A^*) \cap \partial J(x_0), \quad (14.9)$$

then one has for any x_λ solution of (14.8)

$$D_\eta(x_\lambda|x_0) \leq \frac{1}{2} \left(\frac{\|w\|}{\sqrt{\lambda}} + \sqrt{\lambda} \|p\| \right)^2. \quad (14.10)$$

Furthermore, one has the useful bound

$$\|Ax_\lambda - y\| \leq \|w\| + (\sqrt{2} + 1) \|p\| \lambda. \quad (14.11)$$

Proof. The optimality of x_λ for (14.8) implies

$$\frac{1}{2\lambda} \|Ax_\lambda - y\|^2 + J(x_\lambda) \leq \frac{1}{2\lambda} \|Ax_0 - y\|^2 + J(x_0) = \frac{1}{2\lambda} \|w\|^2 + J(x_0).$$

Hence, using $\langle \eta, x_\lambda - x_0 \rangle = \langle p, Ax_\lambda - Ax_0 \rangle = \langle p, Ax_\lambda - y + w \rangle$, one has

$$\begin{aligned} D_\eta(x_\lambda|x_0) &= J(x_\lambda) - J(x_0) - \langle \eta, x_\lambda - x_0 \rangle \leq \frac{1}{2\lambda} \|w\|^2 - \frac{1}{2\lambda} \|Ax_\lambda - y\|^2 - \langle p, Ax_\lambda - y \rangle - \langle p, w \rangle \\ &= \frac{1}{2\lambda} \|w\|^2 - \frac{1}{2\lambda} \|Ax_\lambda - y + \lambda p\|^2 + \lambda \|p\|^2 - \langle p, w \rangle \\ &\leq \frac{1}{2\lambda} \|w\|^2 + \frac{\lambda}{2} \|p\|^2 + \|p\| \|w\| = \frac{1}{2} \left(\frac{\|w\|}{\sqrt{\lambda}} + \sqrt{\lambda} \|p\| \right)^2. \end{aligned}$$

From the second line above, since $D_\eta(x_\lambda|x_0) \geq 0$, one has using Cauchy-Schwartz

$$\|Ax_\lambda - y + \lambda p\|^2 \leq \|w\|^2 + 2\lambda^2 \|p\|^2 + 2\lambda \|p\| \|w\| \leq \|w\|^2 + 2\sqrt{2} \|p\| \|w\| \lambda + 2\lambda^2 \|p\|^2 = \left(\|w\| + \sqrt{2} \lambda \|p\| \right)^2.$$

Hence

$$\|Ax_\lambda - y\| \leq \|Ax_\lambda - y + \lambda p\| + \lambda \|p\| \leq \|w\| + \sqrt{2} \lambda \|p\| + \lambda \|p\|.$$

□

Choosing $\lambda = \|w\|/\|p\|$ in (14.10), one thus obtain a linear rate in term of Bregman divergence $D_\eta(x_\lambda|x_0) \leq 2\|w\|\|p\|$. For the simple case of a quadratic regularized $J(x) = \|x\|^2/2$, as used in Section ??, one sees that the source conditions (14.9) simply reads

$$x_0 \in \text{Im}(A^*)$$

which is equivalent to (10.12) with exponent $\beta = \frac{1}{2}$, and under this condition, (14.10) gives the following sub-linear rate in term of the ℓ^2 norm

$$\|x_0 - x_\lambda\| \leq 2\sqrt{\|w\|\|p\|}.$$

[ToDo: This seems inconsistent, this should corresponds to $\beta = 1$ to obtain the same rates in both theorems!]

Note that the “source condition” (14.9) is equivalent to x_0 such that $Ax_0 = y$ is a solution to the constraint problem

$$\min_{Ax=y} J(x).$$

So simply being a solution of the constraint noiseless problem thus implies a linear rate for the resolution of the noisy problem in term of the Bregman divergence.

14.2.2 Linear Rates in Norms for ℓ^1 Regularization

The issue with the control (14.10) of the error in term of Bregman divergence is that it is not “distance-like” for regularizers J which are not strictly convex. This is in particular the case for the ℓ^1 norm $J = \|\cdot\|_1$ which we now study.

The following fundamental lemma shows however that this Bregman divergence for ℓ^1 behave like a distance (and in fact controls the ℓ^1 norm) on the indexes where η does not saturate.

Lemma 4. For $\eta \in \partial \|\cdot\|_1(x_0)$, let $J \stackrel{\text{def.}}{=} \text{sat}(\eta)$. Then

$$D_\eta(x|x_0) \geq (1 - \|\eta_{J^c}\|_\infty) \|(x - x_0)_{J^c}\|_1. \quad (14.12)$$

Proof. Note that $x_{0,J^c} = 0$ since $\text{supp}(x_0) \subset \text{sat}(\eta)$ by definition of the sub-differential of the ℓ^1 norm. Since the ℓ^1 norm is separable, each term in the sum defining $D_\eta(x|x_0)$ is positive, hence

$$\begin{aligned} D_\eta(x|x_0) &= \sum_i |x_i| - |x_{0,i}| - \eta_i(x_i - x_{0,i}) \geq \sum_{i \in J^c} |x_i| - |x_{0,i}| - \eta_i(x_i - x_{0,i}) \\ &= \sum_{i \in J^c} |x_i| - \eta_i x_i \geq \sum_{i \in J^c} (1 - |\eta_i|) |x_i| \geq (1 - \|\eta_{J^c}\|_\infty) \sum_{i \in J^c} |x_i| = (1 - \|\eta_{J^c}\|_\infty) \sum_{i \in J^c} |x_i - x_{0,i}|. \end{aligned}$$

□

The quantity $1 - \|\eta_{J^c}\|_\infty > 0$ controls how much η is “inside” the sub-differential. The larger this coefficients, the better is the control of the Bregman divergence.

The following theorem uses this lemma to state the convergence rate of the sparse regularized solution, under the same hypothesis as Proposition 48 (with $x^* = x_0$).

Theorem 42. *If there exists*

$$\eta \in \mathcal{D}_0(Ax_0, x_0) \quad (14.13)$$

and $\ker(A_J) = \{0\}$ where $J \stackrel{\text{def}}{=} \text{sat}(\eta)$ then choosing $\lambda = c\|w\|$, there exists C (depending on c) such that any solution x_λ of $\mathcal{P}(Ax_0 + w)$ satisfies

$$\|x_\lambda - x_0\| \leq C\|w\|. \quad (14.14)$$

Proof. We denote $y = Ax_0 + w$. The optimality of x_λ in $(\mathcal{P}_\lambda(y))$ implies

$$\frac{1}{2\lambda}\|Ax_\lambda - y\|^2 + \|x_\lambda\|_1 \leq \frac{1}{2\lambda}\|Ax_0 - y\|^2 + \|x_0\|_1 = \frac{1}{2\lambda}\|w\|^2 + \|x_0\|_1$$

and hence

$$\|Ax_\lambda - y\|^2 \leq \|w\|^2 + 2\lambda\|x_0\|_1$$

Using the fact that A_J is injective, one has $A_J^+ A_J = \text{Id}_J$, so that

$$\begin{aligned} \|(x_\lambda - x_0)_J\|_1 &= \|A_J^+ A_J(x_\lambda - x_0)_J\|_1 \leq \|A_J^+\|_{1,2} \|A_J x_{\lambda,J} - y + w\| \leq \|A_J^+\|_{1,2} (\|A_J x_{\lambda,J} - y\| + \|w\|) \\ &\leq \|A_J^+\|_{1,2} (\|Ax_\lambda - y\| + \|A_{J^c} x_{\lambda,J^c}\| + \|w\|) \\ &\leq \|A_J^+\|_{1,2} (\|Ax_\lambda - y\| + \|A_{J^c}\|_{2,1} \|x_{\lambda,J^c} - x_{0,J^c}\|_1 + \|w\|) \\ &\leq \|A_J^+\|_{1,2} \left(\|w\| + (\sqrt{2} + 1)\|p\|\lambda + \|A_{J^c}\|_{2,1} \|x_{\lambda,J^c} - x_{0,J^c}\|_1 + \|w\| \right) \end{aligned}$$

where we used $x_{0,J^c} = 0$ and (14.11). One plug this bound in the decomposition, and using (14.12) and (14.10)

$$\begin{aligned} \|x_\lambda - x_0\|_1 &= \|(x_\lambda - x_0)_J\|_1 + \|(x_\lambda - x_0)_{J^c}\|_1 \\ &\leq \|(x_\lambda - x_0)_{J^c}\|_1 (1 + \|A_J^+\|_{1,2} \|A_{J^c}\|_{2,1}) + \|A_J^+\|_{1,2} \left((\sqrt{2} + 1)\|p\|\lambda + 2\|w\| \right) \\ &\leq \frac{D_\eta(x|x_0)}{1 - \|\eta_{J^c}\|_\infty} (1 + \|A_J^+\|_{1,2} \|A_{J^c}\|_{2,1}) + \|A_J^+\|_{1,2} \left((\sqrt{2} + 1)\|p\|\lambda + 2\|w\| \right) \\ &\leq \frac{\frac{1}{2} \left(\frac{\|w\|}{\sqrt{\lambda}} + \sqrt{\lambda}\|p\| \right)^2}{1 - \|\eta_{J^c}\|_\infty} (1 + \|A_J^+\|_{1,2} \|A_{J^c}\|_{2,1}) + \|A_J^+\|_{1,2} \left((\sqrt{2} + 1)\|p\|\lambda + 2\|w\| \right). \end{aligned}$$

Thus setting $\lambda = c\|w\|$, one obtains the constant

$$C \stackrel{\text{def}}{=} \frac{\frac{1}{2} \left(\frac{1}{\sqrt{c}} + \sqrt{c}\|p\| \right)^2}{1 - \|\eta_{J^c}\|_\infty} (1 + \|A_J^+\|_{1,2} \|A_{J^c}\|_{2,1}) + \|A_J^+\|_{1,2} \left((\sqrt{2} + 1)\|p\|c + 2 \right).$$

□

Note that this theorem does not imply that x_λ is a unique solution, only x_0 is unique in general. The condition (14.13) is often called a “source condition”, and is strengthened by imposing a non-degeneracy $\ker(A_J) = \{0\}$. This non-degeneracy imply some stability in ℓ^2 sense (14.14). The result (14.14) shows a linear rate, i.e. the (possibly multi-valued) inverse map $y \mapsto x_\lambda$ is Lipschitz continuous.

It should be compared with Theorem 31 on linear methods for inverse problem regularization, which only gives sub-linear rate. The sources conditions in the linear (10.12) and non-linear (14.13) cases are however very different. In the linear case, for $\beta = 1/2$, it reads $x_0 \in \text{Im}(A^*) = \ker(A)^\perp$, which is mandatory because linear method cannot recover anything in $\ker(A)$. On contrary, the non-linear source condition only requires that η to be in $\text{Im}(A^*)$, and is able (in the favorable cases of course) to recover information in $\ker(A)$.

14.2.3 Sparsistency

Theorem 42 is abstract in the sense that it rely on hypotheses which are hard to check. The crux of the problem, to be able to apply this theorem, is to be able to “construct” a valid certificate (14.13). We now give a powerful “recipe” which – when it works – not only give a sufficient condition for linear rate, but also provides “support stability”.

For any solution x_λ of $(\mathcal{P}_\lambda(y))$, as already done in (14.2), we define the (unique, independent of the chosen solution) dual certificate

$$\eta_\lambda \stackrel{\text{def.}}{=} A^* p_\lambda \quad \text{where} \quad p_\lambda \stackrel{\text{def.}}{=} \frac{y - Ax_\lambda}{\lambda}.$$

The following proposition shows that p_λ converge to a very specific dual certificate of the constrained problem, which we coined “minimal norm” certificate.

Proposition 50. *If $y = Ax_0$ where x_0 is a solution to $(\mathcal{P}_\lambda(y = Ax_0))$, one has*

$$p_\lambda \rightarrow p_0 \stackrel{\text{def.}}{=} \underset{p \in \mathbb{R}^P}{\operatorname{argmin}} \{ \|p\| ; A^* p \in \mathcal{D}_0(y, x_0) \}. \quad (14.15)$$

The vector $\eta_0 \stackrel{\text{def.}}{=} A^* p_0$ is called the “minimum norm certificate”.

Proof. This follows from the fact that p_λ is the unique solution to (14.6) and then applying the same proof as the one done in Proposition 23 to study the small λ limit of penalized problems. \square

This proposition shows that, while dual certificate $\mathcal{D}_0(y, x_0)$ for $\lambda = 0$ are non-unique, taking the limit as $\lambda \rightarrow 0$ singles-out a specific one, which is of paramount importance to study stability of the support when the noise and λ are small.

A major difficulty in computing (14.24) is that it should satisfy the non-linear constraint $\|\eta_0\|_\infty$. One thus can “simplify” this definition by removing this ℓ^∞ constraint and define the so-called “minimum norm certificate”

$$\eta_F \stackrel{\text{def.}}{=} A^* p_F \quad \text{where} \quad p_F \stackrel{\text{def.}}{=} \underset{p \in \mathbb{R}^P}{\operatorname{argmin}} \{ \|p\| ; A_I^* p = \operatorname{sign}(x_{0,I}) \}. \quad (14.16)$$

The notation “ η_F ” refers to the “Fuchs” certificate, which we named in honour of J-J. Fuchs who first used it to study ℓ^1 minimization.

We insists that p_F is not necessarily a valid certificate (hence the naming “pre-certificate”) since one does not have in general $\|\eta_F\|_\infty \leq 1$. The vector p_F is a least square solution to the linear system $A_I^* p = \operatorname{sign}(x_{0,I})$, and it can thus be compute in closed form using the pseudo-inverse $p_F = A_I^{*,+} \operatorname{sign}(x_{0,I})$ (see Proposition (21)). In case $\ker(A_I) = \{0\}$, one has the simple formula

$$p_F = A_I(A_I^* A_I)^{-1} \operatorname{sign}(x_{0,I}).$$

Denoting $C \stackrel{\text{def.}}{=} A^* A$ the “correlation” matrix, one has the nice formula

$$\eta_F = C_{\cdot, I} C_{I, I}^{-1} \operatorname{sign}(x_{0,I}). \quad (14.17)$$

The following proposition relates η_F to η_0 , and shows that η_F can be used as a “proxy” for η_0

Proposition 51. *If $\|\eta_F\|_\infty \leq 1$, then $p_F = p_0$ and $\eta_F = \eta_0$.*

The condition $\|\eta_F\|_\infty \leq 1$ implies that x_0 is solution to $(\mathcal{P}_0(y))$. The following theorem shows that if one strengthen this condition to impose a non-degeneracy on η_F , then one has linear rate with a stable support in the small noise regime.

Remark 8 (Operator norm). In the proof, we use the $\ell^p - \ell^q$ matrix operator norm, which is defined as

$$\|B\|_{p,q} \stackrel{\text{def.}}{=} \max \{ \|Bu\|_q ; \|u\|_p \leq 1 \}.$$

For $p = q$, we denote $\|B\|_p \stackrel{\text{def.}}{=} \|B\|_{p,p}$. For $p = 2$, $\|B\|_2$ is the maximum singular value, and one has

$$\|B\|_1 = \max_j \sum_i |B_{i,j}| \quad \text{and} \quad \|B\|_\infty = \max_i \sum_j |B_{i,j}|.$$

Theorem 43. *If*

$$\|\eta_F\|_\infty \leq 1 \quad \text{and} \quad \|\eta_{F,I^c}\|_\infty < 1,$$

and $\ker(A_I) = \{0\}$, then there exists C, C' such that if $\max(\|w\|, \|w\|/\lambda) \leq C$, then the solution x_λ of $(\mathcal{P}_\lambda(y))$ is unique, is supported in I , and in fact

$$x_{\lambda,I} = x_{0,I} + A_I^+ w - \lambda(A_I^* A_I)^{-1} \text{sign}(x_{0,I}^*). \quad (14.18)$$

In particular, $\|x_\lambda - x_0\| = O(\|w\|)$.

Proof. In the following we denote $T \stackrel{\text{def.}}{=} \min_{i \in I} |x_{0,i}|$ the signal level, and $\delta \stackrel{\text{def.}}{=} \|A^* w\|_\infty$ which is the natural way to measure the noise amplitude in the sparse setting. We define $s \stackrel{\text{def.}}{=} \text{sign}(x_0)$, and consider the “ansatz” (14.18) and thus define the following candidate solution

$$\hat{x}_I \stackrel{\text{def.}}{=} x_{0,I} + A_I^+ w - \lambda(A_I^* A_I)^{-1} s_I, \quad (14.19)$$

and $\hat{x}_{I^c} = 0$. The goal is to show that \hat{x} is indeed the unique solution of $(\mathcal{P}_\lambda(y))$.

Step 1. The first step is to show sign consistency, i.e. that $\text{sign}(\hat{x}) = s$. This is true if $\|x_{0,I} - \hat{x}_I\|_\infty < T$, and is thus implied by

$$\|x_{0,I} - \hat{x}_I\|_\infty \leq K \|A_I^* w\|_\infty + K\lambda < T \quad \text{where} \quad K \stackrel{\text{def.}}{=} \|(A_I^* A_I)^{-1}\|_\infty, \quad (14.20)$$

where we used the fact that $A_I^+ = (A_I^* A_I)^{-1} A_I^*$.

Step 2. The second step is to check the first order condition of Proposition 47, i.e. $\|\hat{\eta}_{I^c}\|_\infty < 1$, where $\lambda \hat{\eta} = A^*(y - A\hat{x})$. This implies indeed that \hat{x} is the unique solution of $(\mathcal{P}_\lambda(y))$. One has

$$\begin{aligned} \lambda \hat{\eta} &= A^*(A_I x_{0,I} + w - A_I (x_{0,I} + A_I^+ w - \lambda(A_I^* A_I)^{-1} s_I)) \\ &= A^*(A_I A_I^+ - \text{Id})w + \lambda \eta_F. \end{aligned}$$

The condition $\|\hat{\eta}_{I^c}\|_\infty < 1$ is thus implied by

$$\|A_{I^c}^* A_I (A_I^* A_I)^{-1}\|_\infty \|A_I^* w\|_\infty + \|A_{I^c}^* w\|_\infty + \lambda \|\eta_{F,I^c}\|_\infty \leq R \|A_I^* w\|_\infty - S\lambda < 0 \quad (14.21)$$

$$R \stackrel{\text{def.}}{=} KL + 1 \quad \text{and} \quad S \stackrel{\text{def.}}{=} 1 - \|\eta_{F,I^c}\|_\infty > 0$$

where we denoted $L \stackrel{\text{def.}}{=} \|A_{I^c}^* A_I\|_\infty$, and also we used the hypothesis $\|\eta_{F,I^c}\|_\infty < 1$.

Conclusion. Putting (14.20) and (14.21) together shows that \hat{x} is the unique solution if (λ, w) are such that the two linear inequations are satisfied

$$\mathcal{R} = \left\{ (\delta, \lambda) ; \delta + \lambda < \frac{T}{K} \quad \text{and} \quad R\delta - S\lambda < 0 \right\}$$

This region \mathcal{R} is triangular-shaped, and includes the following “smaller” simpler triangle

$$\tilde{\mathcal{R}} = \left\{ (\delta, \lambda) ; \frac{\delta}{\lambda} < \frac{S}{R} \quad \text{and} \quad \lambda < \lambda_{\max} \right\} \quad \text{where} \quad \lambda_{\max} \stackrel{\text{def.}}{=} \frac{T(KL + 1)}{K(R + S)}. \quad (14.22)$$

□




Figure 14.3: Display of certificate η_F for a $A \in \mathbb{R}^{P \times n}$, $n = 64$, with independent Gaussian entries.

It is important to realize that Theorem 43 operates in a “small noise” regime, i.e. $\|w\|$ (and hence λ) needs to be small enough for the support to be identifiable (otherwise small amplitude component of x_0 will be killed by the regularization). In contrast, Theorem 42 is “global” and holds for any noise level $\|w\|$. The price to pay is that one has no controls about the support (and one does not even know whether x_λ is unique) and then the constants involved are more pessimistic.

A nice feature of this proof is that it gives access to explicit constants, involving the three key parameters K, L, S , which controls:

- K accounts for the conditioning of the operator on the support I ;
- L accounts for the worst correlation between atoms inside and outside the support ;
- S accounts for how much the certificates η_F are non-degenerate.

The constants on $\|A^*w\|/\lambda$ and on λ are given by (14.22). Choosing (which is in practice impossible, because it requires knowledge about the solution) the smallest possible λ gives $\lambda = \delta \frac{S}{R}$ and in this regime the error is bounded in ℓ^∞ (using other error norms would simply lead to using other matrix norms)

$$\|x_0 - x_\lambda\|_\infty \leq \left(1 + \frac{KL + 1}{S}\right) K\delta.$$

The crux of the analysis of the performance (in terms of support stability) of ℓ^1 regularization is to be able to say whether, for some class of signal x_0 of interest, η_F is a valid certificate, i.e. $\|\eta_F\|_\infty \leq 1$. Figure 14.4 displays numerically what one obtains when A is random. One sees that η_F is non-degenerate when P is large enough. Section 15.2 performs a mathematical analysis of this phenomenon.

14.3 Sparse Deconvolution Case Study

Chapter 15 studies the particular case where A is random, in which case it is possible to make very precise statements about whether η_F is a valid certificate.

Another interesting case study, which shows the limitation of this approach, is the case of super-resolution. To simplify the analysis, we assume “continuous measurement”, and replace the measurement space \mathbb{R}^P by functions $L^2(\mathbb{R}^d)$ (for simplicity, we treat here $d = 1$). The measurements read

$$Ax = \sum_{i=1}^N x_i a_i(\cdot) \in L^2(\mathbb{R})$$

where the $a_i : \mathbb{R} \rightarrow \mathbb{R}$ are smooth functions. A typical example is the deconvolution problem, where $a_i = \varphi(\cdot - z_i)$ where $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ is the smoothing kernel and $(z_i)_{i=1}^N$ is a discretization grid, and for simplicity we assume $x_i = i/N \in [0, 1]$.

In this case, this forward model correspond to convolution of measures supported on the grid

$$Ax = \varphi \star m_{z,x} \quad \text{where} \quad m_{z,x} \stackrel{\text{def.}}{=} \sum_{i=1}^N x_i \delta_{z_i}.$$

We denote the “continuous” covariance

$$\mathcal{C}(z, z') \stackrel{\text{def.}}{=} \langle \varphi(\cdot - z), \varphi(\cdot - z') \rangle_{L^2(\mathbb{R})} = \int_{\mathbb{R}} \varphi(t - z) \varphi(t - z') dt = (\varphi \star \bar{\varphi})(z - z')$$

where $\bar{\varphi}(t) = \varphi(-t)$, so that the discrete covariance is $C = (\mathcal{C}(z_i, z'_i))_{(i,i') \in I^2} \in \mathbb{R}^{N \times N}$ and $C_{I,I} = (\mathcal{C}(z_i, z'_i))_{(i,i') \in I^2} \in \mathbb{R}^{I \times I}$.

Using (14.17), one sees that η_F is obtained as a sampling on the grid of a “continuous” certificate $\tilde{\eta}_F$

$$\eta_F = (\tilde{\eta}_F(z_i))_{i=1}^N \in \mathbb{R}^N,$$

$$\text{where } \tilde{\eta}_F(x) = \sum_{i \in I} b_i \mathcal{C}(x, z_i) \quad \text{where} \quad b_I = C_{I,I}^{-1} \text{sign}(x_{0,I}), \quad (14.23)$$

so that η_F is a linear combination of I basis functions $(\mathcal{C}(x, z_i))_{i \in I}$.

The question is whether $\|\eta_F\|_{\ell^\infty} \leq 1$. If the grid is fine enough, i.e. N large enough, this can only hold if $\|\tilde{\eta}_F\|_{L^\infty} \leq 1$. The major issue is that $\tilde{\eta}_F$ is only constrained by construction to interpolate $\text{sign}(x_{0,i})$ are points $z_{0,i}$ for $i \in I$. So nothing prevents $\tilde{\eta}_F$ to go outside $[-1, 1]$ around each interpolation point. Figure ?? illustrates this fact.

In order to guarantee this property of “local” non-degeneracy around the support, one has to impose on the certificate the additional constraint $\eta'(z_i) = 0$ for $i \in I$. This leads to consider a minimum pre-certificate with vanishing derivatives

$$\eta_V \stackrel{\text{def.}}{=} A^* p_V \quad \text{where} \quad p_V \argmin_{p \in L^2(\mathbb{R})} \{ \|p\|_{L^2(\mathbb{R})} ; \tilde{\eta}(z_I) = \text{sign}(x_{0,I}), \tilde{\eta}'(z_I) = \mathbf{0}_I \}. \quad (14.24)$$

where we denoted $\tilde{\eta} = \bar{\psi} \star p$. Similarly to (14.23), this vanishing pre-certificate can be written as a linear combination, but this time of $2|I|$ basis functions

$$\tilde{\eta}_V(x) = \sum_{i \in I} b_i \mathcal{C}(x, z_i) + c_i \partial_2 \mathcal{C}(x, z_i),$$

where $\partial_2 \mathcal{C}$ is the derivative of \mathcal{C} with respect to the second variable, and (b, c) are solution of a $2|I| \times 2|I|$ linear system

$$\begin{pmatrix} b \\ c \end{pmatrix} = \begin{pmatrix} (\mathcal{C}(x_i, x_{i'}))_{i,i' \in I^2} & (\partial_2 \mathcal{C}(x_i, x_{i'}))_{i,i' \in I^2} \\ (\partial_1 \mathcal{C}(x_i, x_{i'}))_{i,i' \in I^2} & (\partial_1 \partial_2 \mathcal{C}(x_i, x_{i'}))_{i,i' \in I^2} \end{pmatrix}^{-1} \begin{pmatrix} \text{sign}(x_{0,I}) \\ \mathbf{0}_I \end{pmatrix}.$$

The associated continuous pre-certificate is $\tilde{\eta}_V = \bar{\psi} \star p_V$, and η_V is a sampling on the grid of $\tilde{\eta}_V$. Figure 14.4 shows that this pre-certificate η_V is much better behaved than η_F . If $\|\eta_V\|_\infty \leq 1$, one can apply (42) and thus obtain a linear convergence rate with respect to the ℓ^2 norm on the grid. But for very fine grid, since one is interested in sparse solution, the ℓ^2 norm becomes meaningless (because the L^2 norm is not defined on measures). Since η_V is different from η_F , one cannot directly applies Theorem 43: the support is not stable on discrete grids, which is a fundamental property of super-resolution problems (as opposed to compressed sensing problems). The way to recover interesting results is to use and analyze methods without grids. Indeed, after removing the grid, one can show that η_V becomes the minimum norm certificate (and is the limit of η_λ).




Figure 14.4: Display of “continuous” certificate η_F and η_V for A being a convolution operator.

Bibliography

- [1] P. Alliez and C. Gotsman. Recent advances in compression of 3d meshes. In N. A. Dodgson, M. S. Floater, and M. A. Sabin, editors, *Advances in multiresolution for geometric modelling*, pages 3–26. Springer Verlag, 2005.
- [2] P. Alliez, G. Ucelli, C. Gotsman, and M. Attene. Recent advances in remeshing of surfaces. In *AIM@SHAPE report*. 2005.
- [3] Amir Beck. *Introduction to Nonlinear Optimization: Theory, Algorithms, and Applications with MATLAB*. SIAM, 2014.
- [4] Stephen Boyd, Neal Parikh, Eric Chu, Borja Peleato, and Jonathan Eckstein. Distributed optimization and statistical learning via the alternating direction method of multipliers. *Foundations and Trends® in Machine Learning*, 3(1):1–122, 2011.
- [5] Stephen Boyd and Lieven Vandenbergh. *Convex optimization*. Cambridge university press, 2004.
- [6] E. Candès and D. Donoho. New tight frames of curvelets and optimal representations of objects with piecewise C^2 singularities. *Commun. on Pure and Appl. Math.*, 57(2):219–266, 2004.
- [7] E. J. Candès. The restricted isometry property and its implications for compressed sensing. *Compte Rendus de l’Académie des Sciences, Serie I*(346):589–592, 2006.
- [8] E. J. Candès, L. Demanet, D. L. Donoho, and L. Ying. Fast discrete curvelet transforms. *SIAM Multiscale Modeling and Simulation*, 5:861–899, 2005.
- [9] A. Chambolle. An algorithm for total variation minimization and applications. *J. Math. Imaging Vis.*, 20:89–97, 2004.
- [10] Antonin Chambolle, Vicent Caselles, Daniel Cremers, Matteo Novaga, and Thomas Pock. An introduction to total variation for image analysis. *Theoretical foundations and numerical methods for sparse recovery*, 9(263-340):227, 2010.
- [11] Antonin Chambolle and Thomas Pock. An introduction to continuous optimization for imaging. *Acta Numerica*, 25:161–319, 2016.
- [12] S.S. Chen, D.L. Donoho, and M.A. Saunders. Atomic decomposition by basis pursuit. *SIAM Journal on Scientific Computing*, 20(1):33–61, 1999.
- [13] F. R. K. Chung. Spectral graph theory. *Regional Conference Series in Mathematics, American Mathematical Society*, 92:1–212, 1997.
- [14] Philippe G Ciarlet. Introduction à l’analyse numérique matricielle et à l’optimisation. 1982.
- [15] P. L. Combettes and V. R. Wajs. Signal recovery by proximal forward-backward splitting. *SIAM Multiscale Modeling and Simulation*, 4(4), 2005.

- [16] P. Schroeder et al. D. Zorin. Subdivision surfaces in character animation. In *Course notes at SIGGRAPH 2000*, July 2000.
- [17] I. Daubechies, M. Defrise, and C. De Mol. An iterative thresholding algorithm for linear inverse problems with a sparsity constraint. *Commun. on Pure and Appl. Math.*, 57:1413–1541, 2004.
- [18] I. Daubechies and W. Sweldens. Factoring wavelet transforms into lifting steps. *J. Fourier Anal. Appl.*, 4(3):245–267, 1998.
- [19] D. Donoho and I. Johnstone. Ideal spatial adaptation via wavelet shrinkage. *Biometrika*, 81:425–455, Dec 1994.
- [20] Heinz Werner Engl, Martin Hanke, and Andreas Neubauer. *Regularization of inverse problems*, volume 375. Springer Science & Business Media, 1996.
- [21] M. Figueiredo and R. Nowak. An EM Algorithm for Wavelet-Based Image Restoration. *IEEE Trans. Image Proc.*, 12(8):906–916, 2003.
- [22] M. S. Floater and K. Hormann. Surface parameterization: a tutorial and survey. In N. A. Dodgson, M. S. Floater, and M. A. Sabin, editors, *Advances in multiresolution for geometric modelling*, pages 157–186. Springer Verlag, 2005.
- [23] Simon Foucart and Holger Rauhut. *A mathematical introduction to compressive sensing*, volume 1. Birkhäuser Basel, 2013.
- [24] I. Guskov, W. Sweldens, and P. Schröder. Multiresolution signal processing for meshes. In Alyn Rockwood, editor, *Proceedings of the Conference on Computer Graphics (Siggraph99)*, pages 325–334. ACM Press, August8–13 1999.
- [25] A. Khodakovsky, P. Schröder, and W. Sweldens. Progressive geometry compression. In *Proceedings of the Computer Graphics Conference 2000 (SIGGRAPH-00)*, pages 271–278, New York, July 23–28 2000. ACM Press.
- [26] L. Kobbelt. $\sqrt{3}$ subdivision. In Sheila Hoffmeyer, editor, *Proc. of SIGGRAPH’00*, pages 103–112, New York, July 23–28 2000. ACM Press.
- [27] M. Lounsbery, T. D. DeRose, and J. Warren. Multiresolution analysis for surfaces of arbitrary topological type. *ACM Trans. Graph.*, 16(1):34–73, 1997.
- [28] S. Mallat. *A Wavelet Tour of Signal Processing, 3rd edition*. Academic Press, San Diego, 2009.
- [29] Stephane Mallat. *A wavelet tour of signal processing: the sparse way*. Academic press, 2008.
- [30] D. Mumford and J. Shah. Optimal approximation by piecewise smooth functions and associated variational problems. *Commun. on Pure and Appl. Math.*, 42:577–685, 1989.
- [31] Neal Parikh, Stephen Boyd, et al. Proximal algorithms. *Foundations and Trends® in Optimization*, 1(3):127–239, 2014.
- [32] Gabriel Peyré. *L’algèbre discrète de la transformée de Fourier*. Ellipses, 2004.
- [33] Gabriel Peyré and Marco Cuturi. Computational optimal transport. 2017.
- [34] J. Portilla, V. Strela, M.J. Wainwright, and Simoncelli E.P. Image denoising using scale mixtures of Gaussians in the wavelet domain. *IEEE Trans. Image Proc.*, 12(11):1338–1351, November 2003.
- [35] E. Praun and H. Hoppe. Spherical parametrization and remeshing. *ACM Transactions on Graphics*, 22(3):340–349, July 2003.

- [36] L. I. Rudin, S. Osher, and E. Fatemi. Nonlinear total variation based noise removal algorithms. *Phys. D*, 60(1-4):259–268, 1992.
- [37] Filippo Santambrogio. Optimal transport for applied mathematicians. *Birkäuser, NY*, 2015.
- [38] Otmar Scherzer, Markus Grasmair, Harald Grossauer, Markus Haltmeier, Frank Lenzen, and L Sirovich. *Variational methods in imaging*. Springer, 2009.
- [39] P. Schröder and W. Sweldens. Spherical Wavelets: Efficiently Representing Functions on the Sphere. In *Proc. of SIGGRAPH 95*, pages 161–172, 1995.
- [40] P. Schröder and W. Sweldens. Spherical wavelets: Texture processing. In P. Hanrahan and W. Purgathofer, editors, *Rendering Techniques '95*. Springer Verlag, Wien, New York, August 1995.
- [41] C. E. Shannon. A mathematical theory of communication. *The Bell System Technical Journal*, 27(3):379–423, 1948.
- [42] A. Sheffer, E. Praun, and K. Rose. Mesh parameterization methods and their applications. *Found. Trends. Comput. Graph. Vis.*, 2(2):105–171, 2006.
- [43] Jean-Luc Starck, Fionn Murtagh, and Jalal Fadili. *Sparse image and signal processing: Wavelets and related geometric multiscale analysis*. Cambridge university press, 2015.
- [44] W. Sweldens. The lifting scheme: A custom-design construction of biorthogonal wavelets. *Applied and Computation Harmonic Analysis*, 3(2):186–200, 1996.
- [45] W. Sweldens. The lifting scheme: A construction of second generation wavelets. *SIAM J. Math. Anal.*, 29(2):511–546, 1997.