

Our math formulas, like $x^n + y^n = z^n$, and

$$\sum_{i=1}^n \sin x + i^{\sin x} + i^{i^{\sin x}}$$

are going to be using the MathTime Professional 2 fonts, but the text font is just Computer Modern (the letters for ‘sin’ are going to come from cmr10, cmr7 and cmr5).

Here are some math formulas that should all work out OK.

$$\begin{array}{ccccc} A, \ldots, Z & a, \ldots, z & \Gamma, \ldots, \Omega & \Gamma, \ldots, \Omega & \alpha, \ldots, \omega \\ 2^{A, \ldots, Z} & 2^{a, \ldots, z} & 2^{\Gamma, \ldots, \Omega} & 2^{\Gamma, \ldots, \Omega} & 2^{\alpha, \ldots, \omega} \\ 2^{2^{A, \ldots, Z}} & 2^{2^{a, \ldots, z}} & 2^{2^{\Gamma, \ldots, \Omega}} & 2^{\Gamma, \ldots, \Omega} & 2^{\alpha, \ldots, \omega} \end{array}$$

$$\begin{array}{l} \aleph_\alpha \times \aleph_\beta = \beta \iff \alpha \leq \beta \\ 2^{\aleph_\alpha \times \aleph_\beta = \beta} \iff \alpha \leq \beta \\ 2^{2^{\aleph_\alpha \times \aleph_\beta = \beta}} \iff \alpha \leq \beta \end{array}$$

$$\begin{array}{l} \forall \varepsilon > \alpha, \Gamma_\alpha \hookrightarrow \Gamma_\varepsilon \\ 2^{\forall \varepsilon > \alpha, \Gamma_\alpha \hookrightarrow \Gamma_\varepsilon} \\ 2^{2^{\forall \varepsilon > \alpha, \Gamma_\alpha \hookrightarrow \Gamma_\varepsilon}} \end{array}$$

$$\begin{array}{l} |x-a| < \delta \implies |f(x)-l| < \varepsilon \\ 2^{|x-a| < \delta \implies |f(x)-l| < \varepsilon} \\ 2^{2^{|x-a| < \delta \implies |f(x)-l| < \varepsilon}} \end{array}$$

$$\begin{array}{l} \underbrace{V \times \cdots \times V}_k \times \underbrace{V \times \cdots \times V}_l \rightarrow \underbrace{V \times \cdots \times V}_{k+l} \\ 2^{\underbrace{V \times \cdots \times V}_k \times \underbrace{V \times \cdots \times V}_l \rightarrow \underbrace{V \times \cdots \times V}_{k+l}} \\ 2^{2^{\underbrace{V \times \cdots \times V}_k \times \underbrace{V \times \cdots \times V}_l \rightarrow \underbrace{V \times \cdots \times V}_{k+l}}} \end{array}$$

$$\begin{array}{l} \{x|x \neq x\} = \emptyset \qquad (A \cap B)^\circ \subset A^\circ \cap B^\circ \\ 2^{\{x|x \neq x\} = \emptyset} \qquad (A \cap B)^\circ \subset A^\circ \cap B^\circ \\ 2^{2^{\{x|x \neq x\} = \emptyset}} \qquad (A \cap B)^\circ \subset A^\circ \cap B^\circ \end{array}$$

$$\begin{array}{l} \omega = v + v(x,y) \, dx + w(x,y) \, dy + d\kappa \\ 2^{\omega = v + v(x,y) \, dx + w(x,y) \, dy + d\kappa} \\ 2^{2^{\omega = v + v(x,y) \, dx + w(x,y) \, dy + d\kappa}} \end{array}$$

$$d\omega = dv + \left(\frac{\partial w}{\partial x} - \frac{\partial v}{\partial y}\right) dx \wedge dy$$

$$2^{d\omega = dv + \left(\frac{\partial w}{\partial x} - \frac{\partial v}{\partial y}\right) dx \wedge dy}$$

$$2^2^{d\omega = dv + \left(\frac{\partial w}{\partial x} - \frac{\partial v}{\partial y}\right) dx \wedge dy}$$

$$\hat{x} + \widehat{X} + \widehat{xy} + \widehat{xyz} + \vec{A}$$

$$2^{\hat{x}+\widehat{X}+\widehat{xy}+\widehat{xyz}+\vec{A}}$$

$$2^{2^{\hat{x}+\widehat{X}+\widehat{xy}+\widehat{xyz}+\vec{A}}}$$

$$R_{ijkl}=-R_{jikl}=-R_{ijlk}=R_{klij}$$

$$2^{R_{ijkl}=-R_{jikl}=-R_{ijlk}=R_{klij}}$$

$$2^2^{R_{ijkl}=-R_{jikl}=-R_{ijlk}=R_{klij}}$$

$$(f\circ g)'(x)=f'(g(x))\cdot g'(x)$$

$$2^{(f\circ g)'(x)=f'(g(x))\cdot g'(x)}$$

$$2^2^{(f\circ g)'(x)=f'(g(x))\cdot g'(x)}$$

$$f(x)=\left\{\begin{array}{ll} |x| & x>a \\ -|x| & x\leq a \end{array}\right.$$

$$2^{f(x)=\left\{\begin{array}{ll} |x| & x>a \\ -|x| & x\leq a \end{array}\right.}$$

$$2^2^{f(x)=\left\{\begin{array}{ll} |x| & x>a \\ -|x| & x\leq a \end{array}\right.}$$

$$\int_{-\infty}^{\infty}e^{-x\cdot x}\,dx=\sqrt{\pi}$$

$$2^{\int_{-\infty}^{\infty}e^{-x\cdot x}\,dx=\sqrt{\pi}}$$

$$2^2^{\int_{-\infty}^{\infty}e^{-x\cdot x}\,dx=\sqrt{\pi}}$$

$$X=\sum_i \xi^i \frac{\partial}{\partial x^i} + \sum_j x^j \frac{\partial}{\partial \dot{x}^j}$$

$$2^{X=\sum_i \xi^i \frac{\partial}{\partial x^i} + \sum_j x^j \frac{\partial}{\partial \dot{x}^j}}$$

$$2^2^{X=\sum_i \xi^i \frac{\partial}{\partial x^i} + \sum_j x^j \frac{\partial}{\partial \dot{x}^j}}$$

Bold letters in math can be taken from the Times bold symbols:

$$A_{\mathbf{X}}(f) = \mathbf{X}(\mathbf{f}) = 2^{2^{\mathbf{X}(\mathbf{g})}}$$

We can also get ‘calligraphic’ letters:

$$\mathcal{A}, \mathcal{B}, \dots, \mathcal{Z}$$

Compare

$$X_f + X_j + X_p + X_t + X_y + X_A + X_B + X_D + X_H + X_I + X_K + X_L + X_M + X_P + X_X$$

with the following (with no adjustments):

$$X_f + X_j + X_p + X_t + X_y + X_A + X_B + X_D + X_H + X_I + X_K + X_L + X_M + X_P + X_X$$

We have the special accent

$$\overset{\circ}{x}$$

and can replace

$$\dot{I} + \ddot{I}$$

with

$$\dot{I} + \ddot{I}$$

There are

$$\hat{A} + \hat{A} + \hat{A} + \hat{A} + + \hat{M} + \hat{M} + \hat{M} + \hat{M} + \widehat{xy} + \widehat{xyz} + \widehat{xyzw} + \overline{x + y + z + \dots + w}$$

and

$$\tilde{A} + \tilde{A} + \tilde{A} + \tilde{A} + + \tilde{M} + \tilde{M} + \tilde{M} + \tilde{M} + \widetilde{xy} + \widetilde{xyz} + \widetilde{xyzw} + \overline{x + y + z + \dots + w}$$

and

$$\check{A} + \check{A} + \check{A} + \check{A} + + \check{M} + \check{M} + \check{M} + \check{M} + \overline{\check{xy}} + \overline{\check{xyz}} + \overline{\check{xyzw}} + \overline{x + y + z + \dots + w}$$

and

$$\bar{M} + \bar{M} + \bar{M} + \overline{x + y + z}$$

We have

$$\alpha_c^{-1} \cdot \alpha_c' = \begin{pmatrix} 0 & 0 & \dots & -\kappa_1 \\ 1 & 0 & & -\kappa_2 \\ 0 & 1 & & \vdots \\ \vdots & \vdots & & -\kappa_{n-1} \\ 0 & 0 & \dots & 1 & 0 \end{pmatrix}$$

versus

$$\alpha_c^{-1} \cdot \alpha_c' = \begin{pmatrix} 0 & 0 & \dots & -\kappa_1 \\ 1 & 0 & & -\kappa_2 \\ 0 & 1 & & \vdots \\ \vdots & \vdots & & -\kappa_{n-1} \\ 0 & 0 & \dots & 1 & 0 \end{pmatrix}$$

Similarly, instead of having to rely on an extensible square root symbol, we can also get individually designed ones:

$$\sqrt{\sum_{i=1}^n (y^i - x^i)^2} \quad \text{vs.} \quad \sqrt{\sum_{i=1}^n (y^i - x^i)^2}$$