

GSE245908 | CHC20 | PGAM5-high Mac-RNA / immune subgroup neighborhoods


All T-lineage (baseline)
27 High sources; 64 edges




Exhaustion-supported T
0 High sources; 0 edges




Early activation-supported T
3 High sources; 3 edges




CD8 / cytotoxic T-related
0 High sources; 0 edges




CCR7/TCF7 naive-memory T
11 High sources; 12 edges




FOXP3/IKZF2 Treg-related
0 High sources; 0 edges




KLRD1/GNLY NK-related
1 High sources; 1 edges




MS4A1/CD79A B-related
1 High sources; 1 edges




MZB1/JCHAIN plasma-related
14 High sources; 15 edges




CD1C/FCER1A cDC2-related
8 High sources; 12 edges




CLEC9A/XCR1 cDC1-related
0 High sources; 0 edges




CLEC4C/IL3RA pDC-related
0 High sources; 0 edges




FCN1/VCAN inflammatory monocyte
30 High sources; 98 edges



FCGR3B/CSF3R neutrophil-related
12 High sources; 13 edges



TPSB2/CPA3 mast-related
1 High sources; 1 edges



Original H&E coordinates
Blue: target RNA gate
Gold rings: PGAM5-high Mac-RNA
Purple lines: direct spot neighbors

RNA gates can overlap.
Zero edges != absent cell type.
No cell-specific communication
is established.

Main QC and original PGAM5 Q75 labels are unchanged. Non-self hex-grid neighbors only; no spatial location inferred from UMAP.
Target programs are provisional mixed-spot RNA annotations; not pathology ROIs, pure single cells, or interaction probabilities.