

GSE245908 | CHC23 | PGAM5-high Mac-RNA / immune subgroup neighborhoods


All T-lineage (baseline)
8 High sources; 20 edges




Exhaustion-supported T
1 High sources; 2 edges




Early activation-supported T
0 High sources; 0 edges




CD8 / cytotoxic T-related
4 High sources; 5 edges




CCR7/TCF7 naive-memory T
1 High sources; 3 edges




FOXP3/IKZF2 Treg-related
2 High sources; 2 edges




KLRD1/GNLY NK-related
2 High sources; 2 edges




MS4A1/CD79A B-related
0 High sources; 0 edges




MZB1/JCHAIN plasma-related
7 High sources; 19 edges




CD1C/FCER1A cDC2-related
1 High sources; 1 edges




CLEC9A/XCR1 cDC1-related
0 High sources; 0 edges




CLEC4C/IL3RA pDC-related
0 High sources; 0 edges




FCN1/VCAN inflammatory monocyte
6 High sources; 14 edges



FCGR3B/CSF3R neutrophil-related
4 High sources; 6 edges



TPSB2/CPA3 mast-related
4 High sources; 4 edges



Original H&E coordinates
Blue: target RNA gate
Gold rings: PGAM5-high Mac-RNA
Purple lines: direct spot neighbors

RNA gates can overlap.
Zero edges != absent cell type.
No cell-specific communication
is established.

Main QC and original PGAM5 Q75 labels are unchanged. Non-self hex-grid neighbors only; no spatial location inferred from UMAP.
Target programs are provisional mixed-spot RNA annotations; not pathology ROIs, pure single cells, or interaction probabilities.