Candidate ligand-receptor RNA co-detection (descriptive)

CHC20 | High sources to immune subgroup targets

CHC23 | High sources to immune subgroup targets

All T-lineage (baseline)
Exhaustion-supported T
Early activation-supported T
CD8 / cytotoxic T-related
CCR7/TCF7 naive-memory T
FOXP3/IKZF2 Treg-related
KLRD1/GNLY NK-related
MS4A1/CD79A B-related
MZB1/JCHAIN plasma-related
CD1C/FCER1A cDC2-related
CLEC9A/XCR1 cDC1-related
CLEC4C/IL3RA pDC-related
FCN1/VCAN inflammatory monocyte
FCGR3B/CSF3R neutrophil-related
TPSB2/CPA3 mast-related

SPP1--CD44
TGFB1--TGFBR1_TGFBR2
IL10--IL10RA_IL10RB
PVR--TIGIT
LGALS9--HAVCR2
CD274--PDCD1
CD80--CD28
CXCL9--CXCR3
CCL2--CCR2
CSF1--CSF1R
TNF--TNFRSF1A
TNF--TNFRSF1B
TNFSF13B--TNFRSF13C
TNFSF13--TNFRSF17
CXCL12--CXCR4

Eligible edges with ligand + all receptor subunits (%)
100
80
60
40
20
0

15 fixed pairs from LIANA snapshot 1.10.0; no communication significance test. Grey: no eligible edges; yellow zero: edges without RNA support.
All-T includes T subgroups; subgroup gates and targets can overlap. A 100% value from 1-2 edges is not replicated evidence. Mixed-spot receptor attribution remains unresolved.