
PGAM5-high vs all other macrophage-RNA spots
Equal-section median adjusted effects; 4 independent patients
MHC-II antigen presentation
Inflammatory myeloid RNA
CCL / CXCL8 chemokines
CXCR3 ligands
Checkpoint ligands
IL10 / TGFB1 / LGALS9
Costimulation ligands
Scavenger / LPL state
C1Q complement (gate overlap)
Uptake proxy (gate overlap)
SPP1 / MMP12 remodeling
Hypoxia response
Cycling
+0.09 -0.10 +0.05 +0.19
+0.03 -0.03 -0.21 -0.19
-0.06 -0.15 +0.46 -0.27
+0.07 +0.07 -0.30 +0.02
+0.03 -0.05 +0.14 +0.19
-0.005 +0.22 +0.15 -0.12
-0.02 -0.21 -0.11 -0.08
-0.11 -0.003 -0.52 -0.15
+0.01 -0.17 -0.25 +0.10
+0.22 -0.22 +0.30 -0.02
+0.09 -0.02 -0.27 -0.01
-0.03 +0.10 -0.15 -0.17
+0.14 -0.04 +0.28 +0.04
A009
B013
C004
C008
0.8
0.6
0.4
0.2
0.0
−0.2
−0.4
−0.6
−0.8
Adjusted coefficient / within-section outcome SD