# 学术文献阅读

2020年9月5日

## 目录


- □文献获取
- □文献管理
- □文献阅读

## 文献获取

- 搜索关键词
- 人名/课题组
- 跟踪订阅
- 他人推荐
- 已有文献拓展


## 文献获取-关键词

### Google Scholar




## 文献获取-人名搜索

- Google scholar citation
- Homepage
- Researchgate







## 文献获取-订阅








检索 来源出版物 列表 SciVal »







#### 作者详情



## 文献获取-他人推荐

H.H. Heenen, J.A. Gauthier, H.H. Kristoffersen, T. Ludwig, and K. Chan, J. Chem. Phys. **152**, 144703 (2020).

# Google Scholar

I		Q
	Articles	

## 文献获取-已有文献拓展

#### References

- M. Pumera, Graphene-based nanomaterials for energy storage, Energy Environ. Sci. 4 (2011) 668–674, https://doi.org/10.1039/C0EE00295J.
- [2] N. Choi, Z. Chen, S.A. Freunberger, X. Ji, Y. Sun, K. Amine, G. Yushin, L.F. Nazar, J. Cho, P.G. Bruce, Challenges facing lithium batteries and electrical double-layer capacitors, Angew. Chem. 51 (2012) 9994–10024, https://doi.org/10.1002/anie. 201201429.
- [3] F. Shen, X. Chen, P. Gao, G. Chen, Electrochemical removal of fluoride ions from industrial wastewater, Chem. Eng. Sci. 58 (2003) 987–993, https://doi.org/10. 1016/S0009-2509(02)00639-5.
- [4] G. Chen, Electrochemical technologies in wastewater treatment, Separ. Purif. Technol. 38 (2004) 11–41, https://doi.org/10.1016/j.seppur.2003.10.006.
- [5] Y. Yavuz, A.S. Koparal, U.B. Ogutveren, Treatment of petroleum refinery wastewater by electrochemical methods, Desalination 258 (2010) 201–205, https://doi.org/10.1016/j.desal.2010.03.013.
- [6] N.R. Council, Research Opportunities in Corrosion Science and Engineering, The National Academies Press, Washington, DC, 2011https://doi.org/10.17226/ 13032.
- [7] Y. Liu, D. Gokcen, U. Bertocci, T.P. Moffat, Self-terminating growth of platinum films by electrochemical deposition, Science 338 (6112) (2012) 1327–1330, https://doi.org/10.1126/science.1228925.
- [8] K.T. Butler, G.S. Gautam, P. Canepa, Designing interfaces in energy materials applications with first-principles calculations, npj Comput. Mater. 5 (2019), https://doi.org/10.1038/s41524-019-0160-9.


## 其它

- 研之成理
- 材料人
- X-MOL
- 蔻享学术(学术报告视频)
- http://gen.lib.rus.ec/(英文专著)
- Sci-hub.tw (英文文献)


## 文献管理

- 整理文献
- 方便使用
- 引用和输出

## 文献管理器-Mendeley







but do diffuse away if they are described oxidatively from gold. field enhancement | buried interface | spectroelectrochemistry ctrochemistry is one of the most important disciplines in nodern science and industry, involved in thousands of ap-

most DC analytical tools lack the sensitivity to access molecular-

frequency vibrational spectroscopy and surface plasmon resonance

at EC interfaces, we demonstrate the feasibility of measuring in situ

and real-time vibrational spectra during IC reactions at noble metal

electrodes. Application of the technique to EC reactions at a gold surface helps in understanding how the surface in a basis solution

is coolised and reduced during a cyclic voltammetry cycle. Study of

description of a third self-assembled monologer from gold through

DC reactions in a basis solution shows that the described think by

reductive reaction remain as an ordered layer near the gold interface.

effections including butteries, electrobois, electromethesis, and

many others (1). I file all chemical reserious, the anamhigaous

Electrochemical (EQ) reactions are crucial in many applications, per due to the SF resonance (U.I., UZ). Thus, SF-enhanced SFVS may allow us to gain a comprehensive and molecular-level view of level information of reactants and products. By combining sum: EC interfaces


In this paper, we report our recent study on EC reactions at gold electrodes in agreeous solutions using SP-columned numfrequency obstational spectroscopy (SP/SFVS). The raw SP-SFVS spectra showed combined resonant characteristics of SP and SFVS, providing in sits and real-time information of EC on with and without molecular adsorbates. We found that the SP-SPVS signal followed the exidation and reduction procemes donoted by the CV of an EC cycle, and could help-our understanding of how, for example, a gold surface is oxidized and reduced during the EC cycle. Using alianothiol as an example, we were also able to study adsorption and desorption of an alkanethiol self-assembled monolayer (SAM) on gold manipulated by EC. Since its discovery in the 1990s, EC of thirds on gold has been extensively studied (13-20) for applications in muso technology and bioscience (21, 22). However, due to lack of in situ spectroscopic tools, its microscopic understanding remains

applications, yet most EC analytical tools lack the sensitivity to access notecular level information of reactants and products. By combring sum-frequency vibrational spectroscopy and surface plasmon. reconunce at IIC interfaces, we demonstrate the feedbility of measuring in altu and real-time vibrational spectra during EC reactions at noble metal-electrodes Application of the technique to EC reactions at a gold surface helps in understanding how the surface in a basic solution is oslitical and reduced during a cyclivolumentry cycle. Study of description of a thici self-essentialed monologie from gold through EC reactions in a basic solution shows that the described thinly by soluctive reaction remain as an ordered layer near the gold interface, but do diffuse every if they are descr.

Author Enywords

Add File...

## 文献管理器-Mendeley



https://www.mendeley.com/referencemanagement/web-importer#id 1

## 文献管理器-其它

- EndNote
- Papers3/Readcube
- Zotero

达到主要目的为主,方便使用和引用(Word 和 Latex)

## 文献阅读

- 粗读
- 精读
- 拓展阅读

## 粗读文献-Title & Journal

基本检索 被引参考文献检索 高级检索			
CO2 reduction	◎ 标题	*	检索检索提示
		+添加行   重设	
时间跨度 所有年份 (1950 - 2020) 更多设置 ▼			
排序方式: 日期 IF 被引频次 使用次数 相关性 更多▼			1 / 651 >
□ 选择页面 □ 导出 添加到标记结果列表			<b>听检索结果</b> 建引文报告
Optimizing electricity mix for CO2 emissions reduction: A 作者: Kang, Jidong; Ng, Tsan Sheng; Su, Bin	A robust input-output linear pro		E次: 0 所有数据库)
EUROPEAN JOURNAL OF OPERATIONAL RESEARCH 卷: 287 其	明: 1 页: 280-292 出版年: NOV 16	2020 使用2	で数~
<b>◇</b> 5-F-X 出版商处的全文 查看摘要▼			
Ethylene Selectivity in Electrocatalytic CO2 Reduction of Study	n Cu Nanomaterials: A Crystal P	hase-Dependent	
作者: Chen, Ye; Fan, Zhanxi; Wang, Jiong; 等.  JOURNAL OF THE AMERICAN CHEMICAL SOCIETY 卷: 142 期:	: 29 页: 12760-12766 出版年: JU	L 22 2020	

## 粗读文献-Journal

- 综合类: Nature, Science, 大子刊, Nature communication, PNAS,
- 化学类: JACS, Angew. Chem. Int. Ed., ACS C atalysis
- 物理化学: JPC系列, PCCP, JCP
- 电化学: Electrochimica Acta, Electrochemical communication
- 其它相关

## 文献阅读-文献组成

### 粗读

Abstract

Title

- Figures and Tables
- Introduction
- Conclusion

- Methods
- Results and Discussion

精读

- Reference
- **Support Information**
- Authors (& affiliation)

### 粗读文献-开始阅读



Research Article

pubs.acs.org/acscatalysis

# Insight into the Microenvironments of the Metal-Ionic Liquid Interface during Electrochemical CO<sub>2</sub> Reduction

Hyung-Kyu Lim,  $^{\dagger_{\odot}}$  Youngkook Kwon,  $^{\ddagger,\$}$  Han Seul Kim,  $^{\parallel}$  Jiwon Jeon,  $^{\parallel}$  Yong-Hoon Kim,  $^{\parallel_{\odot}}$  Jung-Ae Lim, Beom-Sik Kim,  $^{\ddagger,\$}$  Jina Choi,  $^{*,\ddagger}$  and Hyungjun Kim $^{*,\parallel,\perp_{\odot}}$ 

Supporting Information

Department of Chemical Engineering, Kangwon National University, Chuncheon, Gangwon-do 24341, Republic of Korea


<sup>&</sup>lt;sup>‡</sup>Carbon Resource Institute, Korea Research Institute of Chemical Technology, 141 Gajeong-ro, Yuseong-gu, Daejeon 34114, Republic of Korea

<sup>§</sup>Advanced Materials and Chemical Engineering, University of Science & Technology, 217 Gajeong-ro, Yuseong-gu, Daejeon 34113, Republic of Korea

Graduate School of EEWS, Korea Advanced Institute of Science and Technology, 291 Daehak-ro, Yuseong-gu, Daejeon 34141, Republic of Korea

<sup>&</sup>lt;sup>1</sup>Department of Chemistry, Korea Advanced Institute of Science and Technology, 291 Daehak-ro, Yuseong-gu, Daejeon 34141, Republic of Korea


## 粗读文献-Figures/Tables



✔ 从图像与表格中看与你的研究是否相关

### 粗读文献-Abstract

ABSTRACT: Recently, many experimental and theoretical efforts are being intensified to develop high-performance catalysts for electrochemical CO<sub>2</sub> conversion. Beyond the catalyst material screening, it is also critical to optimize the surrounding reaction medium. From vast experiments, inclusion of room-temperature ionic liquid (RTIL) in the electrolyte is found to be beneficial for CO<sub>2</sub> conversion; however, there is no unified picture of the role of RTIL, prohibiting further optimization of the reaction medium. Using a state-of-the-art multiscale simulation, we here unveil the atomic origin of the catalytic promotion effect of RTIL during CO<sub>2</sub> conversion. Unlike the conventional belief, which assumes a specific intermolecular coordination by the RTIL component, we find that the



promotion effect is collectively manifested by tuning the reaction microenvironment. This mechanism suggests the critical importance of the bulk properties (e.g., resistance, gas solubility and diffusivity, viscosity, etc.) over the detailed chemical variations of the RTIL components in designing the optimal electrolyte components, which is further supported by our experiments. This fundamental understanding of complex electrochemical interfaces will help in the development of more advanced electrochemical CO<sub>2</sub> conversion catalytic systems in the future.

KEYWORDS: reaction mechanism, electrocatalysis, ionic liquids, multiscale simulation, solid-liquid interface

### ✔ 从关键词和简介中推断研究内容与结论

## 粗读文献-Conclusions

#### CONCLUSION

On the basis of the atomistic details elucidated from our stateof-the-art simulations and systematic experiments, the catalytic promotion effect of RTILs is manifested by the collective participation of both cations and anions of RTILs. The RTIL at the EDL interface provides a suitable microenvironment for solvating the intermediate, which can be well preserved over the variations in chemical details because the solvation energy is dominated by the nonspecific interaction. We further anticipate that our comprehensive study of the nature of the complex electrochemical interface will be useful for designing enhanced electrochemical CO<sub>2</sub> reduction systems in the future. For example, the bulk properties of the reaction medium, such as the electrical resistance, gas solubility, gas diffusivity, and viscosity, are more critically optimized than a specific interaction pair. In addition, a certain amount of water is indispensable in facilitating the proton transfer by developing a percolated hydrogen bond network and improving the mass transport of RTIL components (Table S6) by screening the strong interionic electrostatic interaction.36

✓ 查看研究 结果是否 感兴趣

## 精读文献-Introduction背景

#### INTRODUCTION

There have been various technical explorations of the sustainable future of the global carbon-based economy. The development of CO<sub>2</sub> conversion technologies is imperative to successfully recycle the anthropogenically emitted greenhouse gas to produce future carbon resources of fuels and chemicals. Electrochemical CO<sub>2</sub> conversion is considered one of the most feasible and promising technologies because of its high reactivity under ambient conditions, its wide product window according to the type of catalyst, and its applicability in commercial processes. Thus, many experimental and theoretical studies have extensively developed high-performance electrochemical catalyst systems for CO<sub>2</sub> conversion to achieve a minimized overpotential, improved product selectivity, improved stability, etc. <sup>1-4</sup>

Beyond many efforts to optimize the catalyst material, 5-7 there was an interesting report in 2011 by Rosen et al.8 They

#### REFERENCES

- Ganesh, I. Conversion of Carbon Dioxide into Methanol a Potential Liquid Fuel: Fundamental Challenges and Opportunities (a Review). Renewable Sustainable Energy Rev. 2014, 31, 221–257.
- (2) Jhong, H. R.; Ma, S. C.; Kenis, P. J. A. Electrochemical Conversion of CO<sub>2</sub> to Useful Chemicals: Current Status, Remaining Challenges, and Future Opportunities. Curr. Opin. Chem. Eng. 2013, 2, 191–199.
- (3) Oloman, C.; Li, H. Electrochemical Processing of Carbon Dioxide. ChemSusChem 2008, 1, 385-391.

showed that the activity of the electrochemical CO<sub>2</sub>-to-CO conversion reaction can be remarkably enhanced by an electrolyte mixture of 18 mol% 1-ethyl-3-methylimidazolium tetrafluoroborate (EMIm-BF<sub>4</sub>), one of the room-temperature ionic liquids (RTILs), and 82 mol % water. After this seminal work, many experimental reports have shown that the addition of RTIL to an electrolyte generally promotes the CO<sub>2</sub> reduction activity regardless of the choice of catalyst material (Ag, Au, or Pt). Because the electrochemical catalytic reaction is manifested by the interfacial chemistry, which occurs at the heterogeneous interface between the catalyst surface and the liquid medium, this study successfully demonstrated the importance of a particular design of the reaction medium.

Revised: November 6, 2017 Revised: January 5, 2018 Published: February 6, 2018

(19) Lim, H. K.; Lee, H.; Kim, H. A Seamless Grid-Based Interface for Mean-Field QM/MM Coupled with Efficient Solvation Free Energy Calculations. J. Chem. Theory Comput. 2016, 12, 5088-5099.

(20) Nakano, H.; Yamamoto, T. Accurate and Efficient Treatment of Continuous Solute Charge Density in the Mean-Field QM/MM Free Energy Calculation. J. Chem. Theory Comput. 2013, 9, 188–203.

(21) Lin, S. T.; Blanco, M.; Goddard, W. A. The Two-Phase Model for Calculating Thermodynamic Properties of Liquids from Molecular Dynamics: Validation for the Phase Diagram of Lennard-Jones Fluids. J. Chem. Phys. 2003, 119, 11792—11805.

(22) Dimeter C Test Dentilel Alexanders for Chest Dense

## 精读文献-Introduction科学问题

However, there is little understanding of how RTIL participates in and promotes CO2 reduction chemistry, which limits further the systematic design of the catalyst system. Rosen et al. attributed the improved reactivity to a greater proton availability by the hydrolysis of BF<sub>4</sub><sup>-11</sup> and a layer of EMIm+ on the catalyst surface12 from follow-up experimental studies. In addition, the imidazole cation may directly chemically react with CO2 and serve as a proton donor, which promotes the proton-coupled electron transfer (PCET) process. 13,14 However, a recent experimental study has shown that methylation at the C2 position of imidazole promotes the CO2 reduction activity, which excludes the possibility of the direct catalytic role of a proton at the C2 position. 15 The experimental screening approach to identify the catalytic role of RTIL mostly relies on investigating the correlation between the catalytic activity and the chemical modifications of the RTIL. 15,16 Although this approach certainly provides valuable information about how the RTIL participates in the reaction, the concurrent change in the bulk properties of the RTIL after the chemical modifications adds many complications in deciphering the experimental observations. For example, methylation at the C2 position of 1-ethyl-3-methylimidazolium bis(trifluoromethylsulfonyl)imide (EMIm-TFSI) increases the melting temperature from -14.7 to 22.0 °C and changes the viscosity from 21 to 74 mPa s.17 As summarized in our recent review, 18 there have been various efforts to identify mechanisms through various experimental and theoretical studies, but no decisive conclusion has been drawn.

▶ 作者总结的领域内关键科学问题

### 具体解决方法-最后一段

In this study, we unveil the atomic origin of the catalytic promotion effect of RTILs during electrochemical CO2 conversion by means of the first-principles-based multiscale simulation method (DFT-CES). In contrast to the previous conception that a certain type of direct chemical interaction between the RTIL component and the reaction intermediate promotes catalytic activity (which has often been assumed in the interpretation of experimental results), we find that RTIL components participate in the reaction in a collective manner by adjusting the reaction microenvironment during CO2 reduction. On the basis of our mechanism, we designed and performed a series of systematic experiments. This demonstrates the importance of bulk properties such as a bulk conductivity to the detailed chemical interactions of RTIL, suggesting a new design guideline for the reaction medium to enhance CO2 reduction.

## 精读文献-Methods

#### METHODS

Multiscale Simulation Method (DFT-CES). To briefly explain the key aspects of the DFT-CES method for the

Computational Details. First, we built four types of solvent models with different molar ratios of EMIm-BF<sub>4</sub> to water: from pure water (0:100) to pure ionic liquid (100:0) (Table S1). We used the LAMMPS (Large-scale Atomic/

**Electrochemical Measurements.** Electrochemical experiments were performed in a three-electrode system. A silver foil (Alfa Acer, 99.998%) was polished before each measurement

✔ 厘清所用方法,根据自己的研究目的决定理解程度。

## 精读文献-Results and Discussion

- 线性讲述
- 段内第一句
- 结合图表快速定位
- Support Information

#### RESULTS AND DISCUSSION

Reaction Thermodynamics. Using the DFT-CES/2PT method, we investigate the reaction thermodynamics for the CO<sub>2</sub>-to-CO reduction on the Ag(111) surface when the electrolyte phase is water or an EMIm-BF<sub>4</sub>/water mixture (20:80 molar ratio). The simulation cell in Figure S1 includes

Water Molecule Coordinating \*COOH. Surprisingly, we find that the local coordinating environment of \*COOH is

Collective Change in the Reaction Microenvironment. Beyond the comparison of the direct coordinating environ-

#### 4. RESULTS AND DISCUSSION

- **4.1. Convergence of SCRF Iterations.** To check the convergence of DFT-CES, we monitored how  $E_{\text{reorg}}$  from DFT calculations changed over the SCRF iterations. As shown in
- **4.2. Reaction Field Sampled from MD.** After running enough SCRF iterations to converge  $E_{\rm reorge}$  we visualized the reaction field of  $\langle \rho_{\rm solvent} \rangle$  in Figure 3 for two representative

## 精读文献-Authors

# Insight into the Microenvironments of the Metal-Ionic Liquid Interface during Electrochemical CO<sub>2</sub> Reduction

Hyung-Kyu Lim, †© Youngkook Kwon, ‡,§ Han Seul Kim, I Jiwon Jeon, Yong-Hoon Kim, © Jung-Ae Lim, Beom-Sik Kim, Jina Choi, \*\* and Hyungjun Kim\*, I Jiwon Jeon, Yong-Hoon Kim, I Jung-Ae Lim, Beom-Sik Kim, I Jina Choi, \*\* and Hyungjun Kim\*, I Jiwon Jeon, Yong-Hoon Kim, I Jiwon Jeon, I J

#### AUTHOR INFORMATION

#### **Corresponding Authors**


\*Phone: +82-42-860-7396. E-mail: jchoi@krict.re.kr. \*Phone: +82-42-350-1725. E-mail: linus16@kaist.ac.kr.

#### REFERENCES

- (18) Lim, H.-K.; Kim, H. The Mechanism of Room-Temperature Ionic-Liquid-Based Electrochemical CO<sub>2</sub> Reduction: A Review. *Molecules* **2017**, 22, 536.
- (19) Lim, H. K.; Lee, H.; Kim, H. A Seamless Grid-Based Interface for Mean-Field QM/MM Coupled with Efficient Solvation Free Energy Calculations. *J. Chem. Theory Comput.* **2016**, *12*, 5088–5099.
- (36) Jeon, J.; Kim, H.; Goddard, W. A.; Pascal, T. A.; Lee, G. I.; Kang, J. K. The Role of Confined Water in Ionic Liquid Electrolytes for Dye-Sensitized Solar Cells. *J. Phys. Chem. Lett.* **2012**, *3*, 556–559.

## 文献阅读

- ✔ 大部分文献粗读,少量精度(保证文章阅读量);
- ✔ 建议一篇文献一次性看完,即使有很多不懂的地方;
- ✓ 结合自己的数据,隔断时间多次阅读核心的文献,可获得灵感;
- ✓ 通过多种工具,跟踪最新的文献;
- ✓ 定期进行文献总结,综述,厘清目前研究的不足之处 及我们可弥补的地方。



Nothing makes you feel stupid quite like reading a scientific journal article.