

Assignment Assignment-03 due 10/08/2022 at 11:59pm PDT

Problem 1. (2 points)

A game of chance involves rolling an unevenly balanced 4-sided die. The probability that a roll comes up 1 is 0.16, the probability that a roll comes up 1 or 2 is 0.43, and the probability that a roll comes up 2 or 3 is 0.45 . If you win the amount that appears on the die, what is your expected winnings? (Note that the die has 4 sides.)

Answer = _____ dollars.

Answer(s) submitted:

- 2.8

(correct)

Correct Answers:

- 2.8

Problem 2. (3 points)

The mean and standard deviation of a random variable x are -8 and 1 respectively. Find the mean and standard deviation of the given random variables:

(1) $y = x + 4$

$\mu = \underline{\hspace{1cm}}$

$\sigma = \underline{\hspace{1cm}}$

(2) $v = 3x$

$\mu = \underline{\hspace{1cm}}$

$\sigma = \underline{\hspace{1cm}}$

(3) $w = 3x + 4$

$\mu = \underline{\hspace{1cm}}$

$\sigma = \underline{\hspace{1cm}}$

Answer(s) submitted:

- -4
- 1
- -24
- 3
- -20
- 3

(correct)

Correct Answers:

- -4
- 1
- -24
- 3
- -20
- 3

Problem 3. (4 points)

A robot fires 3 shots at a moving target. For the first shot, the probability of hitting the moving target is $1/3$. For subsequent shots beyond the first shot, the probability of hitting the moving target is $1/2$ if the previous shot is a hit (for example, the probability of hitting the moving target on the 3rd shot is $1/2$ if the 2nd shot is a hit) and the probability of hitting the moving target is $1/4$ if the previous shot is a miss.

What is the mean and variance of the number of hits?

Mean (rounded to the nearest whole number): ____

Variance (correct to 2 decimals): ____

Solution:

Mean: We list out all 8 possible outcomes with their probabilities as shown below.

Sequence	No. Hits	Probability
hit/hit/hit	3	$\frac{1}{3} \times \frac{1}{2} \times \frac{1}{2} = \frac{4}{48}$
hit/hit/miss	2	$\frac{1}{3} \times \frac{1}{2} \times \frac{1}{2} = \frac{4}{48}$
hit/miss/hit	2	$\frac{1}{3} \times \frac{1}{2} \times \frac{1}{4} = \frac{2}{48}$
hit/miss/miss	1	$\frac{1}{3} \times \frac{1}{2} \times \frac{3}{4} = \frac{6}{48}$
miss/hit/hit	2	$\frac{2}{3} \times \frac{1}{4} \times \frac{1}{2} = \frac{4}{48}$
miss/hit/miss	1	$\frac{2}{3} \times \frac{1}{4} \times \frac{1}{2} = \frac{4}{48}$
miss/miss/hit	1	$\frac{2}{3} \times \frac{3}{4} \times \frac{1}{4} = \frac{6}{48}$
miss/miss/miss	0	$\frac{2}{3} \times \frac{3}{4} \times \frac{3}{4} = \frac{18}{48}$

We then compute the probabilities for each value of X by adding the appropriate entries in the table above.

X	0	1	2	3
Probability	$18/48 = 3/8$	$16/48 = 1/3$	$10/48 = 5/24$	$4/48 = 1/12$

Finally, we find $E(X) = \sum_{i=0}^3 iP(X=i) = 1$.

Variance: by computing $E(X^2) = \frac{3}{8} \times 0^2 + \frac{1}{3} \times 1^2 + \frac{5}{24} \times 2^2 + \frac{1}{12} \times 3^2$ using the probabilities above, we find the variance by $Var(X) = E(X^2) - E(X)^2 = 0.92$.

Answer(s) submitted:

- 1
- 0.92

(correct)

Correct Answers:

- 1
- 0.92

Problem 4. (2 points)

The following density function describes a random variable X .

$$f(x) = \frac{x-1}{8} \text{ if } 1 < x < 5$$

A. Find the probability that X lies between 2 and 4.

Probability = _____

B. Find the probability that X is less than 3.

Probability = _____

Answer(s) submitted:

- 0.5
- 0.25

(correct)

Correct Answers:

- 0.5
- 0.25

Problem 5. (3 points)

The time that a butterfly lives after emerging from its chrysalis can be modelled by a random variable T , the model here taking the probability that a butterfly survives for more than t days as

$$P(T > t) = \frac{36}{(6+t)^2}, \quad t \geq 0.$$

For these problems, please ensure your answers are accurate to within 3 decimals.

Part a)

What is the probability that a butterfly will die within 6 days of emerging?

—

Part b)

If a large number of butterflies emerge on the same day, after how many days would you expect only 6 % to be alive?

—

Part c)

Calculate the mean lifetime of a butterfly after emerging from its chrysalis.

—

Solution:**Part a)**

We require

Part b)

Now we wish to find t such that $P(T > t) = 0.06$ (i.e. the lifetime exceeds t with probability 6 %). We solve

$$\frac{36}{(6+t)^2} = 0.06,$$

that is,

$$(6+t)^2 = 600.000.$$

Solving this quadratic and taking the positive root, we find $t = 18.495$.

Part c)

The c.d.f. satisfies

$$F(t) = P(T \leq t) = 1 - P(T > t)$$

and so

$$F(t) = \begin{cases} 0 & \text{if } t < 0 \\ 1 - \frac{36}{(6+t)^2} & \text{if } t \geq 0. \end{cases}$$

Hence the p.d.f. can be found by differentiation,

and so

$$f(t) = \begin{cases} 0 & \text{if } t < 0 \\ \frac{72}{(6+t)^3} & \text{if } t \geq 0. \end{cases}$$

Hence, the expectation is given by

$$E(T) = \int_0^{\infty} \frac{72t}{(6+t)^2} dt.$$

Putting $x = 6 + t$, the integral above becomes

$$\int_6^{\infty} \frac{72(x-6)}{x^3} dx = \int_6^{\infty} \left(\frac{72}{x^2} - \frac{432}{x^3} \right) dx = 6.$$

Answer(s) submitted:

- 0.75
- 18.495
- 6

(correct)

Correct Answers:

- 0.75
- 18.495
- 6

Problem 6. (6 points)

The continuous random variable X has cumulative distribution function given by

$$F(x) = \begin{cases} 0 & \text{for } x < 0 \\ x - \frac{1}{4}x^2 & \text{for } 0 \leq x \leq 2 \\ 1 & \text{for } x > 2 \end{cases}$$

Part(a) Find $P(X > \frac{1}{2})$. Give your answer as a decimal, correct to 2 decimal places. ____

Part(b) Two independent observations of X are taken. Find the probability correct to 2 decimal places that one is less than $\frac{1}{2}$ and the other is greater than $\frac{1}{2}$. The order in which we take the observations matters. ____

Part(c) Find the median of X , correct to 2 decimal places. ____

Part(d) Find $E(\sqrt{X})$, correct to 2 decimal places. ____

Part(e) Find the value of q such that $P(X < q) = \frac{1}{4}$. Give your answer as a decimal correct to 3 decimal places. ____

Part(f) Find the value of c correct to one decimal place given that $E(X + c) = 4E(X - c)$. ____

Answer(s) submitted:

- 0.56
- 0.49
- 0.59
- 0.75
- 0.268
- 0.4

(correct)

Correct Answers:

- 0.56
- 0.49
- 0.59
- 0.75
- 0.268
- 0.4