

Chapter 7

- 7.1 Measures of predictive accuracy
- 7.2 Information criteria and cross-validation
 - Instead of 7.2, read:
Vehtari, A., Gelman, A., Gabry, J. (2017). Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC. *Statistics and Computing*. 27(5):1413–1432. [arXiv preprint](#).
- 7.3 Model comparison based on predictive performance
- 7.4 Model comparison using Bayes factors
- 7.5 Continuous model expansion / sensitivity analysis
- 7.5 Example (may be skipped)


Model assesment, selection and inference after selection

- Extra material at <https://avehtari.github.io/modelselection/>
 - Videos, Slides, Notebooks, References
 - The most relevant for the course is the first part of the talk “Model assesment, comparison and selection at Master class in Bayesian statistics, CIRM, Marseille”


Model comparison

- “A popular hypothesis has it that primates with larger brains produce more energetic milk, so that brains can grow quickly” (from Statistical Rethinking)
 - Model 1: formula = kcal.per.g \sim neocortex
 - Model 2: formula = kcal.per.g \sim neocortex + log(mass)

Pointwise comparison LOO models: Model 1




Pointwise comparison LOO models: Model 1



Model 1 $\text{elpd_loo} \approx 3.7$, $\text{SE}=1.8$

Model 2 $\text{elpd_loo} \approx 8.4$, $\text{SE}=2.8$


Pointwise comparison LOO models: Model 1



Model 1 $\text{elpd_loo} \approx 3.7$, $\text{SE}=1.8$

Model 2 $\text{elpd_loo} \approx 8.4$, $\text{SE}=2.8$

Pointwise comparison LOO models




Model comparison:

(negative 'elpd_diff' favors 1st model, positive favors 2nd)

elpd_diff	se
4.7	2.7

Arsenic well example – Model comparison




An estimated difference in elpd_{loo} of 16.4 with SE of 4.4.

see [Vehtari, Gelman & Gabry \(2017a\)](#)

Sometimes cross-validation is not needed

- Posterior predictive checking is often sufficient



Predicting the yields of mesquite bushes.

Gelman, Hill & Vehtari (2019): Regression and Other Stories, Chapter 11.

- BDA3, Chapter 6
- Gabry, Simpson, Vehtari, Betancourt, Gelman (2018). Visualization in Bayesian workflow. JRSS A, [preprint arXiv:1709.01449](https://arxiv.org/abs/1709.01449)
- mc-stan.org/bayesplot/articles/graphical-ppcs.html
- betanalpha.github.io/assets/case_studies/principled_bayesian_workflow.html

Sometimes cross-validation is not needed

- For some very simple cases you may assume that true model is included in the list of models considered (M -closed)

Sometimes cross-validation is not needed

- For some very simple cases you may assume that true model is included in the list of models considered (M -closed)
 - see predictive model selection in M -closed case by San Martini and Spezzaferri (1984)


Sometimes cross-validation is not needed

- For some very simple cases you may assume that true model is included in the list of models considered (M -closed)
 - see predictive model selection in M -closed case by San Martini and Spezzaferri (1984)
 - but you should not force your design of experiment or analysis to stay in the simplified world

Sometimes cross-validation is not needed

- For some very simple cases you may assume that true model is included in the list of models considered (M -closed)
 - see predictive model selection in M -closed case by San Martini and Spezzaferri (1984)
 - but you should not force your design of experiment or analysis to stay in the simplified world
- In nested case, often easier and more accurate to analyse posterior distribution of more complex model directly
avehtari.github.io/modelselection/betablockers.html

Sometimes predictive model comparison can be useful




Marginal posterior intervals

Sometimes predictive model comparison can be useful




Marginal posterior intervals




Joint posterior density

`rstanarm` + `bayesplot`

Sometimes predictive model comparison can be useful



Marginal posterior intervals



Joint posterior density

`rstanarm` + `bayesplot`

see also [Collinear demo](#)

What if one is not clearly better than others?

What if one is not clearly better than others?

- Continuous expansion including all models?
 - and then analyse the posterior distribution directly
avehtari.github.io/modelselection/betablockers.html
 - sparse priors like regularized horseshoe prior instead of variable selection
video, refs and demos at avehtari.github.io/modelselection/

What if one is not clearly better than others?

- Continuous expansion including all models?
 - and then analyse the posterior distribution directly
avehtari.github.io/modelselection/betablockers.html
 - sparse priors like regularized horseshoe prior instead of variable selection
video, refs and demos at avehtari.github.io/modelselection/
- Model averaging with BMA or Bayesian stacking?
mc-stan.org/loo/articles/loo2-example.html

What if one is not clearly better than others?

- Continuous expansion including all models?
 - and then analyse the posterior distribution directly
avehtari.github.io/modelselection/betablockers.html
 - sparse priors like regularized horseshoe prior instead of variable selection
video, refs and demos at avehtari.github.io/modelselection/
- Model averaging with BMA or Bayesian stacking?
mc-stan.org/loo/articles/loo2-example.html
- In a nested case choose simpler if assuming some cost for extra parts?
andrewgelman.com/2018/07/26/parsimonious-principle-vs-integration-uncertainties/

What if one is not clearly better than others?

- Continuous expansion including all models?
 - and then analyse the posterior distribution directly
avehtari.github.io/modelselection/betablockers.html
 - sparse priors like regularized horseshoe prior instead of variable selection
video, refs and demos at avehtari.github.io/modelselection/
- Model averaging with BMA or Bayesian stacking?
mc-stan.org/loo/articles/loo2-example.html
- In a nested case choose simpler if assuming some cost for extra parts?
andrewgelman.com/2018/07/26/parsimonious-principle-vs-integration-uncertainties/
- In a nested case choose more complex if you want to take into account all the uncertainties.
andrewgelman.com/2018/07/26/parsimonious-principle-vs-integration-uncertainties/

Model averaging

- Prefer continuous model expansion

Model averaging

- Prefer continuous model expansion
- If needed integrate over the model space = model averaging

Model averaging

- Prefer continuous model expansion
- If needed integrate over the model space = model averaging
- Bayesian stacking may work better than BMA
 - Yao, Vehtari, Simpson, & Gelman (2018)

Cross-validation and model selection

- Cross-validation can be used for model selection if
 - small number of models
 - the difference between models is clear

Cross-validation and model selection

- Cross-validation can be used for model selection if
 - small number of models
 - the difference between models is clear
- Do not use cross-validation to choose from a large set of models
 - selection process leads to overfitting

Cross-validation and model selection

- Cross-validation can be used for model selection if
 - small number of models
 - the difference between models is clear
- Do not use cross-validation to choose from a large set of models
 - selection process leads to overfitting
- Overfitting in selection process is not unique for cross-validation

Selection induced bias and overfitting

- Selection induced bias in cross-validation
 - same data is used to assess the performance and make the selection
 - the selected model fits more to the data
 - the CV estimate for the selected model is biased
 - recognised already, e.g., by Stone (1974)


Selection induced bias and overfitting

- Selection induced bias in cross-validation
 - same data is used to assess the performance and make the selection
 - the selected model fits more to the data
 - the CV estimate for the selected model is biased
 - recognised already, e.g., by Stone (1974)
- Performance of the selection process itself can be assessed using two level cross-validation, but it does not help choosing better models


Selection induced bias and overfitting

- Selection induced bias in cross-validation
 - same data is used to assess the performance and make the selection
 - the selected model fits more to the data
 - the CV estimate for the selected model is biased
 - recognised already, e.g., by Stone (1974)
- Performance of the selection process itself can be assessed using two level cross-validation, but it does not help choosing better models
- Bigger problem if there is a large number of models as in covariate selection

Selection induced bias in variable selection




Selection induced bias in variable selection




Piironen & Vehtari (2017)

Selection induced bias in variable selection



Piironen & Vehtari (2017)

Selection induced bias in variable selection



Piironen &
Vehtari (2017)

Take-home messages

- It's good to think predictions of observables, because observables are the only ones we can observe
- Cross-validation can simulate predicting and observing new data
- Cross-validation is good if you don't trust your model
- Different variants of cross-validation are useful in different scenarios
- Cross-validation has high variance, and **if** you trust your model you can beat cross-validation in accuracy

Take-home messages

- It's good to think predictions of observables, because observables are the only ones we can observe
- Cross-validation can simulate predicting and observing new data
- Cross-validation is good if you don't trust your model
- Different variants of cross-validation are useful in different scenarios
- Cross-validation has high variance, and **if** you trust your model you can beat cross-validation in accuracy

Take-home messages

- It's good to think predictions of observables, because observables are the only ones we can observe
- Cross-validation can simulate predicting and observing new data
- Cross-validation is good if you don't trust your model
- Different variants of cross-validation are useful in different scenarios
- Cross-validation has high variance, and **if** you trust your model you can beat cross-validation in accuracy

Take-home messages

- It's good to think predictions of observables, because observables are the only ones we can observe
- Cross-validation can simulate predicting and observing new data
- Cross-validation is good if you don't trust your model
- Different variants of cross-validation are useful in different scenarios
- Cross-validation has high variance, and **if** you trust your model you can beat cross-validation in accuracy

Take-home messages

- It's good to think predictions of observables, because observables are the only ones we can observe
- Cross-validation can simulate predicting and observing new data
- Cross-validation is good if you don't trust your model
- Different variants of cross-validation are useful in different scenarios
- Cross-validation has high variance, and **if** you trust your model you can beat cross-validation in accuracy

Rich model vs feature selection?

- If we care only about the predictive performance
 - Include all available prior information
 - Integrate over all uncertainties
 - No need for feature selection

Rich model vs feature selection?

- If we care only about the predictive performance
 - Include all available prior information
 - Integrate over all uncertainties
 - No need for feature selection
- Variable selection can be useful if
 - need to reduce measurement or computation cost in the future
 - improve explainability

Rich model vs feature selection?

- If we care only about the predictive performance
 - Include all available prior information
 - Integrate over all uncertainties
 - No need for feature selection
- Variable selection can be useful if
 - need to reduce measurement or computation cost in the future
 - improve explainability
- Two options for variable selection
 - Find a minimal subset of features that yield a good predictive model
 - Identify all features that have predictive information

Why shrinkage priors alone do not solve the variable selection problem

- A common strategy:
 - Fit model with a shrinkage prior
 - Select variables based on marginal posteriors (of the regression coefficients)

Why shrinkage priors alone do not solve the variable selection problem

- A common strategy:
 - Fit model with a shrinkage prior
 - Select variables based on marginal posteriors (of the regression coefficients)
- Problems
 - Marginal posteriors are difficult with correlated features
 - How to do post-selection inference correctly?

Example

Consider data

$$f \sim \mathbf{N}(0, 1),$$

$$y \mid f \sim \mathbf{N}(f, 1)$$

$$x_j \mid f \sim \mathbf{N}(\sqrt{\rho}f, 1 - \rho), \quad j = 1, \dots, 25,$$

$$x_j \mid f \sim \mathbf{N}(0, 1), \quad j = 26, \dots, 50.$$

Example

Consider data

$$f \sim \mathcal{N}(0, 1),$$

$$y \mid f \sim \mathcal{N}(f, 1)$$

$$x_j \mid f \sim \mathcal{N}(\sqrt{\rho}f, 1 - \rho), \quad j = 1, \dots, 25,$$

$$x_j \mid f \sim \mathcal{N}(0, 1), \quad j = 26, \dots, 50.$$

- y are noisy observations about latent f

Example

Consider data

$$f \sim \mathcal{N}(0, 1),$$

$$y \mid f \sim \mathcal{N}(f, 1)$$

$$x_j \mid f \sim \mathcal{N}(\sqrt{\rho}f, 1 - \rho), \quad j = 1, \dots, 25,$$

$$x_j \mid f \sim \mathcal{N}(0, 1), \quad j = 26, \dots, 50.$$

- y are noisy observations about latent f
- First $p_{\text{rel}} = 25$ features are correlated with ρ and predictive about y

Example

Consider data

$$\begin{aligned}f &\sim \mathcal{N}(0, 1), \\y \mid f &\sim \mathcal{N}(f, 1) \\x_j \mid f &\sim \mathcal{N}(\sqrt{\rho}f, 1 - \rho), & j = 1, \dots, 25, \\x_j \mid f &\sim \mathcal{N}(0, 1), & j = 26, \dots, 50.\end{aligned}$$

- y are noisy observations about latent f
- First $p_{\text{rel}} = 25$ features are correlated with ρ and predictive about y
- Remaining 25 features are irrelevant random noise

Example


Consider data

$$\begin{aligned}f &\sim \mathcal{N}(0, 1), \\y \mid f &\sim \mathcal{N}(f, 1) \\x_j \mid f &\sim \mathcal{N}(\sqrt{\rho}f, 1 - \rho), & j = 1, \dots, 25, \\x_j \mid f &\sim \mathcal{N}(0, 1), & j = 26, \dots, 50.\end{aligned}$$

- y are noisy observations about latent f
- First $p_{\text{rel}} = 25$ features are correlated with ρ and predictive about y
- Remaining 25 features are irrelevant random noise


Generate one data set $\{x^{(i)}, y^{(i)}\}_{i=1}^n$ with $n = 50$ and $\rho = 0.8$ and assess the feature relevances

Example




A) Gaussian prior, posterior median with 50% and 90% intervals

Example




- A) Gaussian prior, posterior median with 50% and 90% intervals
B) Horseshoe prior, same things

Example



- A) Gaussian prior, posterior median with 50% and 90% intervals
- B) Horseshoe prior, same things
- C) Spike-and-slab prior, posterior inclusion probabilities



Example



- A) Gaussian prior, posterior median with 50% and 90% intervals
- B) Horseshoe prior, same things
- C) Spike-and-slab prior, posterior inclusion probabilities

Half of the features relevant, but all marginals substantially overlapping with zero



What happens?





$$p_{\text{rel}} = 2$$

What happens?

Gaussian



Horseshoe






$p_{\text{rel}} = 2$




$p_{\text{rel}} = 5$

What happens?

Gaussian



Horseshoe



$\rho_{\text{rel}} = 2$

$\rho_{\text{rel}} = 5$

$\rho_{\text{rel}} = 25$

Focus on predictive performance

- Two stage approach
 - Construct a best predictive model you can
⇒ *reference model*
 - Variable selection and post-selection inference
⇒ *projection*


Focus on predictive performance

- Two stage approach
 - Construct a best predictive model you can
⇒ *reference model*
 - Variable selection and post-selection inference
⇒ *projection*
- Instead of looking at the marginals, find the minimal subset of features which have (almost) the same predictive performance as the reference model

Reference model improves variable selection

Same data generating mechanism, but


$n = 30$, $p = 500$, $p_{\text{rel}} = 150$, $\rho = 0.5$.



irrelevant x_j , relevant x_j

Sample correlation with y


Reference model improves variable selection



irrelevant x_j , relevant x_j

A) Sample correlation with y vs. sample correlation with f

Reference model improves variable selection



irrelevant x_j , relevant x_j

A) Sample correlation with y vs. sample correlation with f

B) Sample correlation with y vs. sample correlation with f_*

f_* = linear regression fit with 3 supervised principal components

(Iterative) Supervised Principal Components

- Dimension reduction for high dimensional small data with highly correlating features
 - dimension reduction helps to speed up later computation without discarding much information
 - supervised means that features correlating with the target are favored in constructing the principal components
- Piironen and Vehtari (2018). Iterative supervised principal components. 21st AISTATS, PMLR 84:106-114. [Online](#).

Predictive projection, idea

- Model simplification technique

Predictive projection, idea

- Model simplification technique
- Replace full posterior $p(\theta \mid D)$ with some constrained $q(\theta)$ so that the predictive distribution changes as little as possible

Predictive projection, idea

- Model simplification technique
- Replace full posterior $p(\theta \mid D)$ with some constrained $q(\theta)$ so that the predictive distribution changes as little as possible
- Example constraints
 - $q(\theta)$ can have only point mass at some θ_0
 \Rightarrow “Optimal point estimates”


Predictive projection, idea

- Model simplification technique
- Replace full posterior $p(\theta \mid D)$ with some constrained $q(\theta)$ so that the predictive distribution changes as little as possible
- Example constraints
 - $q(\theta)$ can have only point mass at some θ_0
 \Rightarrow “Optimal point estimates”
 - Some features must have exactly zero regression coefficient
 \Rightarrow “Which features can be discarded”

Predictive projection, idea



- Model simplification technique
- Replace full posterior $p(\theta \mid D)$ with some constrained $q(\theta)$ so that the predictive distribution changes as little as possible
- Example constraints
 - $q(\theta)$ can have only point mass at some θ_0
 \Rightarrow “Optimal point estimates”
 - Some features must have exactly zero regression coefficient
 \Rightarrow “Which features can be discarded”
- The decision theoretic idea of conditioning the smaller model inference on the full model can be tracked to Lindley (1968)
 - draw by draw projection introduced by Goutis & Robert (1998), and Dupuis & Robert (2003)
 - see also many related references in a review by Vehtari & Ojanen (2012)

Logistic regression with two features




Full posterior for β_1 and β_2 and contours of predicted class probability

Logistic regression with two features




Projected point estimates for β_1 and β_2

Logistic regression with two features




Projected point estimates, constraint $\beta_1 = 0$

Logistic regression with two features




Projected point estimates, constraint $\beta_2 = 0$

Logistic regression with two features



Draw-by-draw projection, constraint $\beta_1 = 0$

Logistic regression with two features



Draw-by-draw projection, constraint $\beta_2 = 0$

Predictive projection

- Replace full posterior $p(\theta \mid D)$ with some constrained $q(\theta)$ so that the predictive distribution changes as little as possible

Predictive projection

- Replace full posterior $p(\theta \mid D)$ with some constrained $q(\theta)$ so that the predictive distribution changes as little as possible
- As the full posterior $p(\theta \mid D)$ is projected to $q(\theta)$
 - the prior is also projected and there is no need to define priors for submodels separately

Predictive projection

- Replace full posterior $p(\theta \mid D)$ with some constrained $q(\theta)$ so that the predictive distribution changes as little as possible
- As the full posterior $p(\theta \mid D)$ is projected to $q(\theta)$
 - the prior is also projected and there is no need to define priors for submodels separately
 - even if we constrain some coefficients to be 0, the predictive inference is conditioned on the information related features contributed to the reference model

Projective selection

- How to select a feature combination?

Projective selection

- How to select a feature combination?
- For a given model size, choose feature combination with minimal projective loss


Projective selection

- How to select a feature combination?
- For a given model size, choose feature combination with minimal projective loss
- Search heuristics, e.g.
 - Monte Carlo search
 - Forward search
 - L_1 -penalization (as in Lasso)

Projective selection


- How to select a feature combination?
- For a given model size, choose feature combination with minimal projective loss
- Search heuristics, e.g.
 - Monte Carlo search
 - Forward search
 - L_1 -penalization (as in Lasso)
- Use cross-validation to select the appropriate model size
 - need to cross-validate over the search paths

Selection induced bias in variable selection



Piironen & Vehtari (2017)

Selection induced bias in variable selection




Piironen &
Vehtari (2017)

Bodyfat: small p example of projection predictive

Predict bodyfat percentage. The reference value is obtained by immersing person in water. $n = 251$.


Bodyfat: small p example of projection predictive

Predict bodyfat percentage. The reference value is obtained by immersing person in water. $n = 251$.




Bodyfat

Marginal posteriors of coefficients




Bodyfat

Bivariate marginal of weight and height




Bodyfat

The predictive performance of the full and submodels




Bodyfat

Marginals of projected posterior




Bodyfat

Projected posterior is not just the conditional of joint




Bodyfat

Projected posterior is different than posterior conditioned only on selected features




Projection of Gaussian graphical models


- Williams, Piironen, Vehtari, Rast (2018). Bayesian estimation of Gaussian graphical models with projection predictive selection. [arXiv:1801.05725](https://arxiv.org/abs/1801.05725)




BGL




GL




TIGER



BMA



MAP



Projection

CEU genetic network. BGL: Bayesian glasso; GL: glasso; TIGER: tuning insensitive graph estimation and regression; BMA: Bayesian model averaging; MAP: Maximum a posteriori; Projection: projection predictive selection.

More results

- More results projpred vs. Lasso and elastic net:
Piironen, Paasiniemi, Vehtari (2018). Projective Inference in High-dimensional Problems: Prediction and Feature Selection. [arXiv:1810.02406](#)
- More results projpred vs. marginal posterior probabilities:
Piironen and Vehtari (2017). Comparison of Bayesian predictive methods for model selection. Statistics and Computing, 27(3):711-735. [doi:10.1007/s11222-016-9649-y](#).
- projpred for Gaussian graphical models:
Williams, Piironen, Vehtari, Rast (2018). Bayesian estimation of Gaussian graphical models with projection predictive selection. [arXiv:1801.05725](#)
- More results for Bayes SPC:
Piironen and Vehtari (2018). Iterative supervised principal components. 21st AISTATS, PMLR 84:106-114. [Online](#).
- Several case studies for small to moderate dimensional ($p = 4 \dots 100$) small data:
Vehtari (2018). Model assesment, selection and inference after selection. <https://avehtari.github.io/modelselection/>

Take-home messages (part 2)

- Sparse priors do not automate variable selection
 - Don't trust marginal posteriors

Take-home messages (part 2)

- Sparse priors do not automate variable selection
 - Don't trust marginal posteriors
- Reference model + projection can improve feature selection
 - Excellent tradeoff between accuracy and model complexity
 - Useful also for identifying all the relevant features

Take-home messages (part 2)

- Sparse priors do not automate variable selection
 - Don't trust marginal posteriors
- Reference model + projection can improve feature selection
 - Excellent tradeoff between accuracy and model complexity
 - Useful also for identifying all the relevant features
- Well developed for GLMs, but can be used also with other model families

Take-home messages (part 2)

- Sparse priors do not automate variable selection
 - Don't trust marginal posteriors
- Reference model + projection can improve feature selection
 - Excellent tradeoff between accuracy and model complexity
 - Useful also for identifying all the relevant features
- Well developed for GLMs, but can be used also with other model families
- More details and results (+ some theoretical discussion) in the paper
 - Piironen, Paasiniemi, Vehtari (2018). Projective Inference in High-dimensional Problems: Prediction and Feature Selection. [arXiv:1810.02406](https://arxiv.org/abs/1810.02406)

Take-home messages (part 2)

- Sparse priors do not automate variable selection
 - Don't trust marginal posteriors
- Reference model + projection can improve feature selection
 - Excellent tradeoff between accuracy and model complexity
 - Useful also for identifying all the relevant features
- Well developed for GLMs, but can be used also with other model families
- More details and results (+ some theoretical discussion) in the paper
 - Piironen, Paasiniemi, Vehtari (2018). Projective Inference in High-dimensional Problems: Prediction and Feature Selection. [arXiv:1810.02406](https://arxiv.org/abs/1810.02406)
- R-package `projpred` in CRAN and github
<https://github.com/stan-dev/projpred>
(easy to use, e.g. with RStan, RStanARM, brms)

References

References and more at avehtari.github.io/masterclass/ and avehtari.github.io/modelselection/

- Model selection tutorial at StanCon 2018 Asilomar
 - more about projection predictive variable selection
- Regularized horseshoe talk at StanCon 2018 Asilomar
- Several case studies
- References with links to open access pdfs