

# Discrete Mathematics and Its Applications

## Lecture 7: Graphs: Euler, Hamilton and Coloring

MING GAO

DaSE@ECNU

(for course related communications)

mgao@dase.ecnu.edu.cn



Jan. 3, 2019

# Outline

- 1 Euler Paths and Circuits
- 2 Hamilton Paths and Circuits
- 3 Planar Graphs
  - Euler's Formula
  - Homeomorphic
- 4 Graph Coloring
- 5 Take-aways

# Euler circuit and euler path



The town of Königsberg, Prussia (now called Kaliningrad and part of the Russian republic), was divided into four sections by the branches of the Pregel River.



People wondered whether it was possible to start at some location in the town, travel across all the bridges once without crossing any bridge twice, and return to the starting point.

# Euler circuit and euler path

The town of Königsberg, Prussia (now called Kaliningrad and part of the Russian republic), was divided into four sections by the branches of the Pregel River.




People wondered whether it was possible to start at some location in the town, travel across all the bridges once without crossing any bridge twice, and return to the starting point.


## Definition

An **Euler circuit** in a graph  $G$  is a simple circuit containing every edge of  $G$ . An **Euler path** in  $G$  is a simple path containing every edge of  $G$ .

# Example




# Example



**Question:** Which of the directed graphs in figure have an Euler circuit? Of those that do not, which have an Euler path?

# Example




**Question:** Which of the directed graphs in figure have an Euler circuit? Of those that do not, which have an Euler path?

**Solution:**

The graph  $H_2$  has an Euler circuit, for example,  $a, g, c, b, g, e, d, f, a$ .

# Example



**Question:** Which of the directed graphs in figure have an Euler circuit? Of those that do not, which have an Euler path?


**Solution:**

The graph  $H_2$  has an Euler circuit, for example,  $a, g, c, b, g, e, d, f, a$ .

Neither  $H_1$  nor  $H_3$  has an Euler circuit.



# Example



**Question:** Which of the directed graphs in figure have an Euler circuit? Of those that do not, which have an Euler path?

**Solution:**

The graph  $H_2$  has an Euler circuit, for example,  $a, g, c, b, g, e, d, f, a$ .

Neither  $H_1$  nor  $H_3$  has an Euler circuit.

$H_3$  has an Euler path, namely,  $c, a, b, c, d, b$ , but  $H_1$  does not.

# Necessary and sufficient for Euler circuit

## Theorem

A connected multigraph with at least two vertices has an Euler circuit if and only if each of its vertices has even degree.

## Algorithm to find Euler circuit

### ALGORITHM 1 Constructing Euler Circuits.

```

procedure Euler(G: connected multigraph with all vertices of
    even degree)
  circuit := a circuit in G beginning at an arbitrarily chosen
    vertex with edges successively added to form a path that
    returns to this vertex
  H := G with the edges of this circuit removed
  while H has edges
    subcircuit := a circuit in H beginning at a vertex in H that
      also is an endpoint of an edge of circuit
    H := H with edges of subcircuit and all isolated vertices
      removed
    circuit := circuit with subcircuit inserted at the appropriate
      vertex
  return circuit {circuit is an Euler circuit}

```

# Necessary and sufficient for Euler circuit

## Theorem

A connected multigraph with at least two vertices has an Euler circuit if and only if each of its vertices has even degree.

## Algorithm to find Euler circuit

### ALGORITHM 1 Constructing Euler Circuits.

```

procedure Euler( $G$ : connected multigraph with all vertices of
    even degree)
  circuit := a circuit in  $G$  beginning at an arbitrarily chosen
    vertex with edges successively added to form a path that
    returns to this vertex
   $H := G$  with the edges of this circuit removed
  while  $H$  has edges
    subcircuit := a circuit in  $H$  beginning at a vertex in  $H$  that
      also is an endpoint of an edge of circuit
     $H := H$  with edges of subcircuit and all isolated vertices
      removed
    circuit := circuit with subcircuit inserted at the appropriate
      vertex
  return circuit {circuit is an Euler circuit}
  
```

Algorithm 1 provides an efficient algorithm for finding Euler circuits in a connected multigraph  $G$  with all vertices of even degree.

# Necessary and sufficient for Euler path

## Theorem




A connected multigraph has an Euler path but not an Euler circuit if and only if it has exactly two vertices of odd degree.

# Necessary and sufficient for Euler path

## Theorem

A connected multigraph has an Euler path but not an Euler circuit if and only if it has exactly two vertices of odd degree.

## Algorithm to find Euler path


 $G_1$ 

 $G_2$ 

 $G_3$ 

Which graphs shown in the figure have an Euler path?

# Hamilton paths and circuits


## Definition

A simple path in a graph  $G$  that passes through every vertex exactly once is called a **Hamilton path**, and a simple circuit in a graph  $G$  that passes through every vertex exactly once is called a **Hamilton circuit**.

# Hamilton paths and circuits

## Definition

A simple path in a graph  $G$  that passes through every vertex exactly once is called a **Hamilton path**, and a simple circuit in a graph  $G$  that passes through every vertex exactly once is called a **Hamilton circuit**.



Which graphs shown in the figure have an Hamilton path?

# Conditions for the existence of Hamilton circuit

## Dirac's theorem

If  $G$  is a simple graph with  $n$  vertices with  $n \geq 3$  such that the degree of every vertex in  $G$  is at least  $n/2$ , then  $G$  has a Hamilton circuit.



# Conditions for the existence of Hamilton circuit

## Dirac's theorem

If  $G$  is a simple graph with  $n$  vertices with  $n \geq 3$  such that the degree of every vertex in  $G$  is at least  $n/2$ , then  $G$  has a Hamilton circuit.

## Ore's theorem

If  $G$  is a simple graph with  $n$  vertices with  $n \geq 3$  such that  $\deg(u) + \deg(v) \geq n$  for every pair of nonadjacent vertices  $u$  and  $v$  in  $G$ , then  $G$  has a Hamilton circuit.

# Planar graph



## Definition

A graph is called **planar** if it can be drawn in the plane without any edges crossing (where a crossing of edges is the intersection of the lines or arcs representing them at a point other than their common endpoint).

# Planar graph

## Definition

A graph is called **planar** if it can be drawn in the plane without any edges crossing (where a crossing of edges is the intersection of the lines or arcs representing them at a point other than their common endpoint).





Are  $K_4$  and  $Q_3$  planar graphs?



# Planar graph

## Definition


A graph is called **planar** if it can be drawn in the plane without any edges crossing (where a crossing of edges is the intersection of the lines or arcs representing them at a point other than their common endpoint).



Are  $K_4$  and  $Q_3$  planar graphs?




# Example




Is  $K_{3,3}$  a planar graph?

# Example



Is  $K_{3,3}$  a planar graph?

**Solution:** In any planar representation of  $K_{3,3}$ ,  $v_1$  and  $v_2$  must be connected to both  $v_4$  and  $v_5$ . These four edges form a closed curve that splits the plane into two regions,  $R_1$  and  $R_2$ , as shown in Figure (a).  $v_3$  is in either  $R_1$  or  $R_2$ . When  $v_3$  is in  $R_2$ , the inside of the closed curve, the edges between  $v_3$  and  $v_4$  and between  $v_3$  and  $v_5$  separate  $R_2$  into two subregions,  $R_{21}$  and  $R_{22}$ , as shown in Figure (b). Note that there is no way to place  $v_6$ .



# Outline

- 1 Euler Paths and Circuits
- 2 Hamilton Paths and Circuits
- 3 Planar Graphs**
  - Euler's Formula
  - Homeomorphic
- 4 Graph Coloring
- 5 Take-aways

# Euler's formula

## Theorem

Let  $G$  be a connected planar simple graph with  $e$  edges and  $v$  vertices. Let  $r$  be the number of regions in a planar representation of  $G$ . Then  $r = e - v + 2$ .



# Euler's formula

## Theorem

Let  $G$  be a connected planar simple graph with  $e$  edges and  $v$  vertices. Let  $r$  be the number of regions in a planar representation of  $G$ . Then  $r = e - v + 2$ .

## Proof.

**Basic step:** The relationship  $r_1 = e_1 - v_1 + 2$  is true for  $G_1$ , because  $e_1 = 1$ ,  $v_1 = 2$ , and  $r_1 = 1$ .

# Euler's formula

## Theorem

Let  $G$  be a connected planar simple graph with  $e$  edges and  $v$  vertices. Let  $r$  be the number of regions in a planar representation of  $G$ . Then  $r = e - v + 2$ .

## Proof.

**Basic step:** The relationship  $r_1 = e_1 - v_1 + 2$  is true for  $G_1$ , because  $e_1 = 1$ ,  $v_1 = 2$ , and  $r_1 = 1$ .

**Induction step:** Now assume that  $r_k = e_k - v_k + 2$ . Let  $\{a_{k+1}, b_{k+1}\}$  be the edge that is added to  $G_k$  to obtain  $G_{k+1}$ .

# Euler's formula



## Theorem

Let  $G$  be a connected planar simple graph with  $e$  edges and  $v$  vertices. Let  $r$  be the number of regions in a planar representation of  $G$ . Then  $r = e - v + 2$ .

### Proof.

**Basic step:** The relationship  $r_1 = e_1 - v_1 + 2$  is true for  $G_1$ , because  $e_1 = 1$ ,  $v_1 = 2$ , and  $r_1 = 1$ .

**Induction step:** Now assume that  $r_k = e_k - v_k + 2$ . Let  $\{a_{k+1}, b_{k+1}\}$  be the edge that is added to  $G_k$  to obtain  $G_{k+1}$ .



# Proof of Euler's formula

Proof.

**Case I:** Both  $a_{k+1}$  and  $b_{k+1}$  are already in  $G_k$ .

# Proof of Euler's formula

Proof.

**Case I:** Both  $a_{k+1}$  and  $b_{k+1}$  are already in  $G_k$ . These two vertices must be on the boundary of a common region  $R$ , or else it would be impossible to add the edge  $\{a_{k+1}, b_{k+1}\}$  to  $G_k$  without two edges crossing (and  $G_{k+1}$  is planar).

# Proof of Euler's formula

Proof.

**Case I:** Both  $a_{k+1}$  and  $b_{k+1}$  are already in  $G_k$ . These two vertices must be on the boundary of a common region  $R$ , or else it would be impossible to add the edge  $\{a_{k+1}, b_{k+1}\}$  to  $G_k$  without two edges crossing (and  $G_{k+1}$  is planar). The addition of this new edge splits  $R$  into two regions. Consequently, in this case,  $r_{k+1} = r_k + 1$ ,  $e_{k+1} = e_k + 1$ , and  $v_{k+1} = v_k$ . Thus,  $r_{k+1} = e_{k+1} - v_{k+1} + 2$ .

# Proof of Euler's formula

## Proof.

**Case I:** Both  $a_{k+1}$  and  $b_{k+1}$  are already in  $G_k$ . These two vertices must be on the boundary of a common region  $R$ , or else it would be impossible to add the edge  $\{a_{k+1}, b_{k+1}\}$  to  $G_k$  without two edges crossing (and  $G_{k+1}$  is planar). The addition of this new edge splits  $R$  into two regions. Consequently, in this case,  $r_{k+1} = r_k + 1$ ,  $e_{k+1} = e_k + 1$ , and  $v_{k+1} = v_k$ . Thus,  $r_{k+1} = e_{k+1} - v_{k+1} + 2$ .

**Case II:** One of the two vertices of the new edge is not already in  $G_k$ .

# Proof of Euler's formula

## Proof.

**Case I:** Both  $a_{k+1}$  and  $b_{k+1}$  are already in  $G_k$ . These two vertices must be on the boundary of a common region  $R$ , or else it would be impossible to add the edge  $\{a_{k+1}, b_{k+1}\}$  to  $G_k$  without two edges crossing (and  $G_{k+1}$  is planar). The addition of this new edge splits  $R$  into two regions. Consequently, in this case,  $r_{k+1} = r_k + 1$ ,  $e_{k+1} = e_k + 1$ , and  $v_{k+1} = v_k$ . Thus,  $r_{k+1} = e_{k+1} - v_{k+1} + 2$ .

**Case II:** One of the two vertices of the new edge is not already in  $G_k$ . Suppose that  $a_{k+1}$  is in  $G_k$  but that  $b_{k+1}$  is not.



# Proof of Euler's formula

## Proof.

**Case I:** Both  $a_{k+1}$  and  $b_{k+1}$  are already in  $G_k$ . These two vertices must be on the boundary of a common region  $R$ , or else it would be impossible to add the edge  $\{a_{k+1}, b_{k+1}\}$  to  $G_k$  without two edges crossing (and  $G_{k+1}$  is planar). The addition of this new edge splits  $R$  into two regions. Consequently, in this case,  $r_{k+1} = r_k + 1$ ,  $e_{k+1} = e_k + 1$ , and  $v_{k+1} = v_k$ . Thus,  $r_{k+1} = e_{k+1} - v_{k+1} + 2$ .

**Case II:** One of the two vertices of the new edge is not already in  $G_k$ . Suppose that  $a_{k+1}$  is in  $G_k$  but that  $b_{k+1}$  is not. Adding this new edge does not produce any new regions, because  $b_{k+1}$  must be in a region that has  $a_{k+1}$  on its boundary. Consequently,  $r_{k+1} = r_k$ ,  $e_{k+1} = e_k + 1$ , and  $v_{k+1} = v_k + 1$ . Thus,  $r_{k+1} = e_{k+1} - v_{k+1} + 2$ .

# Proof of Euler's formula

## Proof.

**Case I:** Both  $a_{k+1}$  and  $b_{k+1}$  are already in  $G_k$ . These two vertices must be on the boundary of a common region  $R$ , or else it would be impossible to add the edge  $\{a_{k+1}, b_{k+1}\}$  to  $G_k$  without two edges crossing (and  $G_{k+1}$  is planar). The addition of this new edge splits  $R$  into two regions. Consequently, in this case,  $r_{k+1} = r_k + 1$ ,  $e_{k+1} = e_k + 1$ , and  $v_{k+1} = v_k$ . Thus,  $r_{k+1} = e_{k+1} - v_{k+1} + 2$ .

**Case II:** One of the two vertices of the new edge is not already in  $G_k$ . Suppose that  $a_{k+1}$  is in  $G_k$  but that  $b_{k+1}$  is not. Adding this new edge does not produce any new regions, because  $b_{k+1}$  must be in a region that has  $a_{k+1}$  on its boundary. Consequently,  $r_{k+1} = r_k$ ,  $e_{k+1} = e_k + 1$ , and  $v_{k+1} = v_k + 1$ . Thus,  $r_{k+1} = e_{k+1} - v_{k+1} + 2$ . We have completed the induction argument. Hence,  $r_n = e_n - v_n + 2$  for all  $n$ . □

## Corollary I

If  $G$  is a connected planar simple graph with  $e$  edges and  $v$  vertices, where  $v \geq 3$ , then  $e \leq 3v - 6$ .

## Corollary I

If  $G$  is a connected planar simple graph with  $e$  edges and  $v$  vertices, where  $v \geq 3$ , then  $e \leq 3v - 6$ .

**Proof.**

A connected planar simple graph drawn in the plane divides the plane into regions, say  $r$  of them. The degree of each region is at least three.

## Corollary I

If  $G$  is a connected planar simple graph with  $e$  edges and  $v$  vertices, where  $v \geq 3$ , then  $e \leq 3v - 6$ .

**Proof.**

A connected planar simple graph drawn in the plane divides the plane into regions, say  $r$  of them. The degree of each region is at least three. Note that the sum of the degrees of the regions is exactly twice the number of edges in the graph, because each edge occurs on the boundary of a region exactly twice. It follows that

$$2e = \sum_{\text{all regions } R} \deg(R) \geq 3r.$$

## Corollary 1

If  $G$  is a connected planar simple graph with  $e$  edges and  $v$  vertices, where  $v \geq 3$ , then  $e \leq 3v - 6$ .

**Proof.**

A connected planar simple graph drawn in the plane divides the plane into regions, say  $r$  of them. The degree of each region is at least three. Note that the sum of the degrees of the regions is exactly twice the number of edges in the graph, because each edge occurs on the boundary of a region exactly twice. It follows that

$$2e = \sum_{\text{all regions } R} \deg(R) \geq 3r.$$

Using Euler's formula, we obtain  $e - v + 2 = r \leq (2/3)e$ .

This shows that  $e \leq 3v - 6$ . □

## Corollary II

If  $G$  is a connected planar simple graph, then  $G$  has a vertex of degree not exceeding five.

## Corollary II

If  $G$  is a connected planar simple graph, then  $G$  has a vertex of degree not exceeding five.

**Proof.**

If  $G$  has one or two vertices, the result is true. If  $G$  has at least three vertices, by Corollary 1 we know that  $e \leq 3v - 6$ , so  $2e \leq 6v - 12$ .



## Corollary II

If  $G$  is a connected planar simple graph, then  $G$  has a vertex of degree not exceeding five.

**Proof.**

If  $G$  has one or two vertices, the result is true. If  $G$  has at least three vertices, by Corollary 1 we know that  $e \leq 3v - 6$ , so  $2e \leq 6v - 12$ .

If the degree of every vertex were at least six, then because  $2e = \sum_{v \in V} \deg(v)$  (by the handshaking theorem), we would have  $2e \geq 6v$ . But this contradicts the inequality  $2e \leq 6v - 12$ .

## Corollary II

If  $G$  is a connected planar simple graph, then  $G$  has a vertex of degree not exceeding five.

**Proof.**

If  $G$  has one or two vertices, the result is true. If  $G$  has at least three vertices, by Corollary 1 we know that  $e \leq 3v - 6$ , so  $2e \leq 6v - 12$ .

If the degree of every vertex were at least six, then because  $2e = \sum_{v \in V} \deg(v)$  (by the handshaking theorem), we would have  $2e \geq 6v$ . But this contradicts the inequality  $2e \leq 6v - 12$ .

It follows that there must be a vertex with degree no greater than five. □

## Corollary III

If a connected planar simple graph has  $e$  edges and  $v$  vertices with  $v \geq 3$  and no circuits of length three, then  $e \leq 2v - 4$ .

Proof.

The proof of this corollary is similar to that of Corollary 1, except that in this case the fact that there are no circuits of length three implies that the degree of a region must be at least four.  $\square$

## Corollary III

If a connected planar simple graph has  $e$  edges and  $v$  vertices with  $v \geq 3$  and no circuits of length three, then  $e \leq 2v - 4$ .

Proof.

The proof of this corollary is similar to that of Corollary 1, except that in this case the fact that there are no circuits of length three implies that the degree of a region must be at least four.  $\square$

### Example

**Question:** Determine whether  $K_5$  and  $K_{3,3}$  are planar graphs or not.

## Corollary III

If a connected planar simple graph has  $e$  edges and  $v$  vertices with  $v \geq 3$  and no circuits of length three, then  $e \leq 2v - 4$ .

Proof.

The proof of this corollary is similar to that of Corollary 1, except that in this case the fact that there are no circuits of length three implies that the degree of a region must be at least four.  $\square$

### Example

**Question:** Determine whether  $K_5$  and  $K_{3,3}$  are planar graphs or not.

**Solution:**

For  $K_5$ , we have  $e = 10$  and  $3v - 6 = 9$ . However, the inequality  $e \leq 3v - 6$  is not satisfied. Therefore,  $K_5$  is not planar.

## Corollary III

If a connected planar simple graph has  $e$  edges and  $v$  vertices with  $v \geq 3$  and no circuits of length three, then  $e \leq 2v - 4$ .

**Proof.**

The proof of this corollary is similar to that of Corollary 1, except that in this case the fact that there are no circuits of length three implies that the degree of a region must be at least four.  $\square$

### Example

**Question:** Determine whether  $K_5$  and  $K_{3,3}$  are planar graphs or not.

**Solution:**

For  $K_5$ , we have  $e = 10$  and  $3v - 6 = 9$ . However, the inequality  $e \leq 3v - 6$  is not satisfied. Therefore,  $K_5$  is not planar.

Because  $K_{3,3}$  has no circuits of length three, we have  $e = 9$  and  $2v - 4 = 8$ . Since Corollary 3 is not satisfied,  $K_{3,3}$  is nonplanar.


# Outline

- 1 Euler Paths and Circuits
- 2 Hamilton Paths and Circuits
- 3 Planar Graphs**
  - Euler's Formula
  - Homeomorphic**
- 4 Graph Coloring
- 5 Take-aways

# Homeomorphic

## Definition

If a graph is planar, so will be any graph obtained by removing an edge  $\{u, v\}$  and adding a new vertex  $w$  together with edges  $\{u, w\}$  and  $\{w, v\}$ . Such an operation is called an **elementary subdivision**. The graphs  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  are called **homeomorphic** if they can be obtained from the same graph by a sequence of elementary subdivisions.




**Question:** Determine whether the graphs  $G_1, G_2$ , and  $G_3$  are all homeomorphic or not.



# Homeomorphic application

## Kuratowski's theorem





A graph is nonplanar if and only if it contains a subgraph homeomorphic to  $K_{3,3}$  or  $K_5$ .



**Question:** Determine whether the graph  $G$  shown in the figure is planar.

# Problem formulation

## Motivation



Consider the problem of determining the least number of colors that can be used to color a map so that adjacent regions never have the same color.

# Definitions

## Coloring

A **coloring** of a simple graph is the assignment of a color to each vertex of the graph so that no two adjacent vertices are assigned the same color.

# Definitions

## Coloring

A **coloring** of a simple graph is the assignment of a color to each vertex of the graph so that no two adjacent vertices are assigned the same color.

## Chromatic number

The **chromatic number** of a graph is the least number of colors needed for a coloring of this graph. The chromatic number of a graph  $G$  is denoted by  $\chi(G)$ .


# Definitions

## Coloring

A **coloring** of a simple graph is the assignment of a color to each vertex of the graph so that no two adjacent vertices are assigned the same color.

## Chromatic number

The **chromatic number** of a graph is the least number of colors needed for a coloring of this graph. The chromatic number of a graph  $G$  is denoted by  $\chi(G)$ .



**Question:** What are the chromatic numbers of graphs  $G$  and  $H$ ?

# The four color theorem

## Theorem

The chromatic number of a planar graph is no greater than four.

# The four color theorem

## Theorem

The chromatic number of a planar graph is no greater than four.

### Remarks:

- Note that the four color theorem applies only to planar graphs.
- Nonplanar graphs can have arbitrarily large chromatic numbers.


# The four color theorem

## Theorem

The chromatic number of a planar graph is no greater than four.

### Remarks:

- Note that the four color theorem applies only to planar graphs.
- Nonplanar graphs can have arbitrarily large chromatic numbers.



**Question:** What are the chromatic numbers of graphs  $K_5$ ,  $K_{3,3}$ ,  $C_5$  and  $C_6$ ?



# Take-aways

## Conclusions

- Euler Paths and Circuits
- Hamilton Paths and Circuits
- Planar Graphs
  - Euler's Formula
  - Homeomorphic
- Graph Coloring