

Discrete Mathematics and Its Applications

Lecture 5: Discrete Probability: Probability Basics

MING GAO



DaSE@ ECNU
(for course related communications)
mgao@dase.ecnu.edu.cn

Nov. 22, 2018

Outline

- 1 Introduction
- 2 Sample Space and Events
- 3 Probability and Set Operations
 - Probability of Union
 - Probability of Complement
 - Independence
 - Conditional Probability
- 4 Take-aways

Introduction



Probability as a mathematical framework for:

- reasoning about uncertainty
- developing approaches to inference problems

Experiment and sample space

Experiment

An **experiment** is a procedure that yields one of a given set of possible outcomes.

Experiment and sample space

Experiment

An **experiment** is a procedure that yields one of a given set of possible outcomes.

Sample space

The **sample space**, denoted as Ω , of the experiment is the set of possible outcomes.

Experiment and sample space

Experiment

An **experiment** is a procedure that yields one of a given set of possible outcomes.

Sample space

The **sample space**, denoted as Ω , of the experiment is the set of possible outcomes.

Example

- Roll a die one time,
 $\Omega = \{1, 2, 3, 4, 5, 6\}$.
- We toss a coin twice (Head = H, Tail = T),
 $\Omega = \{HH, HT, TH, TT\}$.

Experiment and sample space

Experiment

An **experiment** is a procedure that yields one of a given set of possible outcomes.

Sample space


The **sample space**, denoted as Ω , of the experiment is the set of possible outcomes.

Example

- Roll a die one time,
 $\Omega = \{1, 2, 3, 4, 5, 6\}$.
- We toss a coin twice (Head = H, Tail = T),
 $\Omega = \{HH, HT, TH, TT\}$.

- “List” (set) of possible outcomes
- List must be:
 - ① Mutually exclusive
 - ② Collectively exhaustive
- Art: to be at the “right” granularity

Continuous sample space



For this case, sample space $\Omega = \{(x, y) | 0 \leq x, y \leq 1\}$. Note that the sample space is infinite and uncountable.

In this course, we only consider the **countable sample spaces**. Thus, we call the learning content to be the discrete probability.

Probability axioms

Event

An **event**, represented as a set, is a subset of the sample space.

Probability axioms

Event

An **event**, represented as a set, is a subset of the sample space.

Example

- Roll an even number,
 $A = \{2, 4, 6\} \subset \Omega$;
- Toss at least one head
 $B =$
 $\{HH, HT, TH\} \subset \Omega$;
- Toss at least three
head $C = \emptyset \subset \Omega$.
- There are $2^{|\Omega|}$ events
for an experiments;
- Events therefore have
all set operations.

Probability axioms

Event

An **event**, represented as a set, is a subset of the sample space.

Example

- Roll an even number,
 $A = \{2, 4, 6\} \subset \Omega$;
- Toss at least one head
 $B = \{HH, HT, TH\} \subset \Omega$;
- Toss at least three head $C = \emptyset \subset \Omega$.
- There are $2^{|\Omega|}$ events for an experiments;
- Events therefore have all set operations.

Axioms

- **Nonnegativity:** $P(A) \geq 0$;
- **Normalization:** $P(\Omega) = 1$ and $P(\emptyset) = 0$;
- **Additivity:** If $A \cap B = \emptyset$, then
 $P(A \cup B) = P(A) + P(B)$.

Furthermore, if $A_i \cap A_j = \emptyset$ for $\forall i \neq j$, then

$$P\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} P(A_i).$$

Finite probability

If S is a finite nonempty sample space of equally likely outcomes, and E is an event, that is, a subset of S , then the probability of E is

$$p(E) = \frac{|E|}{|S|}.$$

Finite probability

If S is a finite nonempty sample space of equally likely outcomes, and E is an event, that is, a subset of S , then the probability of E is

$$p(E) = \frac{|E|}{|S|}.$$

- Let all outcomes be equally likely;
- Computing probabilities \equiv two countings;
 - Counting the successful ways of the event;
 - Counting the size of the sample space;

Examples

Example I

Question: An urn contains four blue balls and five red balls. What is the probability that a ball chosen at random from the urn is blue?

Solution: Let S be the sample space, i.e.,

$$S = \{\textcircled{B}_1, \textcircled{B}_2, \textcircled{B}_3, \textcircled{B}_4, \textcircled{R}_1, \textcircled{R}_2, \textcircled{R}_3, \textcircled{R}_4, \textcircled{R}_5\}.$$

Let E be the event of choosing a blue ball, i.e.,

$$E = \{\textcircled{B}_1, \textcircled{B}_2, \textcircled{B}_3, \textcircled{B}_4\}.$$

In terms of the definition, we can compute the probability as


$$P(E) = \frac{|E|}{|S|} = \frac{4}{9}.$$

Examples Cont'd

Example II

Question: What is the probability that when two dice are rolled, the sum of the numbers on the two dice is 7?

Solution:



(1,1)	(1,2)	(1,3)	(1,4)	(1,5)	(1,6)
(2,1)	(2,2)	(2,3)	(2,4)	(2,5)	(2,6)
(3,1)	(3,2)	(3,3)	(3,4)	(3,5)	(3,6)
(4,1)	(4,2)	(4,3)	(4,4)	(4,5)	(4,6)
(5,1)	(5,2)	(5,3)	(5,4)	(5,5)	(5,6)
(6,1)	(6,2)	(6,3)	(6,4)	(6,5)	(6,6)


Examples Cont'd

Example II

Question: What is the probability that when two dice are rolled, the sum of the numbers on the two dice is 7?

Solution:

- There are a total of 36 possible outcomes when two dice are rolled.




(1,1)	(1,2)	(1,3)	(1,4)	(1,5)	(1,6)
(2,1)	(2,2)	(2,3)	(2,4)	(2,5)	(2,6)
(3,1)	(3,2)	(3,3)	(3,4)	(3,5)	(3,6)
(4,1)	(4,2)	(4,3)	(4,4)	(4,5)	(4,6)
(5,1)	(5,2)	(5,3)	(5,4)	(5,5)	(5,6)
(6,1)	(6,2)	(6,3)	(6,4)	(6,5)	(6,6)

Examples Cont'd

Example II

Question: What is the probability that when two dice are rolled, the sum of the numbers on the two dice is 7?

Solution:



(1,1)	(1,2)	(1,3)	(1,4)	(1,5)	(1,6)
(2,1)	(2,2)	(2,3)	(2,4)	(2,5)	(2,6)
(3,1)	(3,2)	(3,3)	(3,4)	(3,5)	(3,6)
(4,1)	(4,2)	(4,3)	(4,4)	(4,5)	(4,6)
(5,1)	(5,2)	(5,3)	(5,4)	(5,5)	(5,6)
(6,1)	(6,2)	(6,3)	(6,4)	(6,5)	(6,6)

- There are a total of 36 possible outcomes when two dice are rolled.
- There are six successful outcomes, namely, (1, 6), (2, 5), (3, 4), (4, 3), (5, 2), and (6, 1).
- Hence, the probability that a seven comes up when two fair dice are rolled is $6/36 = 1/6$.

Examples Cont'd

Example III

Question: In a lottery, players win a large prize when they pick four random digits that match, in the correct order. A smaller prize is won if only three digits are matched. What is the probability that a player wins the large prize? What is the probability that a player wins the small prize?

Examples Cont'd

Example III

Question: In a lottery, players win a large prize when they pick four random digits that match, in the correct order. A smaller prize is won if only three digits are matched. What is the probability that a player wins the large prize? What is the probability that a player wins the small prize?

Solution: By the product rule, there are $10^4 = 10,000$ ways to choose four digits.

Examples Cont'd

Example III

Question: In a lottery, players win a large prize when they pick four random digits that match, in the correct order. A smaller prize is won if only three digits are matched. What is the probability that a player wins the large prize? What is the probability that a player wins the small prize?

Solution: By the product rule, there are $10^4 = 10,000$ ways to choose four digits.

Large prize case: There is only one way to choose all four digits correctly. Thus, the probability is $1/10,000 = 0.0001$.

Examples Cont'd

Example III

Question: In a lottery, players win a large prize when they pick four random digits that match, in the correct order. A smaller prize is won if only three digits are matched. What is the probability that a player wins the large prize? What is the probability that a player wins the small prize?

Solution: By the product rule, there are $10^4 = 10,000$ ways to choose four digits.

Large prize case: There is only one way to choose all four digits correctly. Thus, the probability is $1/10,000 = 0.0001$.

Small prize case: Exactly one digit must be wrong to get three digits correct, but not all four correct.

Examples Cont'd

Example III

Question: In a lottery, players win a large prize when they pick four random digits that match, in the correct order. A smaller prize is won if only three digits are matched. What is the probability that a player wins the large prize? What is the probability that a player wins the small prize?

Solution: By the product rule, there are $10^4 = 10,000$ ways to choose four digits.

Large prize case: There is only one way to choose all four digits correctly. Thus, the probability is $1/10,000 = 0.0001$.

Small prize case: Exactly one digit must be wrong to get three digits correct, but not all four correct. Hence, there is a total of $\binom{4}{1} \times 9 = 36$ ways to choose four digits with exactly three of the four digits correct. Thus, the probability that a player wins the smaller prize is $36/10,000 = 9/2500 = 0.0036$.

Examples Cont'd

Example IV

Question: Find the probabilities that a poker hand contains four cards of one kind, or a full house (i.e., three of one kind and two of another kind).

Examples Cont'd

Example IV

Question: Find the probabilities that a poker hand contains four cards of one kind, or a full house (i.e., three of one kind and two of another kind).

Solution: There are $C(52, 5)$ different hands of five cards.

Examples Cont'd

Example IV

Question: Find the probabilities that a poker hand contains four cards of one kind, or a full house (i.e., three of one kind and two of another kind).

Solution: There are $C(52, 5)$ different hands of five cards.

Case I: # hands of five cards with four cards of one kind is

$$C(13, 1)C(4, 4)C(48, 1).$$

Examples Cont'd

Example IV

Question: Find the probabilities that a poker hand contains four cards of one kind, or a full house (i.e., three of one kind and two of another kind).

Solution: There are $C(52, 5)$ different hands of five cards.

Case I: # hands of five cards with four cards of one kind is

$$C(13, 1)C(4, 4)C(48, 1).$$

Case II: # hands of three of one kind and two of another kind is

$$P(13, 2)C(4, 3)C(4, 2).$$

Examples Cont'd

Example IV

Question: Find the probabilities that a poker hand contains four cards of one kind, or a full house (i.e., three of one kind and two of another kind).

Solution: There are $C(52, 5)$ different hands of five cards.

Case I: # hands of five cards with four cards of one kind is

$$C(13, 1)C(4, 4)C(48, 1).$$

Case II: # hands of three of one kind and two of another kind is


$$P(13, 2)C(4, 3)C(4, 2).$$

Hence, the probabilities are

$$\frac{C(13, 1)C(4, 4)C(48, 1)}{C(52, 5)} \approx 0.00024, \quad \frac{C(13, 2)C(4, 3)C(4, 2)}{C(52, 5)} \approx 0.0014.$$


Examples Cont'd

Example V



Examples Cont'd

Example V




For this case, sample space

$$\Omega = \{(x, y) | 0 \leq x, y \leq 1\}.$$

Question: How to compute the probability of a point inside the circle area?

Examples Cont'd

Example V



For this case, sample space

$$\Omega = \{(x, y) | 0 \leq x, y \leq 1\}.$$


Question: How to compute the probability of a point inside the circle area?

Let E be an event that the point locates in the circle area C , where

$$C = \{(x, y) | x^2 + y^2 \leq 1 \wedge x, y \geq 0\}.$$

Examples Cont'd

Example V



For this case, sample space

$$\Omega = \{(x, y) | 0 \leq x, y \leq 1\}.$$

Question: How to compute the probability of a point inside the circle area?

Let E be an event that the point locates in the circle area C , where

$$C = \{(x, y) | x^2 + y^2 \leq 1 \wedge x, y \geq 0\}.$$

Then we have

$$P(E) = \frac{S(C)}{S(\Omega)},$$

where $S(\cdot)$ is the area of a plane region.

Assigning probabilities

Let Ω be the sample space of an experiment with a finite or countable number of outcomes. We assign a probability $P(s)$ to each outcome $s \in \Omega$. We require that two conditions be met:

- 1 $0 \leq P(s) \leq 1$ for each $s \in \Omega$;

Assigning probabilities

Let Ω be the sample space of an experiment with a finite or countable number of outcomes. We assign a probability $P(s)$ to each outcome $s \in \Omega$. We require that two conditions be met:

- ① $0 \leq P(s) \leq 1$ for each $s \in \Omega$;
- ② $\sum_{s \in \Omega} P(s) = 1$.

Function P from the set of all outcomes of the sample space Ω to $[0, 1]$ is called a **probability distribution**.

We can model experiments in which outcomes are either equally likely or not equally likely by choosing the appropriate function $P(s)$.

Example I

Question: What probabilities should we assign to the outcomes H (heads) and T (tails) when a fair coin is flipped? What probabilities should be assigned to these outcomes when the coin is biased so that heads comes up twice as often as tails?

Solution:

Fair case: For a fair coin, the probability that heads comes up when the coin is flipped equals the probability that tails comes up, so the outcomes are equally likely, i.e., $P(H) = P(T) = \frac{1}{2}$.

Example I

Question: What probabilities should we assign to the outcomes H (heads) and T (tails) when a fair coin is flipped? What probabilities should be assigned to these outcomes when the coin is biased so that heads comes up twice as often as tails?

Solution:

Fair case: For a fair coin, the probability that heads comes up when the coin is flipped equals the probability that tails comes up, so the outcomes are equally likely, i.e., $P(H) = P(T) = \frac{1}{2}$.

Unfair case: For the biased coin we have $P(H) = 2P(T)$. Since $P(H) + P(T) = 1$, it follows that $P(T) = 1/3$ and $P(H) = 2/3$.

Uniform distribution

Definition of uniform distribution

Suppose that Ω is a set with n elements. The **uniform distribution** assigns the probability $1/n$ to each element of Ω .

Uniform distribution

Definition of uniform distribution

Suppose that Ω is a set with n elements. The **uniform distribution** assigns the probability $1/n$ to each element of Ω .

Definition of event probability

The probability of event E is the sum of the probabilities of the outcomes in E (E is a countable set). That is,

$$P(E) = \sum_{s \in E} P(s).$$

(Note that when E is an infinite set, $\sum_{s \in E} P(s)$ is a convergent infinite series.)

Probability operators

Operators

Let Ω be the sample space, A and B be two events:

- 1 If $A \cap B = \emptyset$, then

$$P(A \cup B) = P(A) + P(B);$$

Probability operators

Operators

Let Ω be the sample space, A and B be two events:

- ① If $A \cap B = \emptyset$, then

$$P(A \cup B) = P(A) + P(B);$$

②

$$P(\overline{A}) = P(\Omega) - P(A) = 1 - P(A);$$

Probability operators

Operators

Let Ω be the sample space, A and B be two events:

- ① If $A \cap B = \emptyset$, then

$$P(A \cup B) = P(A) + P(B);$$

②

$$P(\overline{A}) = P(\Omega) - P(A) = 1 - P(A);$$

- ③ If A and B are independent, then

$$P(A \cap B) = P(A) \cdot P(B);$$

Probability operators

Operators

Let Ω be the sample space, A and B be two events:

- ① If $A \cap B = \emptyset$, then

$$P(A \cup B) = P(A) + P(B);$$

②

$$P(\bar{A}) = P(\Omega) - P(A) = 1 - P(A);$$

- ③ If A and B are independent, then

$$P(A \cap B) = P(A) \cdot P(B);$$

- ④ The conditional probability of A given B , denoted by $P(A|B)$, is computed as

$$P(A|B) = \frac{P(A \cap B)}{P(B)}.$$

Probability of union of two events

Theorem

Let E_1 and E_2 be events in the sample space Ω . Then

$$P(E_1 \cup E_2) = P(E_1) + P(E_2) - P(E_1 \cap E_2).$$

Proof.

Since we have $|E_1 \cup E_2| = |E_1| + |E_2| - |E_1 \cap E_2|$,

Probability of union of two events

Theorem

Let E_1 and E_2 be events in the sample space Ω . Then

$$P(E_1 \cup E_2) = P(E_1) + P(E_2) - P(E_1 \cap E_2).$$

Proof.

Since we have $|E_1 \cup E_2| = |E_1| + |E_2| - |E_1 \cap E_2|$,

$$\begin{aligned} P(E_1 \cup E_2) &= \frac{|E_1 \cup E_2|}{|\Omega|} = \frac{|E_1| + |E_2| - |E_1 \cap E_2|}{|\Omega|} \\ &= \frac{|E_1|}{|\Omega|} + \frac{|E_2|}{|\Omega|} - \frac{|E_1 \cap E_2|}{|\Omega|} \\ &= P(E_1) + P(E_2) - P(E_1 \cap E_2). \end{aligned}$$



Probability of union Cont'd

- If $E_1 \cap E_2 = \emptyset$, then

$$P(E_1 \cup E_2) = P(E_1) + P(E_2);$$

- If $E_i \cap E_j = \emptyset$ for $\forall i, j$, then

$$P\left(\bigcup_{i=1}^{\infty} E_i\right) = \sum_{i=1}^{\infty} P(E_i);$$

-

$$\begin{aligned} P(E_1 \cup E_2 \cup E_3) &= P(E_1) + P(E_2) + P(E_3) \\ &\quad - P(E_1 \cap E_2) - P(E_1 \cap E_3) - P(E_2 \cap E_3) \\ &\quad + P(E_1 \cap E_2 \cap E_3) \end{aligned}$$

Example II

Question: What is the probability that a positive integer selected at random from the set of positive integers not exceeding 100 is divisible by either 2 or 5?

Solution: Let E_1 be the event that the integer selected at random is divisible by 2, and let E_2 be the event that it is divisible by 5.

Example II

Question: What is the probability that a positive integer selected at random from the set of positive integers not exceeding 100 is divisible by either 2 or 5?

Solution: Let E_1 be the event that the integer selected at random is divisible by 2, and let E_2 be the event that it is divisible by 5. Then $E_1 \cup E_2$ is the event that it is divisible by either 2 or 5.

Example II

Question: What is the probability that a positive integer selected at random from the set of positive integers not exceeding 100 is divisible by either 2 or 5?

Solution: Let E_1 be the event that the integer selected at random is divisible by 2, and let E_2 be the event that it is divisible by 5. Then $E_1 \cup E_2$ is the event that it is divisible by either 2 or 5.

$$\begin{aligned}P(E_1 \cup E_2) &= P(E_1) + P(E_2) - P(E_1 \cap E_2) \\&= \frac{50}{100} + \frac{20}{100} - \frac{10}{100} \\&= 0.6\end{aligned}$$

Probability of complement of an event

Theorem

Let E be an event in a sample space Ω . The probability of event $\bar{E} = \Omega - E$, the complementary event of E , is given by

$$P(\bar{E}) = 1 - P(E).$$

Proof.

Since we have $|\Omega| = |E| + |\bar{E}|$,

Probability of complement of an event

Theorem

Let E be an event in a sample space Ω . The probability of event $\bar{E} = \Omega - E$, the complementary event of E , is given by

$$P(\bar{E}) = 1 - P(E).$$

Proof.

Since we have $|\Omega| = |E| + |\bar{E}|$,

$$\begin{aligned} P(\bar{E}) &= \frac{|\bar{E}|}{|\Omega|} = \frac{|\Omega| - |E|}{|\Omega|} \\ &= \frac{|\Omega|}{|\Omega|} - \frac{|E|}{|\Omega|} = 1 - P(E). \end{aligned}$$



Example III

Question: A sequence of 10 bits is randomly generated. What is the probability that at least one of these bits is 0?

Solution: Let E be the event that at least one of the 10 bits is 0.

Example III

Question: A sequence of 10 bits is randomly generated. What is the probability that at least one of these bits is 0?

Solution: Let E be the event that at least one of the 10 bits is 0. Then \bar{E} is the event that all the bits are 1s. Because the sample space Ω is the set of all bit strings of length 10, it follows that

Example III

Question: A sequence of 10 bits is randomly generated. What is the probability that at least one of these bits is 0?

Solution: Let E be the event that at least one of the 10 bits is 0. Then \bar{E} is the event that all the bits are 1s. Because the sample space Ω is the set of all bit strings of length 10, it follows that

$$P(E) = 1 - P(\bar{E}) = 1 - \frac{1}{2^{10}} = \frac{1023}{1024}.$$

Running example

Tossing coins

We toss a coin twice (Head = H, Tail = T), then $\Omega = \{HH, HT, TH, TT\}$.

We define three events:

- ① A : the first toss is H ;
- ② B : the second toss is H ;
- ③ C : the first and second toss give the same results.

Hence, we have

- $P(A) = P(B) = P(C) = \frac{1}{2}$;
- $P(A \cap B) = P(A \cap C) = P(B \cap C) = \frac{|\{HH\}|}{|\Omega|} = \frac{1}{4}$;
- $P(A \cap B \cap C) = \frac{|\{HH\}|}{|\Omega|} = \frac{1}{4}$;

Independence

Definition

- Events E_1 and E_2 are **pair-wise independent** if and only if

$$P(E_1 \cap E_2) = P(E_1)P(E_2);$$

- Events E_1, E_2, \dots, E_n are **mutually independent** if

$$P(E_{i_1} \cap E_{i_2} \cap \dots \cap E_{i_m}) = P(E_{i_1})P(E_{i_2}) \dots P(E_{i_m}),$$

where $i_j, j = 1, 2, \dots, m$, are integers with

$1 \leq i_1 < i_2 < \dots < i_m \leq n$ and $m \geq 2$.

Note that mutually independent must be pair-wise independent, but pair-wise independent may not imply mutually independent (shown in previous example).

Independence

Theorem

Events E and F are pair-wise independent, then

- E and \bar{F} are pair-wise independent;
- \bar{E} and F are pair-wise independent;
- \bar{E} and \bar{F} are pair-wise independent;

Proof.

$$\begin{aligned}P(E) &= P(E \cap (F \cup \bar{F})) = P((E \cap F) \cup (E \cap \bar{F})) \\&= P(E \cap F) + P(E \cap \bar{F}) \\P(E \cap \bar{F}) &= P(E) - P(E \cap F) = P(E) - P(E)P(F) \\&= P(E)(1 - P(F)) = P(E)P(\bar{F})\end{aligned}$$

Hence, we have E and \bar{F} are independent. □

Example IV

Question: Suppose E is the event that a randomly generated bit string of length four begins with a 1 and F is the event that this bit string contains an even number of 1s. Are E and F independent, if the 16 bit strings of length four are equally likely?

Solution: Because there are 16 bit strings of length four, it follows that

$$P(E) = P(F) = 8/16 = 1/2.$$

Example IV

Question: Suppose E is the event that a randomly generated bit string of length four begins with a 1 and F is the event that this bit string contains an even number of 1s. Are E and F independent, if the 16 bit strings of length four are equally likely?

Solution: Because there are 16 bit strings of length four, it follows that

$$P(E) = P(F) = 8/16 = 1/2.$$

Because $E \cap F = \{1111, 1100, 1010, 1001\}$, we see that

$$P(E \cap F) = \frac{4}{16} = \frac{1}{4} = \left(\frac{1}{2}\right)\left(\frac{1}{2}\right) = P(E)P(F).$$

Example IV

Question: Suppose E is the event that a randomly generated bit string of length four begins with a 1 and F is the event that this bit string contains an even number of 1s. Are E and F independent, if the 16 bit strings of length four are equally likely?

Solution: Because there are 16 bit strings of length four, it follows that

$$P(E) = P(F) = 8/16 = 1/2.$$

Because $E \cap F = \{1111, 1100, 1010, 1001\}$, we see that

$$P(E \cap F) = \frac{4}{16} = \frac{1}{4} = \left(\frac{1}{2}\right)\left(\frac{1}{2}\right) = P(E)P(F).$$

Hence, events E and F are independent.

Example V

Question: Assume that each of the four ways a family can have two children is equally likely. Are events E , that a family with two children has two boys, and F , that a family with two children has at least one boy, independent?

Solution: Because $E = \{BB\}$ and $F = \{BB, BG, GB\}$, we have $P(E) = 1/4$ and $P(F) = 3/4$.

Example V

Question: Assume that each of the four ways a family can have two children is equally likely. Are events E , that a family with two children has two boys, and F , that a family with two children has at least one boy, independent?

Solution: Because $E = \{BB\}$ and $F = \{BB, BG, GB\}$, we have $P(E) = 1/4$ and $P(F) = 3/4$.

Obviously, $E \cap F = \{BB\}$, we see that $P(E \cap F) = 1/4$.

Example V

Question: Assume that each of the four ways a family can have two children is equally likely. Are events E , that a family with two children has two boys, and F , that a family with two children has at least one boy, independent?

Solution: Because $E = \{BB\}$ and $F = \{BB, BG, GB\}$, we have $P(E) = 1/4$ and $P(F) = 3/4$.

Obviously, $E \cap F = \{BB\}$, we see that $P(E \cap F) = 1/4$. However, $P(E)P(F) = 3/16$.

That is

$$P(E \cap F) \neq P(E)P(F).$$

Example V

Question: Assume that each of the four ways a family can have two children is equally likely. Are events E , that a family with two children has two boys, and F , that a family with two children has at least one boy, independent?

Solution: Because $E = \{BB\}$ and $F = \{BB, BG, GB\}$, we have $P(E) = 1/4$ and $P(F) = 3/4$.

Obviously, $E \cap F = \{BB\}$, we see that $P(E \cap F) = 1/4$. However, $P(E)P(F) = 3/16$.

That is

$$P(E \cap F) \neq P(E)P(F).$$

Hence, events E and F are not independent.

Example VI

Question: Are the events E , that a family with three children has children of both sexes, and F , that this family has at most one boy, independent? Assume that the eight ways a family can have three children are equally likely.

Example VI

Question: Are the events E , that a family with three children has children of both sexes, and F , that this family has at most one boy, independent? Assume that the eight ways a family can have three children are equally likely.

Solution: Because $E = \{BBG, BGB, BGG, GBB, GBG, GGB\}$, $F = \{BGG, GBG, GGB, GGG\}$, and $E \cap F = \{BGG, GBG, GGB\}$, it follows that $P(E) = 6/8 = 3/4$, $P(F) = 4/8 = 1/2$, and $P(E \cap F) = 3/8$.

Example VI

Question: Are the events E , that a family with three children has children of both sexes, and F , that this family has at most one boy, independent? Assume that the eight ways a family can have three children are equally likely.

Solution: Because $E = \{BBG, BGB, BGG, GBB, GBG, GGB\}$, $F = \{BGG, GBG, GGB, GGG\}$, and $E \cap F = \{BGG, GBG, GGB\}$, it follows that $P(E) = 6/8 = 3/4$, $P(F) = 4/8 = 1/2$, and $P(E \cap F) = 3/8$.

Because $P(E)P(F) = 3/8$, it follows that $P(E \cap F) = P(E)P(F)$, so E and F are independent.

Conditional probability

Definition

Let E and F be events with $P(F) > 0$. The **conditional probability** of E given F , denoted by $P(E|F)$, is defined as


$$P(E|F) = \frac{P(E \cap F)}{P(F)}.$$

Conditional probability

Definition

Let E and F be events with $P(F) > 0$. The **conditional probability** of E given F , denoted by $P(E|F)$, is defined as

$$P(E|F) = \frac{P(E \cap F)}{P(F)}.$$




Conditional probability

Definition

Let E and F be events with $P(F) > 0$. The **conditional probability** of E given F , denoted by $P(E|F)$, is defined as

$$P(E|F) = \frac{P(E \cap F)}{P(F)}.$$



- $P(E|F)$ is the probability of E , given that F occurred
- F is our new sample space;
- $P(E|F)$ is undefined if $P(F) = 0$.

Example VII

Question: What is the conditional probability that a family with two children has two boys, given they have at least one boy? Assume that each of the possibilities BB, BG, GB, and GG is equally likely.

Example VII

Question: What is the conditional probability that a family with two children has two boys, given they have at least one boy? Assume that each of the possibilities BB, BG, GB, and GG is equally likely.

Solution: Let E be the event that a family with two children has two boys, and let F be the event that a family with two children has at least one boy.

Example VII

Question: What is the conditional probability that a family with two children has two boys, given they have at least one boy? Assume that each of the possibilities BB, BG, GB, and GG is equally likely.

Solution: Let E be the event that a family with two children has two boys, and let F be the event that a family with two children has at least one boy.

It follows that $E = \{BB\}$, $F = \{BB, BG, GB\}$, and $E \cap F = \{BB\}$.

Example VII

Question: What is the conditional probability that a family with two children has two boys, given they have at least one boy? Assume that each of the possibilities BB, BG, GB, and GG is equally likely.

Solution: Let E be the event that a family with two children has two boys, and let F be the event that a family with two children has at least one boy.

It follows that $E = \{BB\}$, $F = \{BB, BG, GB\}$, and $E \cap F = \{BB\}$. Because the four possibilities are equally likely, it follows that $P(F) = 3/4$ and $P(E \cap F) = 1/4$.

Example VII

Question: What is the conditional probability that a family with two children has two boys, given they have at least one boy? Assume that each of the possibilities BB, BG, GB, and GG is equally likely.

Solution: Let E be the event that a family with two children has two boys, and let F be the event that a family with two children has at least one boy.

It follows that $E = \{BB\}$, $F = \{BB, BG, GB\}$, and $E \cap F = \{BB\}$. Because the four possibilities are equally likely, it follows that $P(F) = 3/4$ and $P(E \cap F) = 1/4$.

We conclude that


$$P(E|F) = \frac{P(E \cap F)}{P(F)} = \frac{1/4}{3/4} = \frac{1}{3}.$$

Remarks for conditional probability

- $P(E|F) = P(E)$ if events E and F are independent;


Remarks for conditional probability

- $P(E|F) = P(E)$ if events E and F are independent;
- $P(E \cap F) = P(E) \cdot P(F|E) = P(F) \cdot P(E|F)$;



Remarks for conditional probability


- $P(E|F) = P(E)$ if events E and F are independent;
- $P(E \cap F) = P(E) \cdot P(F|E) = P(F) \cdot P(E|F)$;



- $P(E) = P(F_1) \cdot P(E|F_1) + P(F_2) \cdot P(E|F_2) + P(F_3) \cdot P(E|F_3)$ if $F_1 \cup F_2 \cup F_3 = \Omega$ and $F_i \cap F_j = \emptyset$ for $i \neq j$ (**Total probability theorem**);

Remarks for conditional probability

- $P(E|F) = P(E)$ if events E and F are independent;
- $P(E \cap F) = P(E) \cdot P(F|E) = P(F) \cdot P(E|F)$;



- $P(E) = P(F_1) \cdot P(E|F_1) + P(F_2) \cdot P(E|F_2) + P(F_3) \cdot P(E|F_3)$ if $F_1 \cup F_2 \cup F_3 = \Omega$ and $F_i \cap F_j = \emptyset$ for $i \neq j$ (**Total probability theorem**);
- $P(F_i|E) = \frac{P(F_i) \cdot P(E|F_i)}{P(E)} = \frac{P(F_i) \cdot P(E|F_i)}{\sum_j P(F_j) \cdot P(E|F_j)}$ (**Bayes rule**).

Proof of the total probability theorem

Theorem

Let F_i (for $i = 1, 2, \dots, n$) be a partition of sample space Ω , for any event E , then

$$P(E) = \sum_{i=1}^n P(F_i) \cdot P(E|F_i).$$

Proof of the total probability theorem

Theorem

Let F_i (for $i = 1, 2, \dots, n$) be a partition of sample space Ω , for any event E , then

$$P(E) = \sum_{i=1}^n P(F_i) \cdot P(E|F_i).$$

Proof:

$$\begin{aligned} P(E) &= P(E \cap \Omega) = P(E \cap (F_1 \cup F_2 \cup \dots \cup F_n)) \\ &= P((E \cap F_1) \cup (E \cap F_2) \cup \dots \cup (E \cap F_n)) \\ &= P(E \cap F_1) + P(E \cap F_2) + \dots + P(E \cap F_n) \\ &= P(F_1) \cdot P(E|F_1) + P(F_2) \cdot P(E|F_2) + \dots + P(F_n) \cdot P(E|F_n) \\ &= \sum_{i=1}^n P(F_i) \cdot P(E|F_i). \end{aligned}$$

Example VIII

Question: Suppose we draw two cards from a well shuffled deck. What is the probability the second card in the deck is an ace?

Example VIII

Question: Suppose we draw two cards from a well shuffled deck. What is the probability the second card in the deck is an ace?

Solution: Let E and F be events that the first and the second cards are Aces, respectively. Hence E and \overline{E} is a partition of the sample space.

Example VIII

Question: Suppose we draw two cards from a well shuffled deck. What is the probability the second card in the deck is an ace?

Solution: Let E and F be events that the first and the second cards are Aces, respectively. Hence E and \bar{E} is a partition of the sample space.

In terms of the total probability theorem, condition on whether the first card is an ace or not:

$$\begin{aligned} P(F) &= P(E) \cdot P(F|E) + P(\bar{E}) \cdot P(F|\bar{E}) \\ &= \left(\frac{4}{52}\right)\left(\frac{3}{51}\right) + \left(\frac{48}{52}\right)\left(\frac{4}{51}\right) \\ &= \frac{3 \cdot 4 + 48 \cdot 4}{51 \cdot 52} = \frac{4}{52} \end{aligned}$$

Take-aways

Conclusions

- Introduction
- Sample Space and Events
- Probability and Set Operations
 - Probability of Union
 - Probability of Complement
 - Independence
 - Conditional Probability