


# PATTERNS IN RECTANGULATIONS


Michaela A. Polley  
Dartmouth College

Women in Mathematics in New England 2025  
Northampton, MA, USA  
September 20, 2025




What is a rectangulation?




What is a rectangulation?




What is a rectangulation?




What is a rectangulation?




What is a rectangulation?



What is a rectangulation?




What is a rectangulation?







What is a rectangulation?




What is a rectangulation?




What is a rectangulation?




What is a rectangulation?




What is a rectangulation?




What is a rectangulation?



What is a rectangulation?




What is a rectangulation?






Weak and Strong Equivalence




**STRONGLY**  
equivalent

Weak and Strong Equivalence




**WEAKLY**  
equivalent


Weak and Strong Equivalence



$\mathcal{A}$



$\mathcal{B}$



$\mathcal{C}$

# What is combinatorics?

*Combinatorics* is the area of math that studies the properties of discrete structures (such as rectangulations) and tries to answer questions such as:

# What is combinatorics?

*Combinatorics* is the area of math that studies the properties of discrete structures (such as rectangulations) and tries to answer questions such as:

- How many are there? (Enumeration)

# What is combinatorics?

*Combinatorics* is the area of math that studies the properties of discrete structures (such as rectangulations) and tries to answer questions such as:

- How many are there? (Enumeration)
- How can they be constructed? (Algorithms)

# What is combinatorics?

*Combinatorics* is the area of math that studies the properties of discrete structures (such as rectangulations) and tries to answer questions such as:

- How many are there? (Enumeration)
- How can they be constructed? (Algorithms)
- Are there any structures satisfying a given condition? (Existence)

# What is combinatorics?

*Combinatorics* is the area of math that studies the properties of discrete structures (such as rectangulations) and tries to answer questions such as:

- How many are there? (Enumeration)
- How can they be constructed? (Algorithms)
- Are there any structures satisfying a given condition? (Existence)
- What is the best structure under some criteria? (Optimization)




# What is combinatorics?

*Combinatorics* is the area of math that studies the properties of discrete structures (such as rectangulations) and tries to answer questions such as:

- How many are there? (Enumeration)
- How can they be constructed? (Algorithms)
- Are there any structures satisfying a given condition? (Existence)
- What is the best structure under some criteria? (Optimization)

Pattern Avoidance:  $R(\top, \neg, \vdash)$



Pattern Avoidance:  $R(\top, \neg, \vdash)$


## Pattern Avoidance: $R(\top, \dashv, \vdash)$


Pattern Avoidance:  $R(\top, \dashv, \vdash)$


Proposition

The number of rectangulations of size  $n$  that avoid  $\top$ ,  $\dashv$ , and  $\vdash$ , denoted  $R_n(\top, \dashv, \vdash)$  is  $n$ .


# Pattern Avoidance: $R(\top, \dashv, \vdash)$

## Proposition

The number of rectangulations of size  $n$  that avoid  $\top$ ,  $\dashv$ , and  $\vdash$ , denoted  $R_n(\top, \dashv, \vdash)$  is  $n$ .

## Example

For  $n = 5$ , there are five rectangulations that avoid  $\top$ ,  $\dashv$ , and  $\vdash$ :




Pattern Avoidance:  $R(\neg, \vdash)$


Pattern Avoidance:  $R(-\vdash, \vdash)$




Pattern Avoidance:  $R(\dashv, \vdash)$



Proposition

The number of weak rectangulations of size  $n$  that avoid  $\dashv$  and  $\vdash$ , denoted  $R_n^w(\dashv, \vdash)$  is  $2^{n-1}$ .

# Pattern Avoidance: $R(\dashv, \vdash)$

## Proposition

The number of weak rectangulations of size  $n$  that avoid  $\dashv$  and  $\vdash$ , denoted  $R_n^w(\dashv, \vdash)$  is  $2^{n-1}$ .

## Definition

A *composition* of  $n$  is an ordered list of positive integers  $(a_1, a_2, \dots, a_k)$  such that  $a_1 + a_2 + \dots + a_k = n$ .

# Pattern Avoidance: $R(\dashv, \vdash)$

## Proposition

The number of weak rectangulations of size  $n$  that avoid  $\dashv$  and  $\vdash$ , denoted  $R_n^w(\dashv, \vdash)$  is  $2^{n-1}$ .

## Definition

A *composition* of  $n$  is an ordered list of positive integers  $(a_1, a_2, \dots, a_k)$  such that  $a_1 + a_2 + \dots + a_k = n$ .

$$3$$

$$2 + 1$$

$$1 + 2$$

$$1 + 1 + 1$$

# Pattern Avoidance: $R(\dashv, \vdash)$

## Proposition

The number of weak rectangulations of size  $n$  that avoid  $\dashv$  and  $\vdash$ , denoted  $R_n^w(\dashv, \vdash)$  is  $2^{n-1}$ .

## Definition

A *composition* of  $n$  is an ordered list of positive integers  $(a_1, a_2, \dots, a_k)$  such that  $a_1 + a_2 + \dots + a_k = n$ .

$$3$$

$$2 + 1$$

$$1 + 2$$

$$1 + 1 + 1$$

There are  $2^{n-1}$  compositions of  $n$ :

$$\bullet \bullet \bullet$$

$$\bullet \bullet \mid \bullet$$

$$\bullet \mid \bullet \bullet$$

$$\bullet \mid \bullet \mid \bullet$$


Pattern Avoidance:  $R(\dashv, \vdash)$

Proposition

The number of weak rectangulations of size  $n$  that avoid  $\dashv$  and  $\vdash$ , denoted  $R_n^w(\dashv, \vdash)$  is  $2^{n-1}$ .

Proof

Establish a bijection between  $R_n^w(\dashv, \vdash)$  and compositions of  $n$ .




Pattern Avoidance:  $R(\dashv, \vdash)$

Proposition

The number of weak rectangulations of size  $n$  that avoid  $\dashv$  and  $\vdash$ , denoted  $R_n^w(\dashv, \vdash)$  is  $2^{n-1}$ .

Proof

Establish a bijection between  $R_n^w(\dashv, \vdash)$  and compositions of  $n$ .




**Question:** What about strong rectangulations?

Pattern Avoidance:  $R(\dashv, \vdash)$

By complete enumeration (using a computer), we find that the number of *strong* rectangulations of size  $n$  which avoid  $\vdash$  and  $\dashv$  for  $n = 1, \dots, 8$  are 1, 2, 4, 9, 22, 57, 154, 430.

# Pattern Avoidance: $R(\dashv, \vdash)$

By complete enumeration (using a computer), we find that the number of *strong* rectangulations of size  $n$  which avoid  $\vdash$  and  $\dashv$  for  $n = 1, \dots, 8$  are 1, 2, 4, 9, 22, 57, 154, 430.



## The On-Line Encyclopedia of Integer Sequences (OEIS)

Enter a sequence, word, or sequence number:

oeis.org



Pattern Avoidance:  $R(\neg, \vdash)$

The OEIS is supported by [the many generous donors to the OEIS Foundation](#).

0 1 3 6 2 7  
: 13  
: 20  
23 12  
10 22 11 21

THE ON-LINE ENCYCLOPEDIA  
OF INTEGER SEQUENCES<sup>®</sup>

founded in 1964 by N. J. A. Sloane

1, 2, 4, 9, 22, 57, 154, 430

Search

Hints

(Greetings from [The On-Line Encyclopedia of Integer Sequences!](#))

Search: **seq:1,2,4,9,22,57,154,430**

Displaying 1-1 of 1 result found. page 1

Sort: relevance | [references](#) | [number](#) | [modified](#) | [created](#)    Format: long | [short](#) | [data](#)

A287709

Number of Dyck paths of semilength n such that every peak at level y > 1 is preceded by (at least) one peak at level y-1.

+30  
2

1, 1, **1, 2, 4, 9, 22, 57, 154, 430**, 1234, 3625, 10865, 33136, 102598, 321913, 1021963, 3278543, 10617413, 34678693, 114151769, 378436049, 1262822229, 4239469076, 14312153289, 48567846377, 165610404277, 567259571451, 1951218773118, 6738242931451, 23356148951482

([list](#); [graph](#); [refs](#); [listen](#); [history](#); [text](#); [internal format](#))

OFFSET  
0,4

COMMENTS  
Also the number of Dyck paths of semilength (n-1) whose maximum height is attained by the initial ascent. (That is, Dyck paths with prefix U<sup>k</sup>D, k>=1, and maximum height k.) For a(3)=2: UDUD, UUD. For a(4)=3: UDUDUD, UUDUD, UUDUD, UUDUD. (Andrei Asinowski and Vít Jelínek) - [Andrei Asinowski](#), Jun 21 2021  
From [Andrei Asinowski](#), Sep 01 2025: (Start)  
Also the number of strong rectangulations of size (n-1) that avoid the patterns "top" and "bottom" (that is, the T-shape and the upside-down T-shape). (Andrei Asinowski and Michaela Polley, Thm. 13).  
Also the number of (010,101,120,201)-avoiding inversion sequences e of length (n-1) in which all left-to-right maxima e<sub>j</sub> satisfy e<sub>j</sub>=j-1. Also the number of (010,110,120,210)-avoiding inversion sequences of length (n-1) that satisfy this condition. Also the number of (010,100,120,210)-avoiding inversion sequences of length (n-1) that satisfy this condition. (Andrei Asinowski and Michaela Polley, Prop. 14).  
Also the number of (011,201)-avoiding inversion sequences e of length (n-1) in which the set of values is precisely {0,1,2,...,M}, where M is the maximum value of e. (Andrei Asinowski and Michaela Polley, Prop. 15). (End)

REFERENCES  
Andrei Asinowski and Vít Jelínek. Two types of Dyck paths (unpublished manuscript).

LINKS  
Alois P. Heinz, [Table of n, a\(n\) for n = 0..1000](#)  
Andrei Asinowski and Michaela A. Polley, [Patterns in rectangulations. Part I: T-like patterns, inversion sequence classes I\(010, 101, 120, 201\) and I\(011, 201\), and rushed Dyck paths](#), arXiv:2501.11781 [math.CO], 2025. See pp. 1, 7, 25, 27.  
Axel Racher, [Progressive and rushed Dyck paths](#), arXiv:2403.08120 [math.CO], 2024

https://oeis.org/A287709