

# PCA

ML Instruction Team, Fall 2022

CE Department  
Sharif University of Technology

# Motivation

- PCA identifies one or more orthogonal directions that capture the greatest amount of variance in a feature matrix  $X \in \mathbb{R}^{m \times n}$ .
- Assuming zero-mean feature matrix  $X \in \mathbb{R}^{m \times n}$ , the variance of the samples' projections onto a **unit vector**  $v$  is given by:

$$\frac{1}{n} \sum_{i=1}^n (x_i^t v)^2 = \frac{1}{n} \|Xv\|^2 = \frac{1}{n} v^t X^t X v$$

- In light of this consideration, we define the first desired vector  $v_1$  as the solution to the constrained optimization problem:

$$\max_{\|v\|_2=1} v^t X^t X v$$

- We convert this constrained optimization problem into an unconstrained one by writing down its Lagrangian:

$$\mathcal{L}(v) := v^t X^t X v - \lambda(v^t v - 1)$$

# First PC

- First-order necessary conditions for optimal value imply that:

$$0 = \nabla \mathcal{L}(v_1) = 2X^t X v_1 - 2\lambda v_1$$

- Since  $X^t X v_1 = \lambda v_1$ ,  $v_1$  is an **eigenvector** of  $X^t X$  with eigenvalue  $\lambda$ .
- Since we constrain  $\|v_1\|_2^2 = v_1^t v_1 = 1$ , the value of the objective is precisely:

$$v_1^t X^t X v_1 = v_1^t (\lambda v_1) = \lambda v_1^t v_1 = \lambda$$

- The **optimal value** is  $\lambda = \lambda_{\max}(X^t X)$ , which is achieved when  $v_1$  is a **unit eigenvector** of  $X^t X$  corresponding to its **largest** eigenvalue.

# More PCs?

- How to find more direction with the desired property?
  - ▷ Ideally, the subsequent directions found should also be directions of high variance.
  - ▷ They should be orthogonal to the existing ones in order to minimize redundancy.

- We define the  $k$ -th loading vector  $v_k$  as the solution to the constrained optimization problem:

$$\max_v v^t X^t X v \quad \text{subject to} \quad v^t v = 1, v^t v_i = 0, \quad i = 1, \dots, k-1$$

- **Claim:**  $v_k$  is a **unit eigenvector** of  $X^t X$  corresponding to its  $k$ -th **largest** eigenvalue.
- The unit vector that defines the  $k$ -th axis is called the  $k$ -th principal component (PC).

# Evaluation of PCs

- Assuming the singular value decomposition of feature matrix  $X$  as follows:

$$X = U\Sigma V^T = [u_1, u_2, \dots, u_r] \begin{bmatrix} \sigma_1 & & & 0 \\ & \sigma_2 & & \\ & & \ddots & \\ 0 & & & \sigma_r \end{bmatrix} \begin{bmatrix} v_1^T \\ v_2^T \\ \vdots \\ v_r^T \end{bmatrix}$$

The first  $k$  PCs are  $W_k = [v_1, v_2, \dots, v_k]$ .

- Explained Variance Ratio** explains the proportion of the dataset's variance that lies along the axis of each PC.
- PCA can also be viewed as the projection of the sample points to the subspace with the minimum perpendicular distance.

# Other Derivation?

**Thank You!**

**Any Question?**