



Genomes per phylum: 23,681 genomes, 6,163 species names




Source: samples.csv. Species = SPECIES column (exact name match); 'Xxx sp. STRAIN' names collapsed to one 'Xxx sp.'. Missing family: 501 genomes (excluded from family/class charts).

Top 30 classes




■ All genomes (n=23,681)
■ One per species (n=6,163)

Top 30 orders




Source: samples.csv. Species = SPECIES column (exact name match); 'Xxx sp. STRAIN' names collapsed to one 'Xxx sp.'. Missing family: 501 genomes (excluded from family/class charts).

Top 30 families by genome count




■ All genomes (n=23,681)
■ One per species (n=6,163)

Top 30 families by number of species (one genome per species)




Ascomycota: top 30 families by genome count




Source: samples.csv. Species = SPECIES column (exact name match); 'Xxx sp. STRAIN' names collapsed to one 'Xxx sp.'. Missing family: 501 genomes (excluded from family/class charts).

Basidiomycota: top 30 families by genome count



■ All genomes (n=3,275)
■ One per species (n=1,437)

Genomes per species (redundancy), top 30 families



Source: samples.csv. Species = SPECIES column (exact name match); 'Xxx sp. STRAIN' names collapsed to one 'Xxx sp.'. Missing family: 501 genomes (excluded from family/class charts).