

Discrete Curvature Computation

Work Report

MinliangLIN

November 9, 2018


Outline

1 fxzlib of curvature

2 problems


What is curvature?

- A number indicating how bend a curve is, which lays on surface
- How fast does a curve leave the tangent plane?




Motivation

controlling the direction and size of quadrangulation




What is curvature

- A quadratic form of tangent vector $X^T B X$




What is curvature

- A quadratic form of tangent vector $X^T B X$
- B is a **linear map / tensor** from dX to dN



What is curvature

- A quadratic form of tangent vector $X^T B X$
- B is a **linear map / tensor** from dX to dN
- mean curvature $\kappa_H = \text{tr}(B)/2$




How to compute **mean curvature** on vertex?

- multiply by normal vector $\mathbf{K}(\mathbf{x}) = \kappa_H \mathbf{n}$
- $$\int \int_{\mathcal{A}_M} \mathbf{K}(\mathbf{x}) dA = \frac{1}{2} \sum_{j \in N_1(i)} (\cot \alpha_{ij} + \cot \beta_{ij}) (\mathbf{x}_i - \mathbf{x}_j)$$


How to compute **mean curvature** on vertex?

- multiply by normal vector $\mathbf{K}(\mathbf{x}) = \kappa_H \mathbf{n}$
- $\int \int_{\mathcal{A}_M} \mathbf{K}(\mathbf{x}) dA = \frac{1}{2} \sum_{j \in N_1(i)} (\cot \alpha_{ij} + \cot \beta_{ij}) (\mathbf{x}_i - \mathbf{x}_j)$





How to compute **mean curvature** on vertex?

- multiply by normal vector $\mathbf{K}(\mathbf{x}) = \kappa_H \mathbf{n}$
- $\int \int_{\mathcal{A}_M} \mathbf{K}(\mathbf{x}) dA = \frac{1}{2} \sum_{j \in N_1(i)} (\cot \alpha_{ij} + \cot \beta_{ij}) (\mathbf{x}_i - \mathbf{x}_j)$



How to compute **mean curvature** on vertex?

- multiply by normal vector $\mathbf{K}(\mathbf{x}) = \kappa_H \mathbf{n}$
- $\int \int_{\mathcal{A}_M} \mathbf{K}(\mathbf{x}) dA = \frac{1}{2} \sum_{j \in N_1(i)} (\cot \alpha_{ij} + \cot \beta_{ij})(\mathbf{x}_i - \mathbf{x}_j)$



How to get the tensor/matrix of vertex?

- $B = \begin{pmatrix} a & b \\ b & c \end{pmatrix}$
- sampling along different direction
- least square fitting the data constrain to mean curvature

How to get the tensor/matrix of vertex?

- $B = \begin{pmatrix} a & b \\ b & c \end{pmatrix}$
- sampling along different direction
- least square fitting the data constrain to mean curvature
- $E(a, b, c) = \sum_j w_j (\mathbf{d}_{i,j}^T B \mathbf{d}_{i,j} - \kappa_{i,j}^N)^2$ s.t. $a + c = 2\kappa_H, \overbrace{ac - b^2}^{\text{Hessian}} = \kappa_G$

Problems

■ Some potential trouble makers


- 1 Hard constrain of κ_H is replaced by soft constrain.
- 2 w_j s are not used
- 3 Area of computing κ_H is just barycenter area, and cotangents of obtuse angles are not handled, which may cause some negative weight of summation of κ_H .
- 4 [MDSB03] states that “the curvature values computed from the least squares are often less accurate in practice”.
- 5 We compute curvature of facet as the average of vertex with uniform weight

Conclusion

result is bad when tessellation is irregular

Conclusion


result is bad when tessellation is irregular



Conclusion


result is bad when tessellation is irregular

- reflection




Conclusion

result is bad when tessellation is irregular



Conclusion



result is bad when tessellation is irregular



Conclusion

result is bad when tessellation is irregular


■ burst



Conclusion

result is bad when tessellation is irregular

■ spreading



Ask more questions

- Pipeline:

$weight1 * principal_curvature + weight2 * feature_line \rightarrow$
 $constrain \rightarrow metric \rightarrow frame_field \rightarrow quad$

- How to reduce weight of bad *principal_curvature*?

- 1 conflict with feature

- 2 “flat/sphere like” region

current solution: $weight = \min(|e^{pc_1 - pc_2}| - 1, 100)$, still fail
on the previous big triangle.

- 3 other solution: subdivide or other way to change the bad
tessellation.

Ask more questions

- Pipeline:

$weight1 * principal_curvature + weight2 * feature_line \rightarrow$
 $constrain \rightarrow metric \rightarrow frame_field \rightarrow quad$

- How to reduce weight of bad *principal_curvature*?

- 1 conflict with feature


- 2 “flat/sphere like” region

current solution: $weight = \min(|e^{pc_1 - pc_2}| - 1, 100)$, still fail
on the previous big triangle.

- 3 other solution: subdivide or other way to change the bad
tessellation.

- what is the difference between *metric* and *frame_field*?

Other burst



Reference



Mark Meyer, Mathieu Desbrun, Peter Schröder, and Alan H Barr, *Discrete differential-geometry operators for triangulated 2-manifolds*, Visualization and mathematics III, Springer, 2003, pp. 35–57.