




## AP Practice Problems on the FTC

Graph of  $f$ 

5. Let  $f$  be a function defined on the closed interval  $[0, 7]$ . The graph of  $f$ , consisting of four line segments, is shown above. Let  $g$  be the function given by  $g(x) = \int_2^x f(t) dt$ .
- Find  $g(3)$ ,  $g'(3)$ , and  $g''(3)$ .
  - Find the average rate of change of  $g$  on the interval  $0 \leq x \leq 3$ .
  - For how many values  $c$ , where  $0 < c < 3$ , is  $g'(c)$  equal to the average rate found in part (b)? Explain your reasoning.
  - Find the  $x$ -coordinate of each point of inflection of the graph of  $g$  on the interval  $0 < x < 7$ . Justify your answer.




4. Let  $f$  be a function defined on the closed interval  $-5 \leq x \leq 5$  with  $f(1) = 3$ . The graph of  $f'$ , the derivative of  $f$ , consists of two semicircles and two line segments, as shown above.
- For  $-5 < x < 5$ , find all values  $x$  at which  $f$  has a relative maximum. Justify your answer.
  - For  $-5 < x < 5$ , find all values  $x$  at which the graph of  $f$  has a point of inflection. Justify your answer.
  - Find all intervals on which the graph of  $f$  is concave up and also has positive slope. Explain your reasoning.
  - Find the absolute minimum value of  $f(x)$  over the closed interval  $-5 \leq x \leq 5$ . Explain your reasoning.



Graph of  $g'$

5. Let  $g$  be a continuous function with  $g(2) = 5$ . The graph of the piecewise-linear function  $g'$ , the derivative of  $g$ , is shown above for  $-3 \leq x \leq 7$ .
- Find the  $x$ -coordinate of all points of inflection of the graph of  $y = g(x)$  for  $-3 < x < 7$ . Justify your answer.
  - Find the absolute maximum value of  $g$  on the interval  $-3 \leq x \leq 7$ . Justify your answer.
  - Find the average rate of change of  $g(x)$  on the interval  $-3 \leq x \leq 7$ .
  - Find the average rate of change of  $g'(x)$  on the interval  $-3 \leq x \leq 7$ . Does the Mean Value Theorem applied on the interval  $-3 \leq x \leq 7$  guarantee a value of  $c$ , for  $-3 < c < 7$ , such that  $g''(c)$  is equal to this average rate of change? Why or why not?



Graph of  $v$

4. A particle moves along the  $x$ -axis so that its velocity at time  $t$ , for  $0 \leq t \leq 6$ , is given by a differentiable function  $v$  whose graph is shown above. The velocity is 0 at  $t = 0$ ,  $t = 3$ , and  $t = 5$ , and the graph has horizontal tangents at  $t = 1$  and  $t = 4$ . The areas of the regions bounded by the  $t$ -axis and the graph of  $v$  on the intervals  $[0, 3]$ ,  $[3, 5]$ , and  $[5, 6]$  are 8, 3, and 2, respectively. At time  $t = 0$ , the particle is at  $x = -2$ .
- For  $0 \leq t \leq 6$ , find both the time and the position of the particle when the particle is farthest to the left. Justify your answer.
  - For how many values of  $t$ , where  $0 \leq t \leq 6$ , is the particle at  $x = -8$ ? Explain your reasoning.
  - On the interval  $2 < t < 3$ , is the speed of the particle increasing or decreasing? Give a reason for your answer.
  - During what time intervals, if any, is the acceleration of the particle negative? Justify your answer.