

EPITA

Mathematics

Midterm exam S3

November 2022

Duration: 3 hours

Name:

First name:

Class:

MARK:

The marking system is for a mark from 0 to 40. It will be divided by 2, to get a mark from 0 to 20.

Instructions:

- Documents and pocket calculators are not allowed.
 - Write your answers on the stapled sheets provided for answering. No other sheet will be corrected.
 - Please, do not use lead pencils for answering.
-

[illegible]

Let $a \in \mathbb{R}$ such that $a > 0$ and consider the sequence (u_n) defined for all $n \geq 2$ by:
$$u_n = \frac{(-1)^n}{\sqrt{n^a + (-1)^n}}.$$

1. Find $c \in \mathbb{R}$ such that $u_n = \frac{(-1)^n}{n^{a/2}} + \frac{c}{n^{3a/2}} + o\left(\frac{1}{n^{3a/2}}\right)$.

Exercise 3 (8 points)

The purpose of this exercise is to study the sequence (u_n) defined for all $n \in \mathbb{N}^*$ by: $u_n = \frac{2^n n!}{1 \times 3 \times 5 \times \cdots \times (2n-1)}$.

In that purpose, consider the auxiliary sequence (v_n) defined by: $v_n = \ln(u_n) - \frac{1}{2} \ln(n)$.

1. Let $n \in \mathbb{N}^*$. Compute $\frac{u_{n+1}}{u_n}$.

.....

.....

.....

.....

2. Deduce that $v_{n+1} - v_n = \frac{1}{2} \ln\left(1 + \frac{1}{n}\right) - \ln\left(1 + \frac{1}{2n}\right)$.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

3. Find $a \in \mathbb{R}$ such that $v_{n+1} - v_n \sim \frac{a}{n^2}$.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

4. Show that (v_n) converges. Let ℓ denote its limit.

.....

.....

.....

.....

.....

.....

.....

.....

Let (u_n) be a numerical sequence with an alternating sign.

[illegible]

3. Compute the expectation and the variance of Y .

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Exercise 6: power series (10 points)

Our purpose is to find a function f satisfying the following conditions $(C) : \begin{cases} f'' + xf' + f = 0 \\ f(0) = 1 \text{ and } f'(0) = 0 \end{cases}$

In that purpose, assume there exists a power series $\sum a_n x^n$, admitting a radius of convergence $R > 0$, such that:

$$f(x) = \sum_{n=0}^{+\infty} a_n x^n \quad \text{and} \quad f \text{ solution of } (C) \text{ on }]-R, R[$$

Note: the differential equation $f'' + xf' + f = 1$ is an order 2 linear equation with non-constant coefficients. Thus, **don't try to use the methods that you have studied at the S2** about order 2 differential equations: the latter methods only work for constant coefficients equations.

1. Express $f(0)$ and $f'(0)$ as functions of the sequence (a_n) . What can you deduce about the values of a_0 and a_1 ?

.....

.....

.....

.....

.....

2. Define, as functions of (a_n) , two sequences (b_n) and (c_n) such that for all $x \in]-R, R[$,

$$xf'(x) = \sum_{n=0}^{+\infty} b_n x^n \quad \text{and} \quad f''(x) = \sum_{n=0}^{+\infty} c_n x^n$$

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

3. By injecting these expressions of $xf'(x)$ and $f''(x)$ into the equation $f''(x) + xf'(x) + f(x) = 0$, write this equation as:

$$\forall x \in]-R, R[, \sum_{n=0}^{+\infty} d_n x^n = 0$$

where the coefficients (d_n) depend on the sequence (a_n) .

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

4. Keep in mind that the condition $\sum_{n=0}^{+\infty} d_n x^n = 0$ implies that all the d_n equal zero. Using this property, show that

$$a_2 = -\frac{1}{2}, \qquad a_3 = 0 \qquad \text{and, as a general rule:} \qquad \forall \in \mathbb{N}, a_{n+2} = -\frac{a_n}{n+2}$$

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

5. What is the value of a_n when n is odd?

.....

.....

.....

6. Find the value of a_n when n is even.

Hint: you can set $n = 2k$ ($k \in \mathbb{N}$). To start with, express a_{2k} as a function of $a_{2(k-1)}$, then as a function of $a_{2(k-2)}$, and so on, until you get an expression of a_{2k} as a function of a_0 .

.....

.....

.....

.....

.....

.....

.....

.....

7. Deduce $f(x)$. First, express it as a power series, then find an expression with the usual functions.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

8. **(Bonus)** Check that the final expression that you got at previous question is a solution of (C) on the whole set \mathbb{R} .

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....