Problem 2 – MerLean

Abstract

We consider recovery of a conductivity from a single solution of Darcy's equation on the unit ball, observed in Gaussian white noise of variance n^{-1}. For a sufficiently small fixed Hölder–Zygmund ball of smoothness $\alpha > 1$ around the constant conductivity, we prove that the minimax squared L^2 risk has polynomial order $n^{-2\alpha/(2\alpha+d+3)}$ in every dimension $d \geq 2$. The upper bound differs from the lower bound by at most $(\log n)^2$. The parameter class includes integer smoothness indices and has no prescribed conductivity trace. The lower bound uses localized angular perturbations and uniform nonlinear comparison. For the upper bound, inversion along a contracting gradient flow yields an integrated variance estimate that remains valid at the interior critical point. An independent pilot, a finite-dimensional correction, and boundary extrapolation give an estimator on the whole ball.

1 The problem and its rate

Let $D = \{x \in \mathbb{R}^d : |x| < 1\}$, where $d \geq 2$. For a positive conductivity f, let u_f solve

$$\operatorname{div}(f \nabla u_f) = 1 \quad \text{in } D, \qquad u_f = 0 \quad \text{on } \partial D. \tag{1}$$

We use the weak solution: $u_f \in H_0^1(D)$ and $\int_D f \nabla u_f \cdot \nabla \varphi = -\int_D \varphi$ for every $\varphi \in H_0^1(D)$. Uniform positivity of f gives existence and uniqueness by the coercive energy form.

Write $C_*^s(D) = B_{\infty,\infty}^s(D)$ for the inhomogeneous Hölder–Zygmund restriction space. Thus its norm is the infimum of the global $B_{\infty,\infty}^s(\mathbb{R}^d)$ norms of extensions; a concrete definition is given in Section 2. For noninteger $s > 0$ this is the usual Hölder space, with an equivalent norm. At integer indices we retain the Zygmund convention. Fix

$$\mathcal{F}(\alpha, r) = \{f : \|f - 1\|_{C_*^\alpha(D)} \leq r\}, \qquad \alpha > 1. \tag{2}$$

The observation is the Gaussian process

$$Y_n(\varphi) = \langle u_f, \varphi \rangle_{L^2(D)} + n^{-1/2} W(\varphi), \qquad \varphi \in L^2(D), \tag{3}$$

where W is isonormal: it is centered Gaussian with covariance $\mathbb{E}W(\varphi)W(\psi) = \langle \varphi, \psi \rangle_{L^2(D)}$. Our loss is squared error in the conductivity, and the minimax risk is

$$R_n(\alpha, r) = \inf_{\widehat{f}} \sup_{f \in \mathcal{F}(\alpha,r)} \mathbb{E}_f \|\widehat{f} - f\|_{L^2(D)}^2,$$

where the infimum is over measurable $L^2(D)$-valued estimators based on (3).

Theorem 1.1. *For each integer $d \geq 2$ and each $\alpha > 1$, there is $r_0 = r_0(d, \alpha) > 0$ such that, for every fixed $0 < r < r_0$, there are constants $c, C > 0$ with*

$$c\, n^{-2\alpha/(2\alpha+d+3)} \leq R_n(\alpha, r) \leq C (\log n)^2 n^{-2\alpha/(2\alpha+d+3)} \tag{4}$$

for all sufficiently large n. The constants may depend on d, α, r and the fixed norm convention. No boundary values of f are supplied.

The geometry of the solution matters for this rate. At $f = 1$,

$$u_1(x) = \frac{|x|^2 - 1}{2d}, \qquad \nabla u_1(x) = \frac{x}{d}. \tag{5}$$

Its gradient vanishes at the origin. Moreover, angular oscillations supported in a fixed annulus are attenuated by two powers of their frequency, while there are order J^{d-1} disjoint angular perturbations at frequency J. These two facts lead to the denominator $2\alpha + (d-1) + 4$ in (4). A fixed-direction linearization alone does not establish a minimax lower bound on a fixed nonlinear parameter ball; Section 3 controls the nonlinear errors uniformly over the whole testing family.

The upper bound uses a different aspect of the same geometry. For small r, every u_f is strictly convex. Inversion of $a \mapsto \operatorname{div}(a\nabla u_f)$ along the inward gradient flow is bounded on L^2, including at its unique critical point. After smoothing at scale h, averaging along this flow reduces the integrated noise variance from order $n^{-1}h^{-d-4}$ to $n^{-1}h^{-d-3}$. The estimator must account for the unknown flow and the nonlinear dependence on the noise. We therefore construct a pilot from an independent part of the observation, and bound the inverse of a finite-dimensional correction before taking noise expectations.

Related work. Nickl, van de Geer and Wang [3] study penalized least squares in elliptic coefficient problems with Gaussian white noise. Their Section 3.2 gives prediction bounds, a matching lower bound for estimating u_f in H^2, and a conductivity upper bound obtained through stability; see their Theorems 9–11. Those theorems use Sobolev classes with higher smoothness and prescribed boundary jets in (3.1). Prediction loss and squared conductivity loss are distinct, so those results do not give (4). Nickl and Paternain [4, Sections 3.1–3.3] analyze the linearized Darcy map and illustrate its critical-point geometry by an interior minimum on the disk. Their principal model class is a smooth Sobolev neighborhood, and their differentiability theorem is stated for $d \le 3$. Here the flow estimates are proved directly for the full class (2) in all fixed dimensions $d \ge 2$.

The elliptic regularity inputs are established results. We use Johnsen's boundary regularity theory in Besov spaces [2, Theorem 5.2 and Corollary 1.2] to include integer Zygmund indices, and the global gradient estimate of Breit, Cianchi, Diening and Schwarzacher [1, Corollary 2.7] for the nonlinear testing family. The remaining approximation, transport and Gaussian estimates are proved below. The conclusion is local in the conductivity radius and determines the polynomial rate; no optimality of the logarithmic factor is asserted.

2 Regularity and smoothing

Throughout, C denotes a finite constant that may change between occurrences, but is independent of n, h, the frequency J, and the member of (2). We write $a \lesssim b$ when $a \le Cb$ and $a \asymp b$ for two such inequalities. Norms without a displayed domain refer to the domain under discussion.

2.1 Function spaces and elliptic estimates

Choose a smooth inhomogeneous dyadic partition of unity in frequency, and let H_j denote its Fourier multiplier blocks, with $j = 0$ the low-frequency block and $j \ge 1$ supported where $|\xi| \asymp 2^j$. For $s \in \mathbb{R}$,

$$\|H\|_{C^s_*(\mathbb{R}^d)} = \sup_{j \ge 0} 2^{js} \|H_j\|_\infty.$$

Changing this fixed partition gives an equivalent norm. For $s > 0$, an equivalent global norm is

$$\|H\|_\infty + \sup_{0<|z|\le 1} |z|^{-s} \|\Delta_z^N H\|_\infty, \qquad N \in \mathbb{N}, \quad N > s, \tag{6}$$

where $\Delta_z H(x) = H(x+z) - H(x)$ and Δ_z^N is its N-fold iterate. To see the relevant estimates, each block satisfies

$$\|D^k H_j\|_\infty \lesssim 2^{jk} \|H_j\|_\infty, \qquad \|\Delta_z^N H_j\|_\infty \lesssim \min\{1, (2^j|z|)^N\} \|H_j\|_\infty.$$

Summing at $2^j \asymp |z|^{-1}$ gives the forward bound in (6). The reverse follows by writing an annular block as a finite sum of smooth multipliers applied to differences at scale 2^{-j}; on its frequency support one chooses coordinate increments for which the difference multiplier is bounded away from zero. The inverse multipliers have uniformly bounded convolution L^1 norms after rescaling.

Derivatives lower the index by their order. For $s > 0$, decomposition of a product into low–high, high–low and comparable-frequency terms gives

$$\|FG\|_{C_*^s} \lesssim \|F\|_\infty \|G\|_{C_*^s} + \|G\|_\infty \|F\|_{C_*^s}. \tag{7}$$

In the first two sums the low-frequency factor is bounded in L^∞; in the last, the tail above an output scale is summable because $s > 0$. Restriction norms inherit these estimates by taking extensions with norm within a fixed factor of the infimum. In particular, $C_*^s(D) \subset C^k(\overline{D})$ whenever $s > k$, as the derivative block series then converges uniformly.

Let $P = (-\Delta_D)^{-1}$ be the zero-Dirichlet inverse. Energy and Poincaré inequalities give

$$\|PS\|_2 + \|\nabla PS\|_2 \lesssim \|S\|_2, \qquad \|P \operatorname{div} F\|_{H^1} \lesssim \|F\|_2. \tag{8}$$

The divergence inverse in this display is defined weakly.

Lemma 2.1. *For $s > 1$ and $F \in C_*^s(D; \mathbb{R}^d)$,*

$$\|P \operatorname{div} F\|_{C_*^{s+1}(D)} \le C_s \|F\|_{C_*^s(D)}. \tag{9}$$

There is also $\beta_0 > 0$, depending only on d, such that for $0 < \beta < \min(1, \beta_0)$,

$$\|\nabla P \operatorname{div} F\|_{C^\beta(D)} \le C_\beta \|F\|_{C^\beta(D)}. \tag{10}$$

Here C^β has its ordinary full Hölder norm.

Proof. We specify the application of the two regularity results. For (9), apply [2, Theorem 5.2] to $(-\Delta, \gamma_0)$ on the smooth ball, where γ_0 is the boundary trace. In that theorem its order is 2, its class is 1, and its row and column weights are $b = (0, -2)$ and $a = (0)$. The resulting mapping is

$$B_{p,q}^t(D) \longrightarrow B_{p,q}^{t-2}(D) \times B_{p,q}^{t-1/p}(\partial D), \qquad t > 1 + \max(1/p - 1, d/p - d).$$

The interior symbol is $|\xi|^2$. At nonzero tangential frequency, a decaying homogeneous half-space solution is $ce^{-|\xi'|x_d}$, whose boundary trace determines c. This verifies ellipticity of the Dirichlet boundary problem. Differential operators have the transmission property required there.

The weak solution $w = P \operatorname{div} F$ starts in $H_0^1(D) \subset B_{2,2}^1(D)$, an admissible triple since the threshold is $1/2$. Its data lie in $B_{\infty,\infty}^{s-1}(D)$ with zero boundary trace. The target triple $(s+1, \infty, \infty)$ is admissible, its threshold being zero. The inverse-regularity clause of Theorem 5.2, referring to Corollary 1.2, therefore gives $w \in C_*^{s+1}(D)$. These are the restriction spaces defined in [2,

(2.21)–(2.22)]. The solution map is bounded by the closed graph theorem: convergence of F_j in C^s_* implies convergence of the solutions in H^1_0 by (8); convergence in C^{s+1}_* has the same weak limit. Thus its graph is closed. No density of smooth functions in $B^s_{\infty,\infty}$ is used.

For (10), specialize [1, Corollary 2.7] to the scalar case, exponent $p = 2$, dimension d, and the smooth ball. Its equation is $-\Delta w = -\operatorname{div} F$, so replace the datum by $-F$. The permitted range is $0 < \beta < \beta_0(d, 1, 2)$. Its Hölder bound is a seminorm bound, under the convention (2.2) of that paper. Since w has zero trace, $\int_D \nabla w = 0$. Averaging $\nabla w(x) - \nabla w(y)$ over $y \in D$ controls the supremum by the seminorm and proves the full-norm estimate. □

Proposition 2.2. *After choosing r_0 sufficiently small, every $f \in \mathcal{F}(\alpha, r)$, $r < r_0$, satisfies $1/2 \le f \le 3/2$ and*

$$\|u_f - u_1\|_{C^{\alpha+1}_*(D)} \le Cr, \qquad \|u_f - u_1\|_{C^2(\overline{D})} \le Cr. \tag{11}$$

In particular, we may arrange

$$\frac{3}{4d} I \le D^2 u_f \le \frac{5}{4d} I, \qquad \nabla u_f \cdot \nu \ge \frac{5}{8d} \quad \text{on } \partial B(0,R), \quad 3/4 \le R \le 1. \tag{12}$$

Proof. Writing $q = f - 1$ and $w = u_f - u_1$, the equation is

$$w = P \operatorname{div}\{q \nabla (u_1 + w)\}. \tag{13}$$

On the closed zero-trace subspace of $C^{\alpha+1}_*(D)$, Lemma 2.1 and (7) bound the linear part in w by Cr. For $Cr < 1/2$ this is a contraction, with solution of norm at most Cr. Its weak equation identifies it with $u_f - u_1$ by uniqueness. The embedding into C^2 follows by summing $2^{-j(\alpha-1)}$. Reducing r_0 makes the Hessian perturbation at most $1/(4d)$ in operator norm and the gradient perturbation at most $1/(8d)$. Equation (5) gives (12). The supremum embedding similarly gives the conductivity bounds. □

2.2 A smoothing residual with the required bias

Choose a real kernel $K \in C^\infty_c(B(0,1))$ of mass one with all nonconstant moments through an integer $L > 2\alpha + 2$ equal to zero. Such a signed kernel can be made from $L + 1$ distinct dilations of one smooth bump: solve the Vandermonde equations $\sum_i a_i \lambda_i^j = \mathbf{1}_{\{j=0\}}$, $0 \le j \le L$, with dilation factors small enough to preserve the support. Set

$$K_h(x) = h^{-d} K(x/h), \quad \Omega_h = B(0, 1-2h), \quad \Lambda_h = \log(e/h), \quad f_h = K_h * f, \quad u_h = K_h * u_f.$$

We always take $0 < h < 1/8$. These convolutions and their derivatives on Ω_h use only values in D; extensions are needed only to bound them.

For a global $H \in C^t_*$ and $0 < t < L + 1$, Taylor cancellation on a dyadic block gives

$$\|(K_h * -I) H_j\|_\infty \lesssim \min\{(h 2^j)^{L+1}, 1\} \|H_j\|_\infty.$$

Splitting the sum at $2^j \asymp h^{-1}$ proves

$$\|K_h * H - H\|_\infty \lesssim h^t \|H\|_{C^t_*}, \tag{14}$$

including integer t. Derivative versions follow by applying the same estimate to derivatives. Consequently,

$$\|f_h - f\|_{L^2(\Omega_h)} \lesssim h^\alpha, \quad \|f_h\|_{W^{1,\infty}(\Omega_h)} \lesssim 1, \quad \|u_h - u_f\|_{C^k(\overline{\Omega}_h)} \lesssim h^{\alpha+1-k}, \quad k = 1, 2. \tag{15}$$

Lemma 2.3. *Uniformly over $f \in \mathcal{F}(\alpha, r)$,*

$$r_h := \operatorname{div}(f_h \nabla u_h) - 1, \qquad \|r_h\|_{L^2(\Omega_h)} \lesssim h^\alpha \Lambda_h. \tag{16}$$

Proof. For $F \in C_*^\alpha$ and $G \in C_*^{\alpha-1}$ of bounded norm define

$$\mathcal{C}_h(F, G) = K_h * (FG) - (K_h * F)(K_h * G).$$

With $\delta_z F(x) = F(x - z) - F(x)$ and similarly for G, the exact identity is

$$\mathcal{C}_h(F, G) = \int K_h(z) \delta_z F \, \delta_z G \, dz - (K_h * F - F)(K_h * G - G). \tag{17}$$

Let $k = \lceil \alpha \rceil - 1$ and let ϵ_α be 1 if α is an integer and 0 otherwise. The increment Taylor polynomials of F, G have degrees 1 through k and 1 through $k-1$, respectively; the second is zero if $k = 1$. Their remainders are bounded by

$$C|z|^\alpha \log(e/|z|)^{\epsilon_\alpha}, \qquad C|z|^{\alpha-1} \log(e/|z|)^{\epsilon_\alpha}.$$

For clarity, these endpoint Taylor estimates also follow from the dyadic blocks. For noninteger t, summing derivatives gives the usual Hölder remainder. If t is an integer, derivatives through $t-1$ exist continuously; their top-order increments are bounded by $C|z| \log(e/|z|)$. The blocks below $|z|^{-1}$ contribute at most $C|z|$ each, and their number is logarithmic; the high-frequency tail is geometric. Taylor's integral formula then gives the asserted remainder of order t.

The product of the two increment polynomials has no constant term and degree at most $2k-1$, so its integral against K_h vanishes. The largest mixed remainder term is bounded by $Ch\, h^{\alpha-1} \Lambda_h^{\epsilon_\alpha}$, since F is Lipschitz. The product of remainders is bounded by $Ch^{2\alpha-1} \Lambda_h^{2\epsilon_\alpha}$, which is at most $Ch^\alpha \Lambda_h^{\epsilon_\alpha}$ because $\alpha > 1$. By (14), the last product in (17) is $O(h^{2\alpha-1})$. Thus

$$\|\mathcal{C}_h(F, G)\|_\infty \lesssim h^\alpha \Lambda_h. \tag{18}$$

On Ω_h, convolution of (1) now gives

$$r_h = -\mathcal{C}_h(f, \Delta u_f) - \sum_{i=1}^d \mathcal{C}_h(\partial_i u_f, \partial_i f).$$

Proposition 2.2 puts the first factor of each commutator in C_*^α and the second in $C_*^{\alpha-1}$. This proves (16). □

3 The lower bound: angular perturbations

We construct many perturbations in one fixed annulus, so that the radial direction does not contribute to their number. Put $m = d - 1$, $b_0 = (10\sqrt{m})^{-1}$, and use the sphere chart

$$\kappa(y) = (y, \sqrt{1 - |y|^2}), \qquad y \in (-2b_0, 2b_0)^m.$$

Choose nonzero real $\eta \in C_c^\infty((-1/4, 1/4)^m)$ and $a \in C_c^\infty((1/3, 2/3))$. For a large integer J, take centers $y_k \in J^{-1}\mathbb{Z}^m \cap (-b_0/2, b_0/2)^m$, $1 \le k \le M_J$, where $M_J \asymp J^m$. Define functions on the sphere by

$$\phi_k(\kappa(y)) = \eta(J(y - y_k)), \qquad \psi_k = J^{-2}(-\Delta_{\mathbb{S}^{d-1}})\phi_k, \tag{19}$$

extending by zero outside the chart. The supports are disjoint and separated. Uniform bounds on the chart metric give

$$\|\phi_k\|_2 \asymp J^{-m/2}, \quad \|\nabla_{\mathbb{S}^{d-1}}\phi_k\|_2 \asymp J^{1-m/2}, \quad \|\psi_k\|_2 \asymp J^{-m/2}. \tag{20}$$

For the last lower bound, integrate by parts to obtain $J^2\langle\phi_k,\psi_k\rangle = \|\nabla_{\mathbb{S}^{d-1}}\phi_k\|_2^2$ and apply Cauchy–Schwarz. This construction gives the cancellation needed below without a spectral-localization assumption.

For $\sigma \in \{-1,1\}^{M_J}$ set

$$\Phi_\sigma = \sum_k \sigma_k\phi_k, \quad \Psi_\sigma = \sum_k \sigma_k\psi_k, \quad q_\sigma(\rho\omega) = \varepsilon J^{-\alpha}a(\rho)\Psi_\sigma(\omega), \quad f_\sigma = 1 + q_\sigma. \tag{21}$$

Extend q_σ by zero to $\mathbb{R}^d$. Disjoint supports give, for each integer $s \geq 0$, $\|D^s q_\sigma\|_\infty \leq C_s\varepsilon J^{s-\alpha}$. For an integer $N > \alpha$, therefore,

$$\|\Delta_z^N q_\sigma\|_\infty \lesssim \varepsilon J^{-\alpha}\min\{1,(J|z|)^N\}.$$

Equation (6) gives $\|q_\sigma\|_{C_*^\alpha} \lesssim \varepsilon$, including integer α. Choose a fixed $\varepsilon > 0$, depending on r, small enough that all f_σ belong to (2) and $\|q_\sigma\|_\infty \leq 1/2$. If $H(\sigma,\tau)$ is their Hamming distance, orthogonality yields

$$\|f_\sigma - f_\tau\|_2^2 = 4\varepsilon^2 J^{-2\alpha}\left(\int a(\rho)^2\rho^{d-1}\,d\rho\right)\sum_{k:\sigma_k\neq\tau_k}\|\psi_k\|_2^2 \asymp \varepsilon^2 J^{-2\alpha-m}H(\sigma,\tau). \tag{22}$$

Only this testing subclass has conductivity equal to one near the boundary.

3.1 Linear cancellation and a uniform nonlinear bound

Let

$$T = I + d^{-1}x\cdot\nabla, \qquad A = PT.$$

For smooth h, differentiation of (1) at 1 gives the linear perturbation Ah. Equivalently, $-\Delta Ah = \operatorname{div}(h\nabla u_1) = Th$, with zero trace. We will use an exact identity for this operator. Define the fixed radial functions

$$g = a + \rho a'/d, \qquad c = \rho^2 g, \qquad B = c'' + (d-1)c'/\rho.$$

Polar differentiation gives

$$-\Delta(J^{-2}c\phi_k) = g\psi_k - J^{-2}B\phi_k, \qquad A(a\psi_k) = J^{-2}\{c\phi_k + P(B\phi_k)\}.$$

Thus

$$Aq_\sigma = \varepsilon J^{-\alpha-2}\{c\Phi_\sigma + P(B\Phi_\sigma)\}. \tag{23}$$

Energy estimates (8) imply that a single-sign difference $e = q_+ - q_-$ satisfies

$$\|Ae\|_2 \lesssim J^{-2}\|e\|_2, \qquad \|\nabla Ae\|_2 \lesssim J^{-1}\|e\|_2. \tag{24}$$

Indeed, the L^2 norm of $B\phi_k$ is $O(J^{-m/2})$, while differentiation of $c\phi_k$ costs at most one factor J.

Fix $0 < \beta < \min(1, \alpha-1, \beta_0)$ as allowed in Lemma 2.1. The disjoint packet bounds also give

$$\|q_\sigma\|_{C^\beta} \lesssim \varepsilon J^{-\alpha+\beta}, \quad \|B\Phi_\sigma\|_{C^\beta} \lesssim J^\beta, \quad \|\nabla(c\Phi_\sigma)\|_{C^\beta} \lesssim J^{1+\beta}.$$

To apply (10) to a scalar source $S = B\Phi_\sigma$, extended by zero, write it as $\operatorname{div} F$ with

$$F(x) = \left(\int_{-2}^{x_1} S(t, x_2, \dots, x_d)\, dt, 0, \dots, 0 \right).$$

Then $\|F\|_{C^\beta(D)} \lesssim \|S\|_{C^\beta(\mathbb{R}^d)}$. From (23),

$$\|\nabla A q_\sigma\|_{C^\beta} \lesssim \varepsilon J^{-\alpha-1+\beta}. \tag{25}$$

Write $w_\sigma = u_{f_\sigma} - u_1$. Its exact equation is

$$w_\sigma = A q_\sigma + P \operatorname{div}(q_\sigma \nabla w_\sigma). \tag{26}$$

On the zero-trace space $C^{1,\beta}$ with norm $\|\nabla w\|_{C^\beta}$, the last operator has norm at most $C\|q_\sigma\|_{C^\beta} \le C\varepsilon$. Reducing the fixed ε makes it a contraction of norm at most $1/2$. Its solution equals the weak solution by ellipticity. Hence

$$\sup_\sigma \|\nabla w_\sigma\|_\infty \lesssim \varepsilon J^{-\alpha-1+\beta}. \tag{27}$$

For neighboring signs put $\zeta = u_{f_+} - u_{f_-}$ and $R = \zeta - Ae$. Subtracting the PDEs gives, with zero trace,

$$\operatorname{div}((1+q_+)\nabla R) = -\operatorname{div}(e \nabla w_-) - \operatorname{div}(q_+ \nabla A e).$$

By energy, (24) and (27),

$$\begin{aligned} \|R\|_2 &\lesssim \|e\|_2 \|\nabla w_-\|_\infty + \|q_+\|_\infty \|\nabla A e\|_2 \\ &\lesssim J^{-2} \|e\|_2\, \varepsilon \{ J^{1+\beta-\alpha} + J^{1-\alpha} \}. \end{aligned} \tag{28}$$

Both exponents in braces are negative. Equations (22) and (24) now imply the uniform neighboring bound

$$\|u_{f_+} - u_{f_-}\|_2^2 \lesssim \varepsilon^2 J^{-2\alpha-m-4}. \tag{29}$$

3.2 Gaussian testing

The Gaussian shift experiment (3) has exact relative entropy $\mathrm{KL}(\mathbb{P}_{f_+}, \mathbb{P}_{f_-}) = (n/2)\|u_{f_+} - u_{f_-}\|_2^2$. Choose J to be the ceiling of a sufficiently large constant times $n^{1/(2\alpha+m+4)}$. Then (29) makes each neighboring relative entropy at most $1/8$, and Pinsker's inequality gives total variation at most $1/4$.

Here is the full reduction from testing to loss. Put $h_k = \varepsilon J^{-\alpha} a \psi_k$; these functions are orthogonal in $L^2(D)$. For any estimator $\widehat{f}$, infer its kth sign from the sign of $\langle \widehat{f} - 1, h_k \rangle$, with a fixed convention at zero. A wrong sign entails squared projection error at least $\|h_k\|_2^2$. For two parameter vectors differing only at k, the sum of the two testing error probabilities is at least $1 - \mathrm{TV} \ge 3/4$. Averaging uniformly over all sign vectors therefore gives error probability at least $3/8$ per coordinate. Bessel's inequality and (20) yield

$$\sup_{f \in \mathcal{F}(\alpha, r)} \mathbb{E}_f \|\widehat{f} - f\|_2^2 \ge \frac{3}{8} \sum_{k=1}^{M_J} \|h_k\|_2^2 \gtrsim \varepsilon^2 J^{-2\alpha}.$$

Since $m + 4 = d + 3$, this proves the lower bound of Theorem 1.1.

4 Transport inversion and integrated variance

The upper bound requires an inverse that uses the gradient geometry without dividing by its magnitude. We prove that inverse first for a deterministic potential v.

Lemma 4.1. *Let $\Omega = B(0,R)$ with $3/4 \le R \le 1$, and suppose $v \in C^2(\overline{\Omega})$ satisfies*

$$\kappa I \le D^2 v \le K_0 I, \qquad \nabla v \cdot \nu > 0 \text{ on } \partial\Omega, \qquad \kappa > 0. \tag{30}$$

Let Φ_t be the flow of $-\nabla v$, and set

$$j_t(x) = \det D\Phi_t(x) = \exp\left\{-\int_0^t \Delta v(\Phi_s x)\, ds\right\}, \quad T_v a = \operatorname{div}(a\nabla v), \quad R_v g(x) = \int_0^\infty j_t(x) g(\Phi_t x)\, dt.$$

Then R_v extends to a bounded operator on $L^2(\Omega)$ with

$$\|R_v\|_{2\to 2} \le \frac{2}{d\kappa}. \tag{31}$$

For Lipschitz a, $R_v T_v a = a$. Also $T_v R_v g = g$ in distributions for $g \in L^2(\Omega)$. No boundary value of a is required.

Proof. Strict convexity and the outward gradient imply that the unique minimizer z_0 of v lies in Ω, with $\nabla v(z_0) = 0$. The inward flow remains in Ω for all $t \ge 0$. Strong convexity gives

$$|\Phi_t x - z_0| \le e^{-\kappa t}|x - z_0|, \qquad 0 < j_t(x) \le e^{-d\kappa t}. \tag{32}$$

For smooth bounded a,

$$\frac{d}{dt}\{j_t(x) a(\Phi_t x)\} = -j_t(x)(T_v a)(\Phi_t x).$$

The terminal value vanishes by (32), proving the first inverse identity. The same identity holds almost everywhere for Lipschitz a by the characteristic chain rule. Alternatively, approximate in $W^{1,2}$ by smooth functions and use the boundedness below.

For $S_t g = j_t(g \circ \Phi_t)$, changing variables $y = \Phi_t x$ gives

$$\|S_t g\|_2^2 = \int_{\Phi_t(\Omega)} j_t(\Phi_t^{-1} y)|g(y)|^2\, dy \le e^{-d\kappa t}\|g\|_2^2.$$

Minkowski's inequality proves (31) and defines the integral for every $g \in L^2$. Finally S_t is a strongly continuous semigroup on L^2, and its distributional generator is $-T_v$. Its semigroup identity gives $S_\tau R_v g = R_v g - \int_0^\tau S_t g\, dt$. Dividing by τ and letting $\tau \downarrow 0$ proves $T_v R_v g = g$ in distributions. □

For a coefficient $b \in W^{1,\infty}(\Omega_h)$, write $L_b z = \operatorname{div}(b\nabla z)$, where weak derivatives of b are understood. The Hilbert–Schmidt norm of an integral operator with kernel $H(x,y)$ is $(\int\int |H(x,y)|^2\, dy\, dx)^{1/2}$. Its square is the expected squared L^2 norm of the operator applied to isonormal white noise.

Lemma 4.2. *Suppose v satisfies (30) on Ω_h, with fixed bounds κ, K_0, and $\|b\|_{W^{1,\infty}} \le C_0$. The operator from $L^2(D)$ to $L^2(\Omega_h)$ defined by $R_v L_b(K_h * \cdot)$ satisfies*

$$\|R_v L_b K_h\|_{\mathrm{HS}}^2 \lesssim h^{-d-3}. \tag{33}$$

The constant is uniform under these bounds.

Proof. Its kernel is

$$H_h(x,y) = \int_0^\infty j_t(x)\big[b(\Phi_t x)\Delta K_h(\Phi_t x - y) + \nabla b(\Phi_t x)\cdot \nabla K_h(\Phi_t x - y)\big]\,dt.$$

The bracket has $L^2(dy)$ norm at most $Ch^{-d/2-2}$. Two such brackets have disjoint supports when their centers are more than $2h$ apart. Expanding the square and applying Cauchy–Schwarz in y therefore bounds the squared kernel norm by $Ch^{-d-4}I_h$, where

$$I_h = \int_{\Omega_h}\int_0^\infty\int_0^\infty j_s(x)j_t(x)\mathbf{1}_{\{|\Phi_s x - \Phi_t x|\le 2h\}}\,dt\,ds\,dx.$$

By symmetry take $t = s+\tau$. Put $r_s = |\Phi_s x - z_0|$. The reverse triangle inequality and (32) give

$$|\Phi_{s+\tau}x - \Phi_s x| \ge (1 - e^{-\kappa\tau})r_s.$$

If $r_s \ge 4h$, overlap forces $\tau \le Ch/r_s$. If $r_s < 4h$, use $j_{s+\tau}/j_s \le e^{-d\kappa\tau}$ without the indicator. In either case the τ integral is at most $Cj_s\min(1, h/r_s)$. Consequently,

$$\begin{aligned} I_h &\lesssim \int_0^\infty\int_{\Omega_h} j_s(x)^2 \min\left(1, \frac{h}{|\Phi_s x - z_0|}\right)dx\,ds \\ &\le C\int_0^\infty e^{-d\kappa s}ds \int_{\Omega_h}\min\left(1,\frac{h}{|y-z_0|}\right)dy \lesssim h. \end{aligned}$$

For the last inequality, the region of radius h contributes $O(h^d)$, and the remainder contributes at most $Ch\int_h^2 \rho^{d-2}\,d\rho = O(h)$ when $d \ge 2$. This explicitly includes a neighborhood of the critical point. Substitution proves (33). □

5 Approximation and an independent pilot

We next build the two auxiliary objects used by the estimator. A deterministic projection makes the correction finite dimensional and controls one derivative by h^{-1}. A pilot constructed from legitimate observation tests supplies the flow while remaining independent of the noise used in Lemma 4.2.

5.1 A projection without boundary constraints

Lemma 5.1. *There are deterministic spaces $V_h \subset W^{1,\infty}(\Omega_h)$ of dimension $O(h^{-d})$ and projections $Q_h : L^2(\Omega_h) \to V_h$ such that*

$$\|Q_h\|_{2\to 2} \lesssim 1, \qquad \|\nabla a\|_2 \lesssim h^{-1}\|a\|_2 \quad (a \in V_h). \tag{34}$$

For f_h from Section 2, let $f_0 = Q_h f_h$ and $\eta_h = f_h - f_0$. Then

$$\|\eta_h\|_2 + h\|\nabla\eta_h\|_2 \lesssim h^\alpha, \qquad \|f_0\|_{W^{1,\infty}} \lesssim 1. \tag{35}$$

The spaces impose no boundary values.

Proof. We give a construction to keep the boundary and stability properties explicit. Let $P_0 = \{x : \sum_i |x_i| < 1\}$, split into its 2^d sign simplices. Choose a smooth nondecreasing χ, zero for $t \le 1/(4\sqrt{d})$ and one for $t \ge 1/(2\sqrt{d})$, and define

$$F(0) = 0, \qquad F(x) = x\left[1 + \chi(|x|)\left(\frac{|x|_1}{|x|} - 1\right)\right], \qquad |x|_1 = \sum_i |x_i|.$$

Along $x = t\omega$, with $c = |\omega|_1 \in [1, \sqrt{d}]$, the new radius is $\rho = t[1 + \chi(t)(c-1)]$ and $\partial_t \rho = 1 + (c-1)(\chi + t\chi') \geq 1$. At the polytope boundary $t = 1/c$, it equals one. Thus F maps P_0 bijectively onto D. Its Jacobian is $(\rho/t)^{d-1}\partial_t \rho$, bounded above and below. On each sign simplex it is smooth with bounded derivatives of every fixed order; near zero it is the identity. The map and its inverse are globally Lipschitz. Hence $F_R = RF$, $R = 1 - 2h$, has the same uniform properties as a map onto Ω_h.

For completeness, a conforming mesh of uniformly bounded shape ratios can be chosen as follows. On each sign simplex set $z_i = |x_i|$ and $t_i = z_i + \cdots + z_d$. The simplex becomes $1 \geq t_1 \geq \cdots \geq t_d \geq 0$. Subdivide cubes of side $1/m_h$, $m_h = \lceil 1/h \rceil$, by ordering their coordinate fractions, and retain the simplices inside this ordered region. Pull them back by $z_i = t_i - t_{i+1}$, $t_{d+1} = 0$. The faces $z_i = 0$ correspond to $t_i = t_{i+1}$ (or $t_d = 0$), so meshes on adjacent sign simplices agree. There are only finitely many rescaled element shapes, their volumes are comparable to h^d, and a node belongs to a bounded number of elements.

Let S_h be the continuous piecewise polynomial space of degree $p = \lceil \alpha \rceil$, with every Lagrange node retained, including boundary nodes. Set $V_h = \{A \circ F_R^{-1} : A \in S_h\}$. For each global node i, choose one incident simplex S_i. Let ψ_i be the local polynomial dual to that node, meaning $\int_{S_i} \psi_i \varphi_j = \delta_{ij}$ for the local nodal basis functions. Inverting their positive definite mass matrix and rescaling gives

$$\|\psi_i\|_2 \lesssim |S_i|^{-1/2}, \quad \|\psi_i\|_1 \lesssim 1, \quad \int_{S_i} \psi_i = 1.$$

With φ_i now denoting the global nodal functions, define

$$c_i(g) = \int_{S_i} g(F_R y)\psi_i(y)\, dy, \qquad Q_h g = \left(\sum_i c_i(g)\varphi_i \right) \circ F_R^{-1}.$$

Local duality shows that Q_h is a projection. On each target element, Cauchy–Schwarz bounds its coefficients by the L^2 norm on a bounded collection of nearby elements times $h^{-d/2}$. Summing, using bounded overlap and the Jacobian bounds, proves the first estimate in (34). Polynomial inverse estimates on the finite reference shapes give the second. Continuity across faces ensures membership in $W^{1,\infty}$.

We verify the approximation order for the smoothed function. For each dyadic block of an extension H of f, rapid decay of the Fourier transform of K gives, for every fixed q and N,

$$\|D^q(K_h * H_j)\|_\infty \lesssim 2^{j(q-\alpha)}(1 + h2^j)^{-N}.$$

This follows by localizing the smooth Fourier multiplier to its annulus, rescaling, and bounding its inverse Fourier kernel in L^1 by finitely many derivatives. Summing with $q = p + 1 > \alpha$ yields $\|D^{p+1} f_h\|_\infty \lesssim h^{\alpha - p - 1}$; all lower derivatives are bounded by the same envelope for $h < 1$. On each sign simplex composition with F_R preserves this bound. Taylor expansion on an element therefore proves that the ordinary nodal interpolant $I_h f_h \in V_h$ has errors

$$\|f_h - I_h f_h\|_\infty \lesssim h^\alpha, \qquad \|\nabla(f_h - I_h f_h)\|_\infty \lesssim h^{\alpha - 1}.$$

The interpolants agree across faces because their node values agree. Since $Q_h I_h f_h = I_h f_h$, its local coefficient bounds applied to the remainder give the same estimates for $Q_h f_h$, proving the first part of (35).

Finally, Q_h preserves Lipschitz bounds uniformly. On a target element fix a point y_*. Every source simplex contributing to that element is within Ch of y_*, and, because $\int \psi_i = 1$, its coefficient differs from $(g \circ F_R)(y_*)$ by at most $Ch \operatorname{Lip}(g \circ F_R)$. The basis is a partition of unity and its gradients are $O(h^{-1})$. Subtracting that common constant before differentiating gives $\|Q_h g\|_{W^{1,\infty}} \lesssim \|g\|_{W^{1,\infty}}$. Apply this to (15) to finish the proof. □

5.2 Reflection and Gaussian bounds for the pilot

A convolution near the boundary must use only observations in D. We implement it through the adjoint of an explicit extension. Let $s = \alpha + 1 > 2$, choose an integer $M > s$, and put $J_0 = M + 1$. Choose coefficients λ_j, $1 \le j \le J_0$, satisfying

$$\sum_{j=1}^{J_0} \lambda_j (-j)^\ell = 1, \qquad 0 \le \ell \le M.$$

The system is invertible by the Vandermonde determinant. Fix $\delta_0 < 1/(4J_0)$ and a smooth cutoff η_0, equal to one near zero and vanishing before δ_0. Define $Eg = g$ in D, and in the outer collar put

$$Eg((1+t)\omega) = \eta_0(t) \sum_{j=1}^{J_0} \lambda_j g((1-jt)\omega), \quad 0 < t < \delta_0; \tag{36}$$

set it zero further away. The reflected radii are at least $3/4$, so radial changes of variables prove

$$\|E\|_{L^2(D)\to L^2(\mathbb{R}^d)} \lesssim 1. \tag{37}$$

Consequently its adjoint E^* sends every $L^2(\mathbb{R}^d)$ kernel to a legitimate test function on D.

The moment equations match all available normal derivatives at the boundary; tangential derivatives then agree as well. When s is noninteger, the restriction dyadic bound gives the usual $C^{\lfloor s \rfloor, s - \lfloor s \rfloor}$ envelope for u_f. When s is an integer, its derivatives through $s-1$ are continuous and the top derivatives have modulus $Ct\log(e/t)$, by the dyadic split in the proof of Lemma 2.3. Smooth reflections preserve these envelopes. In boundary coordinates an increment crossing the interface can be split at the interface point, so the same modulus holds for (36). Taylor expansion using the moments of K therefore gives

$$\|K_b * Eu_f - u_f\|_{C^k(\overline{D})} \lesssim b^{\alpha+1-k} \log(e/b), \qquad k = 1, 2. \tag{38}$$

This argument needs only these classical or logarithmic regularity envelopes; it does not assume that E is bounded in every integer Zygmund norm.

Introduce an auxiliary white noise W' independent of Y_n, and form

$$Y^{(1)} = Y_n + n^{-1/2} W', \qquad Y^{(2)} = Y_n - n^{-1/2} W'. \tag{39}$$

Their noises are independent and each has variance $2/n$, as follows by computing the joint Gaussian covariance. Denote the corresponding normalized independent white noises by W_1, W_2. For $k_{b,x}(t) = K_b(x - t)$, set

$$v(x) = Y^{(1)}(E^* k_{b,x}), \qquad x \in \overline{D}. \tag{40}$$

Its mean is $K_b * Eu_f$. On Ω_h also set

$$Z = K_h * Y^{(2)} = u_h + N, \qquad N = \sqrt{2/n}\, K_h * W_2. \tag{41}$$

Here $K_h * Y^{(2)}(x)$ means evaluation of $Y^{(2)}$ at $K_h(x - \cdot)$ restricted to D.

Lemma 5.2. *Suppose b and h are fixed positive powers of n^{-1}. For every $A_0 > 0$, with probability at least $1 - C_{A_0} n^{-A_0}$, uniformly over $f \in \mathcal{F}(\alpha, r)$,*

$$\|v - u_f\|_{C^k(\overline{D})} \le C_{A_0} \left[b^{\alpha+1-k} \log(e/b) + n^{-1/2} b^{-d/2-k} \sqrt{\log n} \right], \quad k = 1, 2, \tag{42}$$

$$h^{-1} \|\nabla N\|_\infty + \|\Delta N\|_\infty \le C_{A_0} n^{-1/2} h^{-d/2-2} \sqrt{\log n}. \tag{43}$$

The fields can be chosen to be smooth and separable.

Proof. By (37), a derivative of order k of the centered pilot has standard deviation at most $\sigma_k = Cn^{-1/2}b^{-d/2-k}$. Its increment standard deviation is bounded by $C\sigma_k \min(1, |x-y|/b)$, using one more derivative of the kernel. The same estimates hold at every fixed derivative order. Gaussian moment bounds and a continuity criterion give continuous versions of all these derivative fields; the $L^2(D)$-valued differentiation identities for the kernels identify a smooth modification.

Here a direct net argument gives the required tail, without an expectation bound left implicit. Cover a fixed cube containing $\overline{D}$ first by a grid of spacing b, then by successive refinements of spacing $b2^{-j}$. There are $O(b^{-d}2^{jd})$ points at level j, and each point differs from its parent by a Gaussian increment with standard deviation at most $C\sigma_k 2^{-j}$. Union bounds with thresholds

$$D_{A_0}\sigma_k\sqrt{\log n} \quad \text{on the first grid,} \qquad D_{A_0}\sigma_k 2^{-j}\sqrt{\log n + j} \quad \text{on level } j$$

have total failure probability $O(n^{-A_0})$ when D_{A_0} is sufficiently large: b^{-d} is a fixed power of n, and the Gaussian exponential absorbs both that power and 2^{jd}. The sum of the thresholds is $O(\sigma_k\sqrt{\log n})$. Continuity extends the conclusion to all points; a finite union treats derivatives through order two. Adding (38) proves (42). Apply the same argument to $K_h * W_2$ at orders one and two to obtain (43). □

6 The estimator and the upper bound

Choose

$$q = 2\alpha + d + 3, \qquad h = n^{-1/q}, \qquad b = h^a, \qquad a = \frac{1 + 1/\alpha}{2} \in (1/\alpha, 1). \tag{44}$$

Only sufficiently large n are relevant; for smaller n the estimator may be set to one. Write $\delta = u_h - v$. The pilot and smoothing bounds give

$$h^{-1}\|\nabla\delta\|_\infty + \|\Delta\delta\|_\infty = o(1) \tag{45}$$

on events of probability $1 - O(n^{-A_0})$, for any fixed A_0. We check this choice explicitly. The bias terms have powers $h^{\alpha-1}$, $h^{a\alpha-1}$ and $h^{a(\alpha-1)}$, all tending to zero since $a\alpha - 1 = (\alpha-1)/2 > 0$. With $n^{-1/2} = h^{q/2}$, the pilot noise powers are

$$q/2 - 1 - a(d/2+1), \qquad q/2 - a(d/2+2).$$

Both are at least $\alpha - 1/2 > 0$, since $a < 1$. The second noise in (43) has power $\alpha - 1/2$ as well. Thus all the stated convergences hold despite the logarithmic factors. In particular, v is uniformly C^2-close to u_f with this probability.

6.1 An observable finite-dimensional equation

First check from v whether

$$\frac{1}{2d}I \le D^2 v \le \frac{3}{2d}I \quad \text{on } \overline{\Omega}_h, \qquad \nabla v \cdot \nu \ge \frac{1}{2d} \quad \text{on } \partial\Omega_h. \tag{46}$$

If this fails, output the constant function one. Proposition 2.2 and Lemma 5.2 show that failure has probability $O(n^{-A_0})$.

When (46) holds, R_v is defined by Lemma 4.1. For a smooth z on $\overline{\Omega}_h$, define an operator on V_h by

$$B_z a = Q_h R_v \operatorname{div}(a\nabla z). \tag{47}$$

The projection and inverse inequalities imply, for a fixed deterministic constant C_B,

$$\|B_z\|_{2\to 2} \le C_B \left(h^{-1} \|\nabla z\|_\infty + \|\Delta z\|_\infty \right). \tag{48}$$

Indeed, $\operatorname{div}(a\nabla z) = \nabla a \cdot \nabla z + a\Delta z$ belongs to L^2; continuity of a excludes a face jump distribution.

Use $z = Z - v = \delta + N$. Check whether the observable right side of (48) is at most $1/2$. If not, output one. Otherwise solve the finite-dimensional equation

$$(I + B_{Z-v})\widehat{f}^{\,\mathrm{int}} = Q_h R_v 1 \quad \text{in } V_h. \tag{49}$$

Its inverse exists with norm at most two. Equations (45) and (43) imply that both checks succeed with probability at least $1 - C_{A_0} n^{-A_0}$. The unknown f_h, f_0 and δ are comparison quantities in the proof; only v, Z and the fixed constructions are used in (49).

All these operations are measurable. The smooth Gaussian fields have separable versions, so the suprema and geometry checks can be tested on countable dense sets. Fix a deterministic basis of V_h. Finite-time flows and Jacobians are measurable functions of the C^2 pilot; integrating fixed Borel representatives gives measurable matrix entries. The tail of R_v beyond time T has L^2 operator norm at most $(2/(d\kappa))e^{-d\kappa T/2}$, so its matrix entries are measurable limits of finite-time entries. Matrix inversion on the accepted event is measurable as well.

6.2 Risk on the interior

Recall $f_0 = Q_h f_h$, $\eta_h = f_h - f_0$ and $r_h = \operatorname{div}(f_h \nabla u_h) - 1$. Since f_h is smooth, the identity $R_v T_v f_h = f_h$ gives

$$f_0 + B_\delta f_0 = Q_h R_v 1 + Q_h R_v r_h - Q_h R_v \operatorname{div}(\eta_h \nabla \delta). \tag{50}$$

To verify it directly, write $T_{u_h} f_h = T_v f_h + \operatorname{div}(f_h \nabla \delta) = 1 + r_h$, apply $Q_h R_v$, and split $f_h = f_0 + \eta_h$ only in the second term. Subtracting (50) from (49) yields the exact error identity on the solving event:

$$\begin{aligned}(I + B_{\delta+N})(\widehat{f}^{\,\mathrm{int}} - f_0) = \; & - Q_h R_v r_h + Q_h R_v \operatorname{div}(\eta_h \nabla \delta) \\ & - Q_h R_v \operatorname{div}(f_0 \nabla N).\end{aligned} \tag{51}$$

The first term is $O(h^\alpha \Lambda_h)$ by Lemma 2.3. For the second,

$$\begin{aligned}\|Q_h R_v \operatorname{div}(\eta_h \nabla \delta)\|_2 &\lesssim \|\nabla \eta_h\|_2 \|\nabla \delta\|_\infty + \|\eta_h\|_2 \|\Delta \delta\|_\infty \\ &\lesssim h^\alpha\end{aligned}$$

on the high-probability pilot event, by (35) and (45).

Conditional on the pilot, W_2 is independent, while f_0 is a deterministic function of the fixed true f. Lemmas 4.2 and 5.1 imply on the geometry event

$$\mathbb{E}_f \left[\|Q_h R_v \operatorname{div}(f_0 \nabla N)\|_2^2 \mid v \right] \lesssim n^{-1} h^{-d-3}. \tag{52}$$

The solving event depends on N, but this causes no additional assumption in (52): first use the deterministic inverse bound two in (51); the resulting nonnegative noise square can then be integrated after dropping the solving-event indicator. Thus the variance bound is applied only to a linear operator whose flow is independent of W_2.

Combining these bounds with $\|f_0 - f\|_{L^2(\Omega_h)} \lesssim h^\alpha$ gives interior error, integrated over the good event, at most

$$C\{h^{2\alpha} \Lambda_h^2 + n^{-1} h^{-d-3}\}. \tag{53}$$

We now control the omitted boundary strip and bound the loss on all other events.

6.3 Boundary recovery and removal of randomization

Choose an integer $M_1 > \alpha$ and set $w_k = (-1)^{k-1}\binom{M_1}{k}$, $1 \le k \le M_1$. Define an extension from the interior ball by

$$E_h g = g \text{ on } \Omega_h, \qquad E_h g(\rho\omega) = \sum_{k=1}^{M_1} w_k g((\rho - 2kh)\omega), \quad 1 - 2h < \rho < 1. \tag{54}$$

For sufficiently small h, all sampled points lie in Ω_h and their radii exceed $1/2$. Radial changes of variables prove the uniform bound

$$\|E_h g\|_{L^2(D)} \le C\|g\|_{L^2(\Omega_h)}. \tag{55}$$

For example, each translated annular integral has density ratio $\rho^{d-1}/(\rho - 2kh)^{d-1}$ uniformly bounded; the number and weights of the terms are fixed.

The coefficient identity $\sum_k w_k(-2k)^j = \mathbf{1}_{\{j=0\}}$ for $0 \le j < M_1$ shows that, for a true f, the strip error in (54) is exactly

$$E_h(f|_{\Omega_h})(\rho\omega) - f(\rho\omega) = (-1)^{M_1+1}\Delta^{M_1}_{-2h\omega} f(\rho\omega).$$

Apply (6) to a global extension of f. It gives pointwise error Ch^α for every $\alpha < M_1$, including integer indices without a logarithm. Since the strip has volume $O(h)$,

$$\|E_h(f|_{\Omega_h}) - f\|_{L^2(D)} \lesssim h^{\alpha+1/2}. \tag{56}$$

On successful checks, extend $\widehat{f}^{\,\mathrm{int}}$ by (54) and clip its values to $[1/2, 3/2]$. On failed checks use one. Denote the resulting randomized full-domain estimator by $\widetilde{f}$. Clipping cannot increase squared error against any f in the class, and it bounds squared loss by $|D|$ on every event. Equations (53)–(56) and the probability bounds therefore give

$$\sup_{f \in \mathcal{F}(\alpha, r)} \mathbb{E}_f \|\widetilde{f} - f\|_2^2 \le C\{h^{2\alpha}\Lambda_h^2 + n^{-1}h^{-d-3} + n^{-A_0}\}. \tag{57}$$

Take $A_0 > 2\alpha/q$ and substitute (44) to obtain the upper rate in (4).

Finally, average $\widetilde{f}(Y_n, W')$ over the known law of the auxiliary W' for each fixed Y_n. This is a bounded $L^2(D)$-valued integral and gives a measurable estimator $\widehat{f}(Y_n)$ in the original experiment. Independence of W' from Y_n and Jensen's inequality show that its risk is no greater than that of $\widetilde{f}$. This completes the proof of Theorem 1.1.

Scope of the conclusion. The small-radius condition supplies convexity of the solution uniformly over the parameter class. The argument does not establish the same rate for arbitrary positive conductivities or for other critical-point geometries. The restriction $d \ge 2$ is used in the spatial overlap integral of Lemma 4.2, and $\alpha > 1$ is used both in the nonlinear testing remainder and in the pilot perturbation bounds. The estimator depends on α; adaptation is not considered. The theorem determines the polynomial rate on the stated fixed-radius class, leaving the sharp logarithmic behavior open.

References

[1] D. Breit, A. Cianchi, L. Diening and S. Schwarzacher, *Global Schauder estimates for the p-Laplace system*, Archive for Rational Mechanics and Analysis **243** (2022), 201–255. doi:10.1007/s00205-021-01712-w.

[2] J. Johnsen, *Elliptic boundary problems and the Boutet de Monvel calculus in Besov and Triebel–Lizorkin spaces*, Mathematica Scandinavica **79** (1996), 25–85. doi:10.7146/math.scand.a-12593. Author version: arXiv:1704.08555v1; theorem numbering as in that version.

[3] R. Nickl, S. van de Geer and S. Wang, *Convergence rates for penalized least squares estimators in PDE constrained regression problems*, SIAM/ASA Journal on Uncertainty Quantification **8** (2020), 374–413. doi:10.1137/18M1236137. Author version: arXiv:1809.08818v3.

[4] R. Nickl and G. P. Paternain, *On some information-theoretic aspects of non-linear statistical inverse problems*, Proceedings of the International Congress of Mathematicians 2022, Vol. 7, EMS Press, 5516–5538. doi:10.4171/ICM2022/18.