
Review of Problem 3 – MerLean



First Proof Third Batch
Reviewer 1
October 8–9, 2026

1 Review

The solution only claims a partial result. The solution engages with the algebraic part of the problem, and acknowledges that it does not attempt to engage with the skein theoretic portion.

Correctness. The paper acknowledges a partial result, and that partial result is the easier half of the main problem statement. But what the partial solution accomplished does appear to be correct.

Exposition. I had trouble finding the precise statement that serves as the objective of the partial result. In each section, I found it hard to deduce what goal was accomplished.

Attribution. There are too few papers cited. The main source for stated SL_n skein algebras should be Lê-Sikora, which is not included in the bibliography.

Correctness.

- 1: essentially flawless mathematical argument
- 0.5: minor errors that are easily corrected
- 0: blank solution, or acknowledged partial result
- -0.5: major errors
- -1: proposed proof makes no progress towards the result.

Note that points should be docked for wrong statements/ solutions.
Score: 0

Exposition.

- 1: well-written and clear solution.
- 0: argument that is possible to follow with serious effort
- -1: very poor exposition

Score: 0

Attributions.

- 1: solutions contains fully appropriate attributions to related work and theorems being used.
- 0: moderate issues with missing attributions or hallucinated references
- -1: blatant plagiarism or hallucinations

Score: 0

2 Submission with inline comments

Abstract

The problem considered here asks for an explicit identification of a reduced stated skein algebra of a punctured bigon with a generalized double of the full-weight quantum Borel algebras, followed by a diagonal quotient giving the simply-connected quantum group. We give a partial answer. At a specified generic parameter we compute the ordered double, its full diagonal quotient, and explicit inverse formulas for triangular quantum matrices. These yield a comparison theorem for an algebra satisfying a precise mixed exchange identity and a vector-space factorization. The proof retains the weight lattice and distinguishes gluing all weight Cartans from gluing only simple-root Cartans. The identification of these hypotheses with the designated skein algebra and its arc conventions remains unresolved; consequently neither requested isomorphism for the original objects is asserted.

3 The question and the scope of the reduction

Let $U_q(\mathfrak{b}_+)$ and $U_q(\mathfrak{b}_-)$ denote the upper and lower Borel subalgebras of the simply-connected quantum group $U_q(\mathfrak{sl}_n)$. The question is to construct

$$\overline{\mathscr{S}}_n(\mathfrak{B}^*) \;\cong\; U_q(\mathfrak{b}_-) \bowtie U_q(\mathfrak{b}_+) \tag{1}$$

for the prescribed Drinfeld–Tanisaki pairing

$$\begin{gathered}\langle \hat{K}_r, \hat{K}_u \rangle = q^{1/n-\delta_{ru}}, \qquad \langle E_i, F_j \rangle = \frac{\delta_{ij}}{q^{-1}-q}, \\ \langle \hat{K}_r, F_j \rangle = \langle E_i, \hat{K}_u \rangle = 0,\end{gathered} \tag{2}$$

and then to deduce

$$U_q(\mathfrak{sl}_n) \;\cong\; \overline{\mathscr{S}}_n(\mathfrak{B}^*)/(b_{rr}c_{rr} - 1 : 1 \le r \le n). \tag{3}$$

Here b_{ij} and c_{ij} are the two designated families of rightward horizontal stated arcs. A state is an endpoint label in $\{1, \dots, n\}$. Parentheses in an algebra quotient always denote the two-sided ideal generated by the listed elements.

An explicit solution requires more than a matching commutator. The endpoint order, boundary reduction and multiplication convention on the skein side must determine the same mixed relation as the ordered double. Injectivity must then be proved. Furthermore, the simply-connected form involves weight Cartans, not just simple-root Cartans.

The precise definition of the original $\overline{\mathscr{S}}_n(\mathfrak{B}^*)$, its designated arc diagram, and its coefficient and fractional-power conventions are not fixed here. We therefore keep (1)–(3) as the original targets and distinguish them from all algebras defined below. The results of this paper are a generic algebraic reduction of these targets. The missing geometric and convention identifications are stated in Section 9.

Our main comparison result is Theorem 8.1. It gives every triangular matrix entry as a root vector with its exact scalar and Cartan factor. Its hypotheses include an explicit mixed exchange identity and a multiplication factorization; it does not presume the desired isomorphism. The independent presentation calculation in Theorem 5.3 identifies the resulting algebra with a specified ordered double. Proposition 6.1 proves the diagonal quotient, and Proposition 6.2 explains why gluing only simple-root ratios is insufficient.

The full-weight presentations and pairing used here are recalled from Voigt–Yuncken [3, Definition 2.13, Proposition 2.14, and Section 2.8.1]; we explicitly convert their coproduct convention.

The inverse triangular-matrix calculation uses the type-A root relations recorded by Chen–Shao–Shum [1, Section 3], who attribute those relations to Yamane. We use the relations, rather than their basis theorem. On the geometric side, Lê–Yu [2, Theorems 6.2 and 7.2] provide arc factorization and normality of the ideals generated by bad corner arcs. Their Theorem 6.2 is credited to Lê–Sikora.

> **Reviewer 1**
>
> Lê–Sikora is mentioned, but should be cited in the bibliography.

Section 9 explains how such results could supply a factorization after reduction, without identifying the original diagram by assumption. No novelty claim is made for the underlying quantum-group theory.

4 Weight Cartans and the two presentations

Throughout the algebraic results, fix an integer $n \geq 2$, an indeterminate t, and

$$\mathbb{K} = \mathbb{Q}(t), \qquad q = t^{2n^2}, \qquad d = q - q^{-1}, \qquad q^{1/n} = t^{2n}. \tag{4}$$

Thus q is not a root of unity and $d \neq 0$. All tensor products are over $\mathbb{K}$ unless another base is specified. We use $1 \leq r, u \leq n$ for weight indices and $1 \leq i, j < n$ for simple-root indices. Set

$$w_{ri} = \delta_{ri} - \delta_{r,i+1}, \qquad a_{ij} = 2\delta_{ij} - \delta_{i,j+1} - \delta_{i+1,j}.$$

4.1 Why the determinant-one torus is the weight torus

Let $\varepsilon_1, \dots, \varepsilon_n$ be the standard basis of $\mathbb{Q}^n$ and put

$$\lambda_r = \varepsilon_r - \frac{1}{n}\sum_{u=1}^{n} \varepsilon_u, \qquad W = \{x \in \mathbb{Q}^n : \textstyle\sum_r x_r = 0,\ x_r - x_u \in \mathbb{Z} \text{ for all } r, u\}.$$

The lattice W, with the ordinary dot product, is the type-A_{n-1} weight lattice. Its simple roots are $\alpha_i = \lambda_i - \lambda_{i+1}$.

Lemma 4.1. *The map $[m] \mapsto \sum_r m_r \lambda_r$ identifies $\mathbb{Z}^n/\mathbb{Z}(1, \dots, 1)$ with W. The fundamental weights are $\omega_i = \lambda_1 + \cdots + \lambda_i$. Moreover,*

$$(\lambda_r, \lambda_u) = \delta_{ru} - 1/n, \qquad (\lambda_r, \alpha_i) = w_{ri}.$$

The least positive integer L for which $(W, W) \subseteq L^{-1}\mathbb{Z}$ is n.

Proof. The image of m has coordinates $m_r - \frac{1}{n}\sum_u m_u$. It is zero precisely when m is constant. Conversely, for $x \in W$ take the integers $m_r = x_r - x_n$; their image is x. The displayed inner products follow by expansion. They give $(\omega_i, \alpha_j) = \delta_{ij}$, while $\lambda_1 = \omega_1$, $\lambda_r = \omega_r - \omega_{r-1}$ for $1 < r < n$, and $\lambda_n = -\omega_{n-1}$. Thus these are all the fundamental weights and they generate W. For integer representatives m, l,

$$\left(\sum_r m_r\lambda_r, \sum_r l_r\lambda_r\right) = \sum_r m_r l_r - \frac{(\sum_r m_r)(\sum_r l_r)}{n}.$$

This belongs to $n^{-1}\mathbb{Z}$, and $(\lambda_1, \lambda_1) = (n-1)/n$ has denominator n. □

4.2 Borel and double presentations

Define B_+ to be the algebra generated by commuting units P_r and elements E_i, subject to

$$\prod_r P_r = 1, \qquad P_r E_i P_r^{-1} = q^{w_{ri}} E_i. \tag{5}$$

The root generators satisfy the type-A quantum Serre relations: for $X_i = E_i$,

$$\begin{aligned} X_i X_j &= X_j X_i \quad (|i-j|>1), \\ X_i^2 X_j - (q+q^{-1}) X_i X_j X_i + X_j X_i^2 &= 0 \qquad (|i-j|=1). \end{aligned} \tag{6}$$

Define B_- by commuting units N_r with $\prod_r N_r = 1$, generators F_i, the same root relations, and

$$N_r F_i N_r^{-1} = q^{-w_{ri}} F_i. \tag{7}$$

For $\lambda = \sum m_r \lambda_r$ write $P_\lambda = \prod P_r^{m_r}$ and $N_\lambda = \prod N_r^{m_r}$. Lemma 4.1 and the product-one relations make these well defined. Abbreviate

$$A_i = P_i P_{i+1}^{-1}, \qquad B_i = N_i N_{i+1}^{-1}.$$

Here the indexed element B_i is distinct from the algebra B_-.

Define $\mathcal{A}$ by the two Borel presentations together with

$$\begin{aligned} P_r N_u &= N_u P_r, & P_r F_i &= q^{-w_{ri}} F_i P_r, \\ N_r E_i &= q^{w_{ri}} E_i N_r, & [E_i, F_j] &= \delta_{ij} \frac{A_i - B_i^{-1}}{d}. \end{aligned} \tag{8}$$

As usual, $[x,y] = xy - yx$. Define $\mathcal{T}$ to have one family of commuting units K_r of product one, the roots E_i, F_i, their relations (6), weights $q^{w_{ri}}$ and $q^{-w_{ri}}$, and

$$[E_i, F_j] = \delta_{ij} \frac{L_i - L_i^{-1}}{d}, \qquad L_i = K_i K_{i+1}^{-1}. \tag{9}$$

These are definitions of auxiliary algebras, independent of any skein identification.

5 The pairing and the ordered double

We first realize $B_\pm$ as actual Hopf algebras. A Hopf algebra has a coproduct Δ, a counit ϵ, and an antipode S. We write $\Delta(x) = \sum x_1 \otimes x_2$ and $(\Delta \otimes \mathrm{id})\Delta(x) = \sum x_1 \otimes x_2 \otimes x_3$, suppressing summation indices. All such sums below are finite for a fixed element. A group-like element g satisfies $\Delta(g) = g \otimes g$ and $\epsilon(g) = 1$; its antipode is g^{-1}.

Proposition 5.1. *The presentations $B_\pm$ admit Hopf structures with group-like Cartans and*

$$\begin{aligned} \Delta(E_i) &= E_i \otimes 1 + A_i \otimes E_i, & S(E_i) &= -A_i^{-1} E_i, \\ \Delta(F_i) &= F_i \otimes B_i^{-1} + 1 \otimes F_i, & S(F_i) &= -F_i B_i. \end{aligned} \tag{10}$$

The Cartan counits are one and the root counits are zero. There is a bilinear pairing $\tau : B_+ \times B_- \to \mathbb{K}$ satisfying

$$\begin{aligned} \tau(xx', y) &= \sum \tau(x', y_1)\tau(x, y_2), \\ \tau(x, yy') &= \sum \tau(x_1, y)\tau(x_2, y'), \\ \tau(1, y) &= \epsilon(y), \qquad \tau(x, 1) = \epsilon(x), \end{aligned} \tag{11}$$

and

$$\begin{aligned} \tau(P_r, N_u) = q^{1/n-\delta_{ru}}, \qquad \tau(E_i, F_j) = -\delta_{ij}/d, \\ \tau(P_r, F_j) = \tau(E_i, N_u) = 0. \end{aligned} \tag{12}$$

Proof. Use the full-weight algebra of [3, Definition 2.13] for type A_{n-1}. The source rank is $n-1$, its squared root lengths are two, and its integer L is n by Lemma 4.1. Its parameter s can be taken to be t^{2n}, so its $q = s^L$ agrees with (4). This checks both the fractional-power convention and the non-root-of-unity hypothesis used for its pairing.

The paragraph after [3, Proposition 2.14] identifies the actual Borel subalgebras with their corresponding presentations. Send their K_λ to P_λ or N_λ, respectively, and fix the root generators. The inverse assignments are $P_r \mapsto K_{\lambda_r}$ and $N_r \mapsto K_{\lambda_r}$. Lemma 4.1 checks the torus relations; the weights are $(\lambda_r, \alpha_i) = w_{ri}$, and the type-A Serre relations are exactly (6). These are inverse presentation maps on the actual Borels.

Write Δ_0, S_0 for the transported source Hopf operations. Their root formulas are

$$\begin{aligned} \Delta_0(E_i) = E_i \otimes A_i + 1 \otimes E_i, \qquad \Delta_0(F_i) = F_i \otimes 1 + B_i^{-1} \otimes F_i, \\ S_0(E_i) = -E_i A_i^{-1}, \qquad S_0(F_i) = -B_i F_i. \end{aligned}$$

The square of S_0 scales E_i by q^2 and F_i by q^{-2} and fixes Cartans. These scalings are automorphisms, so S_0 is invertible. Replacing Δ_0 by its opposite coproduct and S_0 by S_0^{-1} gives (10). This operation is called taking the coopposite Hopf algebra; multiplication is unchanged.

By [3, Proposition 2.73], the source Borels possess a pairing $\rho : B_- \times B_+ \to \mathbb{K}$ with

$$\rho(N_\lambda, P_\mu) = q^{(\lambda,\mu)}, \qquad \rho(F_j, E_i) = \delta_{ij}/d$$

and both mixed root–torus values zero. Its two multiplication rules, using Δ_0, are

$$\begin{aligned} \rho(yy', x) = \sum \rho(y, x_{1,0})\rho(y', x_{2,0}), \\ \rho(y, xx') = \sum \rho(y_{1,0}, x')\rho(y_{2,0}, x). \end{aligned}$$

Define $\tau(x, y) = \rho(S_0(y), x)$, as in [3, Definition 2.74]. Since S_0 reverses multiplication and coproduct, these rules give

$$\begin{aligned} \tau(xx', y) = \sum \tau(x, y_{1,0})\tau(x', y_{2,0}), \\ \tau(x, yy') = \sum \tau(x_{1,0}, y')\tau(x_{2,0}, y). \end{aligned}$$

Reversing the coproduct indices gives (11).

For Cartans this construction gives $\tau(P_\lambda, N_\mu) = q^{-(\lambda,\mu)}$. The mixed zero values follow from the corresponding values of ρ. Finally,

$$\begin{aligned} \tau(E_i, F_j) &= -\rho(B_j F_j, E_i) \\ &= -\rho(B_j, E_i)\rho(F_j, A_i) - \rho(B_j, 1)\rho(F_j, E_i) = -\delta_{ij}/d. \end{aligned}$$

Thus existence precedes the calculation of the generator table; it is not inferred from that table alone. □

5.1 An associative product on the ordered tensor space

The chosen order of the double is negative followed by positive. We include the construction to specify its multiplication and to justify the tensor-space comparison used later.

Lemma 5.2. *The vector space $B_- \otimes B_+$ is an associative unital algebra $\mathcal{D}$ under*

$$(y \otimes x)(z \otimes u) = \sum \tau(x_1, z_1)\tau(x_3, S(z_3))\, yz_2 \otimes x_2 u. \tag{13}$$

Both factors embed as algebras and $(y \otimes 1)(1 \otimes x) = y \otimes x$.

Proof. Start with the ordinary tensor-product Hopf algebra $C = B_- \otimes B_+$. Define scalar-valued maps

$$\sigma(y \otimes x, z \otimes u) = \epsilon(y)\tau(x, z)\epsilon(u),$$
$$\bar{\sigma}(y \otimes x, z \otimes u) = \epsilon(y)\tau(x, S(z))\epsilon(u).$$

For maps on a coalgebra, convolution is $(f * g)(a) = \sum f(a_1)g(a_2)$. The second rule in (11) and the two antipode identities show that $\sigma * \bar{\sigma} = \bar{\sigma} * \sigma = \epsilon \otimes \epsilon$. The unit pairing gives $\sigma(a, 1) = \sigma(1, a) = \epsilon(a)$.

The required cocycle identity is

$$\sum \sigma(a_1, b_1)\sigma(a_2 b_2, c) = \sum \sigma(b_1, c_1)\sigma(a, b_2 c_2). \tag{14}$$

For $a = y \otimes x$, $b = z \otimes u$, $c = w \otimes v$, both sides reduce by (11) to

$$\epsilon(y)\epsilon(v) \sum \tau(x_1, z)\tau(u, w_1)\tau(x_2, w_2).$$

Pure tensors span C, so the identity holds generally.

Set $a \star b = \sum \sigma(a_1, b_1) a_2 b_2 \bar{\sigma}(a_3, b_3)$. To verify associativity explicitly, denote the left and right sides of (14) by $L(a, b, c)$ and $R(a, b, c)$. Their convolution inverses on $C^{\otimes 3}$ are

$$L^{-1}(a, b, c) = \sum \bar{\sigma}(a_1 b_1, c_1)\bar{\sigma}(a_2, b_2),$$
$$R^{-1}(a, b, c) = \sum \bar{\sigma}(a_1, b_1 c_1)\bar{\sigma}(b_2, c_2).$$

Indeed, each is obtained by reversing the two convolution factors defining L or R; ordinary multiplication in C is a coalgebra map. If $m_3(a, b, c) = abc$, substitution and cancellation of adjacent convolution inverses give

$$(a \star b) \star c = (L * m_3 * L^{-1})(a, b, c),$$
$$a \star (b \star c) = (R * m_3 * R^{-1})(a, b, c).$$

For instance the first expression expands as

$$\sum \sigma(a_1, b_1)\sigma(a_2 b_2, c_1) a_3 b_3 c_2 \bar{\sigma}(a_4 b_4, c_3)\bar{\sigma}(a_5, b_5).$$

Since $L = R$, associativity follows; normalization gives the unit. Substitution of pure tensors gives (13). It also gives the asserted factor products. The counits provide linear left inverses to the two factor inclusions. □

Theorem 5.3. *The identity assignments on P_r, N_r, E_i, F_i give an algebra isomorphism $\mathcal{A} \cong \mathcal{D}$.*

Proof. The cross product of $x \in B_+$ with $y \in B_-$ is

$$xy = \sum \tau(x_1, y_1)\tau(x_3, S(y_3)) y_2 x_2. \tag{15}$$

If $c_{ru} = 1/n - \delta_{ru}$, then its row and column sums vanish and

$$c_{ri} - c_{r,i+1} = -w_{ri},$$
$$c_{ij} - c_{i,j+1} - c_{i+1,j} + c_{i+1,j+1} = -a_{ij}.$$

Consequently $\tau(P_r, B_i) = \tau(A_i, N_r) = q^{-w_{ri}}$ and $\tau(A_i, B_j) = q^{-a_{ij}}$. Mixed zero values also hold for inverse Cartans, by applying (11) to a Cartan times its inverse. In (15), a pair of group-like Cartans has cancelling scalar factors. A Cartan and a root have only one nonzero term. These observations give the first three relations of (8).

For the remaining relation, the triples in the iterated coproducts are

$$(E_i, 1, 1),\ (A_i, E_i, 1),\ (A_i, A_i, E_i)$$

and

$$(F_j, B_j^{-1}, B_j^{-1}),\ (1, F_j, B_j^{-1}),\ (1, 1, F_j).$$

Writing $h_{ij} = -\delta_{ij}/d$, their nine contributions to (15) are

$$\begin{pmatrix} h_{ij}B_j^{-1} & 0 & 0 \\ 0 & F_jE_i & 0 \\ 0 & 0 & -h_{ij}A_i \end{pmatrix}.$$

The last entry uses $S(F_j) = -F_jB_j$ and $\tau(E_i, F_jB_j) = h_{ij}$. This proves the commutator and defines a homomorphism $\pi : \mathcal{A} \to \mathcal{D}$.

Multiplication defines a linear map $\mu : B_- \otimes B_+ \to \mathcal{A}$ using the canonical maps from the factor presentations. It is surjective: count root letters in a word, then count pairs in which a positive letter precedes a negative letter. Swapping an adjacent pair using (8) decreases the second count in the reordered term. The correction to an E_iF_j product replaces two root letters by Cartans, so decreases the first count. Lexicographic induction on these nonnegative counts puts every word in negative–positive order; inverse Cartans obey the same scalar exchange rules.

By Lemma 5.2, $\pi\mu$ is the canonical vector-space bijection onto $\mathcal{D}$. It follows that μ is injective. If $\pi(a) = 0$, choose v with $\mu(v) = a$; then $\pi\mu(v) = 0$ forces $v = 0$ and $a = 0$. Thus π is an isomorphism. No assertion of confluence is needed. □

6 The full diagonal quotient

Proposition 6.1. *In $\mathcal{A}$, the elements $Z_r = P_rN_r^{-1}$ are central and $\prod_r Z_r = 1$. If $J = (Z_r - 1 : 1 \le r \le n)$, then*

$$J = (P_r - N_r : 1 \le r \le n), \qquad \mathcal{A}/J \cong \mathcal{T}.$$

The algebra $\mathcal{T}$ is isomorphic to the full-weight algebra of [3, Definition 2.13] at (4).

Proof. Cartans commute, and the conjugation weights of P_r and N_r^{-1} cancel on both E_i and F_i. This proves centrality. The product-one relations give $\prod Z_r = 1$. The identities

$$(Z_r - 1)N_r = P_r - N_r, \qquad (P_r - N_r)N_r^{-1} = Z_r - 1$$

give equality of ideals. Send both Cartan families in $\mathcal{A}/J$ to K_r, fixing the roots. Every relation becomes a defining relation of $\mathcal{T}$, including (9). Conversely, send K_r to the class of P_r and fix the roots. The maps are inverse on generators.

For the last assertion send K_r to the source K_{λ_r} and fix E_i, F_i. The inverse sends K_λ to $\prod_r K_r^{m_r}$ for $\lambda = \sum m_r\lambda_r$. Lemma 4.1 proves independence of the representative. The source K_{α_i} becomes L_i, so its commutator, weights and Serre relations coincide with those of $\mathcal{T}$. This comparison concerns that source-defined algebra; an additional convention identification is needed for the target in (3). □

Proposition 6.2. *The ideal* $J_{\mathrm{root}} = (A_i - B_i : 1 \le i < n)$ *is strictly smaller than* J.

Proof. We have $A_i B_i^{-1} = Z_i Z_{i+1}^{-1}$, so $J_{\mathrm{root}} \subseteq J$. Modulo J_{root}, all Z_r coincide with a central element z satisfying $z^n = 1$; this does not force $z = 1$. More precisely, there is a homomorphism

$$\mathcal{A} \longrightarrow \mathbb{K}[\zeta]/(\zeta^n - 1),$$
$$P_r \longmapsto \zeta, \quad N_r \longmapsto 1, \quad E_i, F_i \longmapsto 0.$$

The element ζ is a unit, both determinant relations hold, and both A_i and B_i map to one, so every defining relation is respected. The map kills J_{root} but sends $Z_1 - 1$ to the nonzero class of $\zeta - 1$, whose degree is less than n. □

For example, at $n = 2$ write $P_1 = p = P_2^{-1}$ and $N_1 = m = N_2^{-1}$. Then

$$[E_1, F_1] = \frac{p^2 - m^{-2}}{d},$$

and the full quotient has $p = m = k$ with commutator $(k^2 - k^{-2})/d$. The weight Cartan k itself survives; its square is the simple-root Cartan. Although one of the two relations $Z_r = 1$ follows from the other, both belong to the specified list of diagonal relations.

7 Triangular matrices and explicit inverse maps

A matrix $T = (t_{ab})$ is a *q-quantum matrix* if every submatrix in increasing row and column order, $\left(\begin{smallmatrix} a & b \\ c & e \end{smallmatrix}\right)$, satisfies

$$ab = qba, \quad ac = qca, \quad be = qeb, \quad ce = qec, \quad bc = cb, \quad ae - ea = d\,bc. \tag{16}$$

> **Reviewer 1**
>
> This is different than the standard notation of $\left(\begin{smallmatrix} a & b \\ c & d \end{smallmatrix}\right)$ and is likely to lead to confusion.

Its quantum determinant is

$$\det_q T = \sum_{\sigma \in S_n} (-q)^{\ell(\sigma)} t_{1,\sigma(1)} \cdots t_{n,\sigma(n)},$$

where $\ell(\sigma)$ is the number of inversions of the permutation. Let Q_L be the algebra presented by a lower triangular q-quantum matrix of determinant one, and Q_U the corresponding upper triangular algebra. Upper or lower forbidden entries are set equal to zero, not removed from the relations before substitution.

In either triangular algebra the diagonal entries commute: a diagonal commutator in (16) contains a forbidden entry. Only the identity permutation survives in the determinant. Indeed, in the lower case a surviving term requires $\sigma(r) \le r$ for all r, and equality of the sums forces equality at every r; the upper case is the same with reversed inequalities. Thus if $p_r = t_{rr}$ then

$$\prod_r p_r = 1, \qquad p_r^{-1} = \prod_{u \neq r} p_u. \tag{17}$$

No localization is necessary to invert these entries.

For any root family X_i, define interval root vectors recursively by

$$\begin{aligned} R_{b+1,b}(X) &= X_b, \\ R_{ab}(X) &= R_{a-1,b}(X) X_{a-1} - q^{-1} X_{a-1} R_{a-1,b}(X) \quad (a > b+1). \end{aligned} \tag{18}$$

They are homogeneous of degree $a - b$ in the simple generators.

7.1 From entries to simple generators

Lemma 7.1. *There is a surjective homomorphism $\phi_+ : B_+ \to Q_L$ given by*

$$P_r \longmapsto p_r, \qquad E_i \longmapsto d^{-1}p_{i+1}^{-1}t_{i+1,i}. \tag{19}$$

For all $a > b$ it satisfies

$$\phi_+(R_{ab}(E)) = d^{-1}p_a^{-1}t_{ab}. \tag{20}$$

Proof. Set $u_i = t_{i+1,i}p_{i+1}^{-1}$. Same-row and same-column relations give $p_r t_{i+1,i} = q^{w_{ri}}t_{i+1,i}p_r$; other diagonal commutator corrections vanish by triangularity. Hence the proposed image of E_i is qu_i/d and has the required weights.

For adjacent indices, put

$$z = t_{i+1,i}, \quad w = t_{i+2,i+1}, \quad c = t_{i+2,i},$$
$$h = p_{i+1}, \quad k = p_{i+2}, \quad u = zh^{-1}, \quad v = wk^{-1}, \quad r = ck^{-1}.$$

The relevant relations are

$$zw - wz = dhc, \quad zc = qcz, \quad cw = qwc, \quad hc = ch, \quad zh = qhz,$$
$$hw = qwh, \quad kz = zk, \quad kw = q^{-1}wk, \quad kc = q^{-1}ck.$$

Moving the diagonal inverses to the right gives

$$quv - vu = dr, \qquad ur = qru, \qquad rv = qvr.$$

If $R = quv - vu$, the identities $uR = qRu$ and $Rv = qvR$ expand to the two ordered adjacent relations (6), after division by q. Distant u_i, u_j commute: their raw commutator correction contains a forbidden upper entry and the relevant diagonals commute with the other entry. Uniform scalar multiplication preserves these homogeneous relations. This proves that (19) is a homomorphism.

For $a > b + 1$, the matrix relations also give

$$t_{a-1,b}t_{a,a-1} - t_{a,a-1}t_{a-1,b} = d\,p_{a-1}t_{ab}. \tag{21}$$

Induction on $a - b$, using (17), proves generation by diagonals and adjacent entries, and hence surjectivity.

For the more precise formula, write $h = p_{a-1}$, $k = p_a$, $x = t_{a-1,b}$, $y = t_{a,a-1}$, $z = t_{ab}$. Besides (21), we have $kx = xk$ and $yh^{-1} = qh^{-1}y$. Induction in (18) gives

$$\phi_+(R_{ab}(E)) = d^{-2}h^{-1}k^{-1}(xy - yx) = d^{-1}k^{-1}z.$$

The initial case is (19). □

7.2 The inverse and its source input

The root identities needed to reverse this map can be stated without a basis assertion. Write $A = R_{ab}(E)$ and $B = R_{ce}(E)$, with $b < a$, $e < c$ and $(b, a) <_{\text{lex}} (e, c)$. They are

$$BA = \begin{cases} q^{-1}AB, & b = e < a < c \text{ or } b < e < a = c, \\ AB, & b < e < c < a \text{ or } b < a < e < c, \\ AB - dR_{cb}(E)R_{ae}(E), & b < e < a < c, \\ qAB - qR_{cb}(E), & b < a = e < c. \end{cases} \tag{22}$$

These are precisely the six configurations preceding [1, Theorem 3.1]. To check the normalization, set their parameter to $v = t^{n^2}$ and their rank to $n-1$. Then $v^2 = q$ and $v^8 \neq 1$, as required in their type-A discussion. Their generator x_{ba} is

$$x_{ba} = v^{a-b-1} R_{ab}(E),$$

by their recursive definition. In the crossing case the two product scalings coincide. In the touching case the source correction has one factor v and its long root has one extra factor v relative to the product; together these give the coefficient $v^2 = q$. This proves the precise specialization (22) of the cited identities in the Serre-presented algebra. It uses neither independence of root monomials nor a triangular-coordinate basis.

Proposition 7.2. *The map ϕ_+ is an isomorphism, with inverse*

$$t_{aa} \longmapsto P_a, \qquad t_{ab} \longmapsto dP_a R_{ab}(E) \quad (a > b). \tag{23}$$

There is also an isomorphism $\phi_- : B_- \to Q_U$. Writing y_{ab} for the entries of Q_U and $h_r = y_{rr}$, it is given by

$$N_r \longmapsto h_r^{-1}, \qquad F_i \longmapsto -d^{-1} y_{i,i+1} h_{i+1}^{-1}. \tag{24}$$

Its inverse sends, for $a > b$,

$$y_{aa} \longmapsto N_a^{-1}, \qquad y_{ba} \longmapsto (-1)^{a-b} d q^{a-b-1} R_{ab}(F) N_a^{-1}. \tag{25}$$

Proof. Let $T_{aa} = P_a$, $T_{ab} = dP_a R_{ab}(E)$ for $a > b$, and $T_{ab} = 0$ for $a < b$. Telescoping the simple weights gives

$$P_r R_{ab}(E) P_r^{-1} = q^{\delta_{rb} - \delta_{ra}} R_{ab}(E).$$

Thus the diagonal relations in (16) hold: common columns contribute q, common rows contribute q^{-1} when the diagonal is placed first, and disjoint diagonal commutators have a forbidden correction entry.

For completeness consider two distinct strict lower entries with intervals in the order used in (22). A shared start gives $T_{ab}T_{cb} = qT_{cb}T_{ab}$, since both products have common Cartan factor $d^2 P_a P_c$. A shared end gives $T_{ab}T_{ae} = qT_{ae}T_{ab}$, since both products have common factor $d^2 q P_a^2$. Nested intervals give northeast–southwest commutation. Disjoint intervals commute and their northwest–southeast correction is zero, because $T_{ae} = 0$.

For crossing intervals $b < e < a < c$, the forward and reverse products have common factor $d^2 P_a P_c$. Their difference is

$$d^3 P_a P_c R_{cb}(E) R_{ae}(E) = dT_{ae} T_{cb}.$$

Here the two correction roots are nested and commute, and their indicated Cartan reorderings have exponent zero. For touching intervals $b < a = e < c$, the two products are $d^2 P_a P_c AB$ and $d^2 q^{-1} P_a P_c BA$. Their difference is

$$d^2 P_a P_c R_{cb}(E) = dT_{aa} T_{cb}.$$

These are the remaining commutator relations. The six interval configurations exhaust distinct strict lower entries. If a northwest or southeast entry of a rectangle is forbidden, its northeast entry is also forbidden, so its commutator relation is zero on both sides. Relations with forbidden entries in a common row or column are also zero. Finally, the triangular determinant is $\prod P_a = 1$. Thus (23) defines a homomorphism $\theta_+ : Q_L \to B_+$.

Equation (20) shows that $\phi_+\theta_+$ fixes every matrix entry; the other composite fixes P_r, E_i immediately. This proves the first isomorphism.

Transposition $t_{ab} \mapsto y_{ba}$ preserves the six matrix relations and the triangular determinant. The presentation map

$$\chi : B_- \longrightarrow B_+, \qquad N_r \longmapsto P_r^{-1}, \quad F_i \longmapsto -q^{-1}E_i$$

is an isomorphism: its inverse sends P_r to N_r^{-1} and E_i to $-qF_i$, and the weights and homogeneous root relations are respected. Composing χ, ϕ_+ and transposition gives (24), using $h_{i+1}^{-1}y_{i,i+1} = qy_{i,i+1}h_{i+1}^{-1}$.

The homogeneity of R_{ab}, of degree $m = a - b$, now gives

$$\phi_-(R_{ab}(F)) = (-1)^m q^{1-m} d^{-1} y_{ba} h_a^{-1},$$

which proves (25). In particular its adjacent-entry formula is $y_{i,i+1} \mapsto -dF_iN_{i+1}^{-1}$; both the sign and the placement of the Cartan factor are fixed. □

8 A conditional comparison theorem

We now isolate precisely the matrix information needed from a skein algebra. The hypotheses concern an independently given algebra, and include linear independence through a factorization, not just a set of generator relations.

Theorem 8.1. *Let $\mathcal{H}$ be a unital $\mathbb{K}$-algebra with a lower triangular q-quantum matrix $X = (x_{ij})$ and an upper triangular q-quantum matrix $Y = (y_{kl})$, each of determinant one. Assume:*

1. *The entry assignments from Q_L, Q_U induce a vector-space bijection*

$$Q_L \otimes Q_U \longrightarrow \mathcal{H}, \qquad a \otimes b \longmapsto a(X)b(Y). \tag{26}$$

2. *For all $1 \le i, j, k, l \le n$ the mixed exchange identity is*

$$\begin{aligned} y_{kl}x_{ij} = {} & q^{\delta_{ik}-\delta_{jl}}x_{ij}y_{kl} + d\,\mathbf{1}_{\{k<i\}}q^{-\delta_{jl}}x_{kj}y_{il} \\ & - d\,\mathbf{1}_{\{j<l\}}q^{\delta_{ik}}x_{il}y_{kj} - d^2\mathbf{1}_{\{k<i\}}\mathbf{1}_{\{j<l\}}x_{kl}y_{ij}. \end{aligned} \tag{27}$$

   Here an indicator $\mathbf{1}_{\{C\}}$ equals one if C holds and zero otherwise, and forbidden triangular entries are zero.

Then there is an algebra isomorphism $\Phi : \mathcal{H} \to \mathcal{A} \cong \mathcal{D}$ with

$$\begin{aligned} &\Phi(x_{rr}) = P_r, \qquad \Phi(y_{rr}) = N_r^{-1}, \\ &\Phi(x_{ab}) = dP_aR_{ab}(E) \quad (a > b), \\ &\Phi(y_{ba}) = (-1)^{a-b}dq^{a-b-1}R_{ab}(F)N_a^{-1} \quad (a > b). \end{aligned} \tag{28}$$

If $I_{\mathcal{H}} = (x_{rr}y_{rr} - 1 : 1 \le r \le n)$, then $\Phi(I_{\mathcal{H}}) = J$ and $\mathcal{H}/I_{\mathcal{H}} \cong \mathcal{T}$.

Proof. Put $p_r = x_{rr}$, $h_r = y_{rr}$, $e_i = x_{i+1,i}$ and $f_i = y_{i,i+1}$. By (17), all p_r, h_r are units. The internal matrix weights are

$$p_re_i = q^{w_{ri}}e_ip_r, \qquad h_rf_i = q^{w_{ri}}f_ih_r.$$

Substitute a diagonal entry into (27). Every correction term contains a forbidden entry of X or Y, giving

$$p_rh_u = h_up_r, \qquad p_rf_i = q^{-w_{ri}}f_ip_r, \qquad h_re_j = q^{-w_{rj}}e_jh_r. \tag{29}$$

For example, with $x_{ij} = x_{rr}$, the three correction branches contain respectively x_{kr} with $k < r$, x_{rl} with $r < l$, and x_{kl} with $k < r < l$. The case of diagonal Y follows in the same way from its upper triangularity.

For adjacent entries (27) becomes

$$f_i e_j = q^{\delta_{i,j+1} - \delta_{i,j-1}} e_j f_i + \delta_{ij} d(p_i h_{i+1} - p_{i+1} h_i). \tag{30}$$

Indeed the two linear correction branches survive only for $i = j$; the quadratic branch requires $i = j$ and then contains $x_{i,i+1} = 0$.

Set $u_j = e_j p_{j+1}^{-1}$ and $v_i = f_i h_{i+1}^{-1}$. The internal weights and (29) yield

$$\begin{aligned} u_j v_i &= q^{\delta_{i,j+1} - \delta_{ij}} e_j f_i p_{j+1}^{-1} h_{i+1}^{-1}, \\ v_i u_j &= q^{\delta_{i,j-1} - \delta_{ij}} f_i e_j p_{j+1}^{-1} h_{i+1}^{-1}. \end{aligned}$$

Thus (30) implies $[u_j, v_i] = 0$ for $i \neq j$, whereas

$$[u_i, v_i] = -q^{-1} d(p_i p_{i+1}^{-1} - h_i h_{i+1}^{-1}).$$

It follows that

$$\begin{aligned} &\Psi(P_r) = p_r, \qquad \Psi(N_r) = h_r^{-1}, \\ &\Psi(E_i) = d^{-1} p_{i+1}^{-1} e_i = q u_i / d, \\ &\Psi(F_i) = -d^{-1} f_i h_{i+1}^{-1} = -v_i / d \end{aligned} \tag{31}$$

satisfy the mixed commutator (8). Equations (29) give the other mixed relations. Lemma 7.1 applied to X and to Y^{T} gives the internal weights and Serre relations. Hence $\Psi : \mathcal{A} \to \mathcal{H}$ is a homomorphism.

To prove bijectivity, multiplication $\nu : B_+ \otimes B_- \to \mathcal{A}$ is surjective by the word induction in Theorem 5.3, now exchanging adjacent negative–positive pairs. Solving (8) in that direction still decreases the inversion count or removes two root letters. The composite $\Psi\nu$ is the vector-space bijection (26) transported through the factor isomorphisms of Proposition 7.2. If $\nu(v) = 0$, this composite forces $v = 0$. If $\Psi(a) = 0$, choose v with $\nu(v) = a$; again $v = 0$. The composite also shows Ψ is surjective. Therefore Ψ is bijective, and its inverse is (28) by the explicit factor inverses. This proves the whole-algebra comparison without presuming that the formulas in the opposite direction respect unspecified additional relations.

Each diagonal generator of $I_{\mathcal{H}}$ maps to $P_r N_r^{-1} - 1 = Z_r - 1$. Thus $\Phi(I_{\mathcal{H}}) \subseteq J$. Conversely, in any finite sum $\sum a'_r (Z_r - 1) b'_r$ representing an element of J, lift both coefficients through the surjective map Φ. This gives a preimage in $I_{\mathcal{H}}$ and proves equality of the full two-sided ideals. Proposition 6.1 gives the quotient statement. □

9 Application to the original problem and remaining identifications

Theorem 8.1 reduces the original question to identifying its hypotheses with the actual skein algebra and its designated arcs. The distinction between a factorization before and after boundary reduction matters. The following elementary lemma supplies that bridge when the required ideal normality is known.

Lemma 9.1. *Let H_1, H_2 be unital subalgebras of an algebra H over a field, such that multiplication $H_1 \otimes H_2 \to H$ is a vector-space bijection. Suppose L_i is a two-sided ideal of H_i and $HL_i = L_iH$ for $i = 1, 2$. If K is the ideal they generate in H, multiplication induces a vector-space bijection*

$$(H_1/L_1) \otimes (H_2/L_2) \longrightarrow H/K.$$

Proof. Normality makes L_1H and HL_2 two-sided ideals, so $K = L_1H + HL_2$. Since H is spanned by H_1H_2, the internal ideal properties imply

$$K = L_1H_2 + H_1L_2.$$

Under the multiplication bijection this is $L_1 \otimes H_2 + H_1 \otimes L_2$. Extending bases of the L_i to bases of the H_i identifies that subspace with the kernel of the tensor quotient map. Taking quotients proves the assertion. The factors need not commute; this is not an isomorphism with an algebra tensor product. □

In [2], bad corner arcs have states $i < j$ in the local corner convention; their generated ideal defines the reduced algebra. A punctured surface is essentially bordered when every component has boundary. A disjoint system of boundary-to-boundary arcs is saturated when every complementary component contains exactly one removed point and the system is maximal with this property. For such systems, [2, Theorem 6.2] gives a multiplication factorization by arc-neighborhood algebras. Theorem 7.2, equations (128)–(129), gives normality of the corner ideals and expresses the total bad-arc ideal as their sum. These are possible inputs to Lemma 9.1; their application requires distinguishing reduction at one selected corner of a neighborhood bigon from reduction at both corners, and identifying the corresponding triangular quotients.

For the original problem, a sufficient missing geometric statement is the following: the designated c_{ij} are the entries x_{ij} of a lower triangular matrix, the designated b_{ij} are the entries y_{ij} of an upper triangular matrix, and these entries satisfy (26) and the exact identity (27), with the parameter convention (4). Rightward orientation alone does not determine these claims. Endpoint state order, framing, stacking, the chosen boundary reduction and the two crossing signs must all be checked. We do not supply that identification here.

If it were established, Theorem 8.1 would give the prospective original formulas

$$\begin{aligned} &c_{rr} \mapsto P_r, && b_{rr} \mapsto N_r^{-1}, \\ &c_{ab} \mapsto dP_aR_{ab}(E) && (a > b), \\ &b_{ba} \mapsto (-1)^{a-b}dq^{a-b-1}R_{ab}(F)N_a^{-1} && (a > b), \end{aligned}$$

with the forbidden entries zero. Since p_r and h_r commute by (29), the products $b_{rr}c_{rr}$ and $x_{rr}y_{rr}$ would coincide under this dictionary. The whole-ideal argument would then apply to every original diagonal generator. These are conditional formulas for the designated arcs, not an asserted map from the original skein algebra.

A second identification is needed on the quantum-group side. Proposition 5.1 uses the full-weight source presentation, passes to coopposite Borels, and constructs the ordered multiplication (13). Its generator pairing agrees numerically with (2). Identifying it with the convention intended in (1) still requires a dictionary for the original extended Cartans, coproduct, antipode, both skew rules and the order of the double. The comparison of $\mathcal{T}$ with the source presentation does not by itself provide that dictionary.

Finally, the base in (4) is an explicit generic choice, not a recovered hypothesis of the original problem. The matrix inverse above invokes the source condition $v^8 \neq 1$, with $v^2 = q$. No conclusion

for $q^4 = 1$, a divided-power form, or a restricted quantum group follows from that argument. A specialization to an algebraic value of t cannot be obtained merely by evaluating a $\mathbb{Q}(t)$-isomorphism: a nonzero polynomial vanishing at that value is invertible in $\mathbb{Q}(t)$. A result over a suitable localized coefficient ring, together with its geometric identifications, is needed for such a deduction.

Thus the pairing normalization, auxiliary inverse maps, injectivity mechanism and full diagonal quotient are available under the stated hypotheses. Establishing their correspondence with the original skein algebra, arcs, coefficients and quantum-group conventions remains necessary for both (1) and (3).

References

[1] Yuqun Chen, Hongshan Shao and K. P. Shum, *Rosso–Yamane Theorem on PBW basis of* $U_q(A_N)$, arXiv:0804.0954v1 (2008).


[2] Thang T. Q. Lê and Tao Yu, *Quantum traces for* SL_n*-skein algebras*, arXiv:2303.08082v2 (2025).

[3] Christian Voigt and Robert Yuncken, *Complex semisimple quantum groups and representation theory*, arXiv:1705.05661v4 (2020).

Review of Problem 3 – MerLean

First Proof Third Batch
Reviewer 3

October 8–9, 2026



1 Review

The strategy that the other solutions (both human and machine) is to first give a presentation of $U_q(\mathfrak{b}_-) \bowtie U_q(\mathfrak{b}_+)$, then a presentation of $\overline{\mathscr{S}}_n(\mathfrak{B}^*)$, define an isomorphism between the two (and prove that it works), and finally describe what happens in the quotient. This paper only achieves the first step, which is probably the most basic one. It acknowledges that it only achieves a partial result, and that the result it obtains is not especially novel. What it achieves appears to be correct.

The exposition is somewhat clear, even though it is full of the usual AI trappings (such as "not just this, but that"). There are also very non-human choices, such as naming the entries of a matrix a, b, c and e. I was also confused, as I read the proof, what had been achieved at each step.

The bibliography is clearly insufficient. It only contains 3 papers; the most glaring omissions are the paper actually defining the stated skein algebras and the paper establishing the case $n = 2$. I would also expect a paper to cite some standard references for the definition of the quantum groups involved.

Correctness.

- 1: essentially flawless mathematical argument
- 0.5: minor errors that are easily corrected
- 0: blank solution, or acknowledged partial result
- -0.5: major errors
- -1: proposed proof makes no progress towards the result.

Note that points should be docked for wrong statements/ solutions.
Score: 0

Exposition.

- 1: well-written and clear solution.
- 0: argument that is possible to follow with serious effort
- -1: very poor exposition

Score: 0

Attributions.

- 1: solutions contains fully appropriate attributions to related work and theorems being used.
- 0: moderate issues with missing attributions or hallucinated references
- -1: blatant plagiarism or hallucinations

Score: -1

2 Submission with inline comments

Abstract

The problem considered here asks for an explicit identification of a reduced stated skein algebra of a punctured bigon with a generalized double of the full-weight quantum Borel algebras, followed by a diagonal quotient giving the simply-connected quantum group. We give a partial answer. At a specified generic parameter we compute the ordered double, its full diagonal quotient, and explicit inverse formulas for triangular quantum matrices. These yield a comparison theorem for an algebra satisfying a precise mixed exchange identity and a vector-space factorization. The proof retains the weight lattice and distinguishes gluing all weight Cartans from gluing only simple-root Cartans. The identification of these hypotheses with the designated skein algebra and its arc conventions remains unresolved; consequently neither requested isomorphism for the original objects is asserted.

3 The question and the scope of the reduction

Let $U_q(\mathfrak{b}_+)$ and $U_q(\mathfrak{b}_-)$ denote the upper and lower Borel subalgebras of the simply-connected quantum group $U_q(\mathfrak{sl}_n)$. The question is to construct

$$\overline{\mathscr{S}}_n(\mathfrak{B}^*) \;\cong\; U_q(\mathfrak{b}_-) \bowtie U_q(\mathfrak{b}_+) \tag{1}$$

for the prescribed Drinfeld–Tanisaki pairing

$$\begin{gathered}\langle \hat{K}_r, \hat{K}_u \rangle = q^{1/n-\delta_{ru}}, \qquad \langle E_i, F_j \rangle = \frac{\delta_{ij}}{q^{-1}-q}, \\ \langle \hat{K}_r, F_j \rangle = \langle E_i, \hat{K}_u \rangle = 0,\end{gathered} \tag{2}$$

and then to deduce

$$U_q(\mathfrak{sl}_n) \;\cong\; \overline{\mathscr{S}}_n(\mathfrak{B}^*)/(b_{rr}c_{rr} - 1 : 1 \le r \le n). \tag{3}$$

Here b_{ij} and c_{ij} are the two designated families of rightward horizontal stated arcs. A state is an endpoint label in $\{1, \dots, n\}$. Parentheses in an algebra quotient always denote the two-sided ideal generated by the listed elements.

An explicit solution requires more than a matching commutator. The endpoint order, boundary reduction and multiplication convention on the skein side must determine the same mixed relation as the ordered double. Injectivity must then be proved. Furthermore, the simply-connected form involves weight Cartans, not just simple-root Cartans.

The precise definition of the original $\overline{\mathscr{S}}_n(\mathfrak{B}^*)$, its designated arc diagram, and its coefficient and fractional-power conventions are not fixed here. We therefore keep (1)–(3) as the original targets and distinguish them from all algebras defined below. The results of this paper are a generic algebraic reduction of these targets. The missing geometric and convention identifications are stated in Section 9.

Our main comparison result is Theorem 8.1. It gives every triangular matrix entry as a root vector with its exact scalar and Cartan factor. Its hypotheses include an explicit mixed exchange identity and a multiplication factorization; it does not presume the desired isomorphism. The independent presentation calculation in Theorem 5.3 identifies the resulting algebra with a specified ordered double. Proposition 6.1 proves the diagonal quotient, and Proposition 6.2 explains why gluing only simple-root ratios is insufficient.

The full-weight presentations and pairing used here are recalled from Voigt–Yuncken [3, Definition 2.13, Proposition 2.14, and Section 2.8.1]; we explicitly convert their coproduct convention.

The inverse triangular-matrix calculation uses the type-A root relations recorded by Chen–Shao–Shum [1, Section 3], who attribute those relations to Yamane.

> **Reviewer 3**
>
> It is a bit bizarre to cite a paper for something and then acknowledge that the paper credits that to someone else, rather than directly crediting that person

We use the relations, rather than their basis theorem. On the geometric side, Lê–Yu [2, Theorems 6.2 and 7.2] provide arc factorization and normality of the ideals generated by bad corner arcs. Their Theorem 6.2 is credited to Lê–Sikora. Section 9 explains how such results could supply a factorization after reduction, without identifying the original diagram by assumption. No novelty claim is made for the underlying quantum-group theory.

4 Weight Cartans and the two presentations

Throughout the algebraic results, fix an integer $n \geq 2$, an indeterminate t, and

$$\mathbb{K} = \mathbb{Q}(t), \qquad q = t^{2n^2}, \qquad d = q - q^{-1}, \qquad q^{1/n} = t^{2n}. \tag{4}$$

Thus q is not a root of unity and $d \neq 0$. All tensor products are over $\mathbb{K}$ unless another base is specified. We use $1 \leq r, u \leq n$ for weight indices and $1 \leq i, j < n$ for simple-root indices. Set

$$w_{ri} = \delta_{ri} - \delta_{r,i+1}, \qquad a_{ij} = 2\delta_{ij} - \delta_{i,j+1} - \delta_{i+1,j}.$$

4.1 Why the determinant-one torus is the weight torus

Let $\varepsilon_1, \dots, \varepsilon_n$ be the standard basis of $\mathbb{Q}^n$ and put

$$\lambda_r = \varepsilon_r - \frac{1}{n}\sum_{u=1}^{n} \varepsilon_u, \qquad W = \{x \in \mathbb{Q}^n : \textstyle\sum_r x_r = 0,\ x_r - x_u \in \mathbb{Z} \text{ for all } r, u\}.$$

The lattice W, with the ordinary dot product, is the type-A_{n-1} weight lattice. Its simple roots are $\alpha_i = \lambda_i - \lambda_{i+1}$.

Lemma 4.1. *The map $[m] \mapsto \sum_r m_r \lambda_r$ identifies $\mathbb{Z}^n/\mathbb{Z}(1, \dots, 1)$ with W. The fundamental weights are $\omega_i = \lambda_1 + \cdots + \lambda_i$. Moreover,*

$$(\lambda_r, \lambda_u) = \delta_{ru} - 1/n, \qquad (\lambda_r, \alpha_i) = w_{ri}.$$

The least positive integer L for which $(W, W) \subseteq L^{-1}\mathbb{Z}$ is n.

Proof. The image of m has coordinates $m_r - \frac{1}{n}\sum_u m_u$. It is zero precisely when m is constant. Conversely, for $x \in W$ take the integers $m_r = x_r - x_n$; their image is x. The displayed inner products follow by expansion. They give $(\omega_i, \alpha_j) = \delta_{ij}$, while $\lambda_1 = \omega_1$, $\lambda_r = \omega_r - \omega_{r-1}$ for $1 < r < n$, and $\lambda_n = -\omega_{n-1}$. Thus these are all the fundamental weights and they generate W. For integer representatives m, l,

$$\left(\sum_r m_r \lambda_r, \sum_r l_r \lambda_r\right) = \sum_r m_r l_r - \frac{(\sum_r m_r)(\sum_r l_r)}{n}.$$

This belongs to $n^{-1}\mathbb{Z}$, and $(\lambda_1, \lambda_1) = (n-1)/n$ has denominator n. □

4.2 Borel and double presentations

Define B_+ to be the algebra generated by commuting units P_r and elements E_i, subject to

$$\prod_r P_r = 1, \qquad P_r E_i P_r^{-1} = q^{w_{ri}} E_i. \tag{5}$$

The root generators satisfy the type-A quantum Serre relations: for $X_i = E_i$,

$$\begin{aligned} X_i X_j &= X_j X_i \quad (|i-j|>1), \\ X_i^2 X_j - (q+q^{-1}) X_i X_j X_i + X_j X_i^2 &= 0 \qquad (|i-j|=1). \end{aligned} \tag{6}$$

Define B_- by commuting units N_r with $\prod_r N_r = 1$, generators F_i, the same root relations, and

$$N_r F_i N_r^{-1} = q^{-w_{ri}} F_i. \tag{7}$$

For $\lambda = \sum m_r \lambda_r$ write $P_\lambda = \prod P_r^{m_r}$ and $N_\lambda = \prod N_r^{m_r}$. Lemma 4.1 and the product-one relations make these well defined. Abbreviate

$$A_i = P_i P_{i+1}^{-1}, \qquad B_i = N_i N_{i+1}^{-1}.$$

Here the indexed element B_i is distinct from the algebra B_-.

Define $\mathcal{A}$ by the two Borel presentations together with

$$\begin{aligned} P_r N_u &= N_u P_r, & P_r F_i &= q^{-w_{ri}} F_i P_r, \\ N_r E_i &= q^{w_{ri}} E_i N_r, & [E_i, F_j] &= \delta_{ij} \frac{A_i - B_i^{-1}}{d}. \end{aligned} \tag{8}$$

As usual, $[x,y] = xy - yx$. Define $\mathcal{T}$ to have one family of commuting units K_r of product one, the roots E_i, F_i, their relations (6), weights $q^{w_{ri}}$ and $q^{-w_{ri}}$, and

$$[E_i, F_j] = \delta_{ij} \frac{L_i - L_i^{-1}}{d}, \qquad L_i = K_i K_{i+1}^{-1}. \tag{9}$$

These are definitions of auxiliary algebras, independent of any skein identification.

5 The pairing and the ordered double

We first realize $B_\pm$ as actual Hopf algebras. A Hopf algebra has a coproduct Δ, a counit ϵ, and an antipode S. We write $\Delta(x) = \sum x_1 \otimes x_2$ and $(\Delta \otimes \mathrm{id})\Delta(x) = \sum x_1 \otimes x_2 \otimes x_3$, suppressing summation indices. All such sums below are finite for a fixed element. A group-like element g satisfies $\Delta(g) = g \otimes g$ and $\epsilon(g) = 1$; its antipode is g^{-1}.

Proposition 5.1. *The presentations $B_\pm$ admit Hopf structures with group-like Cartans and*

$$\begin{aligned} \Delta(E_i) &= E_i \otimes 1 + A_i \otimes E_i, & S(E_i) &= -A_i^{-1} E_i, \\ \Delta(F_i) &= F_i \otimes B_i^{-1} + 1 \otimes F_i, & S(F_i) &= -F_i B_i. \end{aligned} \tag{10}$$

The Cartan counits are one and the root counits are zero. There is a bilinear pairing $\tau : B_+ \times B_- \to \mathbb{K}$ satisfying

$$\begin{aligned} \tau(xx', y) &= \sum \tau(x', y_1)\tau(x, y_2), \\ \tau(x, yy') &= \sum \tau(x_1, y)\tau(x_2, y'), \\ \tau(1, y) &= \epsilon(y), \qquad \tau(x, 1) = \epsilon(x), \end{aligned} \tag{11}$$

and

$$\begin{aligned} \tau(P_r, N_u) = q^{1/n-\delta_{ru}}, \qquad \tau(E_i, F_j) = -\delta_{ij}/d, \\ \tau(P_r, F_j) = \tau(E_i, N_u) = 0. \end{aligned} \tag{12}$$

Proof. Use the full-weight algebra of [3, Definition 2.13] for type A_{n-1}. The source rank is $n-1$, its squared root lengths are two, and its integer L is n by Lemma 4.1. Its parameter s can be taken to be t^{2n}, so its $q = s^L$ agrees with (4). This checks both the fractional-power convention and the non-root-of-unity hypothesis used for its pairing.

The paragraph after [3, Proposition 2.14] identifies the actual Borel subalgebras with their corresponding presentations. Send their K_λ to P_λ or N_λ, respectively, and fix the root generators. The inverse assignments are $P_r \mapsto K_{\lambda_r}$ and $N_r \mapsto K_{\lambda_r}$. Lemma 4.1 checks the torus relations; the weights are $(\lambda_r, \alpha_i) = w_{ri}$, and the type-A Serre relations are exactly (6). These are inverse presentation maps on the actual Borels.

Write Δ_0, S_0 for the transported source Hopf operations. Their root formulas are

$$\begin{aligned} \Delta_0(E_i) = E_i \otimes A_i + 1 \otimes E_i, \qquad \Delta_0(F_i) = F_i \otimes 1 + B_i^{-1} \otimes F_i, \\ S_0(E_i) = -E_i A_i^{-1}, \qquad S_0(F_i) = -B_i F_i. \end{aligned}$$

The square of S_0 scales E_i by q^2 and F_i by q^{-2} and fixes Cartans. These scalings are automorphisms, so S_0 is invertible. Replacing Δ_0 by its opposite coproduct and S_0 by S_0^{-1} gives (10). This operation is called taking the coopposite Hopf algebra; multiplication is unchanged.

By [3, Proposition 2.73], the source Borels possess a pairing $\rho : B_- \times B_+ \to \mathbb{K}$ with

$$\rho(N_\lambda, P_\mu) = q^{(\lambda,\mu)}, \qquad \rho(F_j, E_i) = \delta_{ij}/d$$

and both mixed root–torus values zero. Its two multiplication rules, using Δ_0, are

$$\begin{aligned} \rho(yy', x) = \sum \rho(y, x_{1,0})\rho(y', x_{2,0}), \\ \rho(y, xx') = \sum \rho(y_{1,0}, x')\rho(y_{2,0}, x). \end{aligned}$$

Define $\tau(x, y) = \rho(S_0(y), x)$, as in [3, Definition 2.74]. Since S_0 reverses multiplication and coproduct, these rules give

$$\begin{aligned} \tau(xx', y) = \sum \tau(x, y_{1,0})\tau(x', y_{2,0}), \\ \tau(x, yy') = \sum \tau(x_{1,0}, y')\tau(x_{2,0}, y). \end{aligned}$$

Reversing the coproduct indices gives (11).

For Cartans this construction gives $\tau(P_\lambda, N_\mu) = q^{-(\lambda,\mu)}$. The mixed zero values follow from the corresponding values of ρ. Finally,

$$\begin{aligned} \tau(E_i, F_j) &= -\rho(B_j F_j, E_i) \\ &= -\rho(B_j, E_i)\rho(F_j, A_i) - \rho(B_j, 1)\rho(F_j, E_i) = -\delta_{ij}/d. \end{aligned}$$

Thus existence precedes the calculation of the generator table; it is not inferred from that table alone. □

5.1 An associative product on the ordered tensor space

The chosen order of the double is negative followed by positive. We include the construction to specify its multiplication and to justify the tensor-space comparison used later.

Lemma 5.2. *The vector space $B_- \otimes B_+$ is an associative unital algebra $\mathcal{D}$ under*

$$(y \otimes x)(z \otimes u) = \sum \tau(x_1, z_1)\tau(x_3, S(z_3))\, yz_2 \otimes x_2 u. \tag{13}$$

Both factors embed as algebras and $(y \otimes 1)(1 \otimes x) = y \otimes x$.

Proof. Start with the ordinary tensor-product Hopf algebra $C = B_- \otimes B_+$. Define scalar-valued maps

$$\sigma(y \otimes x, z \otimes u) = \epsilon(y)\tau(x, z)\epsilon(u),$$
$$\bar{\sigma}(y \otimes x, z \otimes u) = \epsilon(y)\tau(x, S(z))\epsilon(u).$$

For maps on a coalgebra, convolution is $(f * g)(a) = \sum f(a_1)g(a_2)$. The second rule in (11) and the two antipode identities show that $\sigma * \bar{\sigma} = \bar{\sigma} * \sigma = \epsilon \otimes \epsilon$. The unit pairing gives $\sigma(a, 1) = \sigma(1, a) = \epsilon(a)$.

The required cocycle identity is

$$\sum \sigma(a_1, b_1)\sigma(a_2 b_2, c) = \sum \sigma(b_1, c_1)\sigma(a, b_2 c_2). \tag{14}$$

For $a = y \otimes x$, $b = z \otimes u$, $c = w \otimes v$, both sides reduce by (11) to

$$\epsilon(y)\epsilon(v) \sum \tau(x_1, z)\tau(u, w_1)\tau(x_2, w_2).$$

Pure tensors span C, so the identity holds generally.

Set $a \star b = \sum \sigma(a_1, b_1) a_2 b_2 \bar{\sigma}(a_3, b_3)$. To verify associativity explicitly, denote the left and right sides of (14) by $L(a, b, c)$ and $R(a, b, c)$. Their convolution inverses on $C^{\otimes 3}$ are

$$L^{-1}(a, b, c) = \sum \bar{\sigma}(a_1 b_1, c_1)\bar{\sigma}(a_2, b_2),$$
$$R^{-1}(a, b, c) = \sum \bar{\sigma}(a_1, b_1 c_1)\bar{\sigma}(b_2, c_2).$$

Indeed, each is obtained by reversing the two convolution factors defining L or R; ordinary multiplication in C is a coalgebra map. If $m_3(a, b, c) = abc$, substitution and cancellation of adjacent convolution inverses give

$$(a \star b) \star c = (L * m_3 * L^{-1})(a, b, c),$$
$$a \star (b \star c) = (R * m_3 * R^{-1})(a, b, c).$$

For instance the first expression expands as

$$\sum \sigma(a_1, b_1)\sigma(a_2 b_2, c_1) a_3 b_3 c_2 \bar{\sigma}(a_4 b_4, c_3)\bar{\sigma}(a_5, b_5).$$

Since $L = R$, associativity follows; normalization gives the unit. Substitution of pure tensors gives (13). It also gives the asserted factor products. The counits provide linear left inverses to the two factor inclusions. □

Theorem 5.3. *The identity assignments on P_r, N_r, E_i, F_i give an algebra isomorphism $\mathcal{A} \cong \mathcal{D}$.*

Proof. The cross product of $x \in B_+$ with $y \in B_-$ is

$$xy = \sum \tau(x_1, y_1)\tau(x_3, S(y_3)) y_2 x_2. \tag{15}$$

If $c_{ru} = 1/n - \delta_{ru}$, then its row and column sums vanish and

$$c_{ri} - c_{r,i+1} = -w_{ri},$$
$$c_{ij} - c_{i,j+1} - c_{i+1,j} + c_{i+1,j+1} = -a_{ij}.$$

Consequently $\tau(P_r, B_i) = \tau(A_i, N_r) = q^{-w_{ri}}$ and $\tau(A_i, B_j) = q^{-a_{ij}}$. Mixed zero values also hold for inverse Cartans, by applying (11) to a Cartan times its inverse. In (15), a pair of group-like Cartans has cancelling scalar factors. A Cartan and a root have only one nonzero term. These observations give the first three relations of (8).

For the remaining relation, the triples in the iterated coproducts are

$$(E_i, 1, 1),\ (A_i, E_i, 1),\ (A_i, A_i, E_i)$$

and

$$(F_j, B_j^{-1}, B_j^{-1}),\ (1, F_j, B_j^{-1}),\ (1, 1, F_j).$$

Writing $h_{ij} = -\delta_{ij}/d$, their nine contributions to (15) are

$$\begin{pmatrix} h_{ij}B_j^{-1} & 0 & 0 \\ 0 & F_jE_i & 0 \\ 0 & 0 & -h_{ij}A_i \end{pmatrix}.$$

The last entry uses $S(F_j) = -F_jB_j$ and $\tau(E_i, F_jB_j) = h_{ij}$. This proves the commutator and defines a homomorphism $\pi : \mathcal{A} \to \mathcal{D}$.

Multiplication defines a linear map $\mu : B_- \otimes B_+ \to \mathcal{A}$ using the canonical maps from the factor presentations. It is surjective: count root letters in a word, then count pairs in which a positive letter precedes a negative letter. Swapping an adjacent pair using (8) decreases the second count in the reordered term. The correction to an E_iF_j product replaces two root letters by Cartans, so decreases the first count. Lexicographic induction on these nonnegative counts puts every word in negative–positive order; inverse Cartans obey the same scalar exchange rules.

By Lemma 5.2, $\pi\mu$ is the canonical vector-space bijection onto $\mathcal{D}$. It follows that μ is injective. If $\pi(a) = 0$, choose v with $\mu(v) = a$; then $\pi\mu(v) = 0$ forces $v = 0$ and $a = 0$. Thus π is an isomorphism. No assertion of confluence is needed. □

6 The full diagonal quotient

Proposition 6.1. *In $\mathcal{A}$, the elements $Z_r = P_rN_r^{-1}$ are central and $\prod_r Z_r = 1$. If $J = (Z_r - 1 : 1 \le r \le n)$, then*

$$J = (P_r - N_r : 1 \le r \le n), \qquad \mathcal{A}/J \cong \mathcal{T}.$$

The algebra $\mathcal{T}$ is isomorphic to the full-weight algebra of [3, Definition 2.13] at (4).

Proof. Cartans commute, and the conjugation weights of P_r and N_r^{-1} cancel on both E_i and F_i. This proves centrality. The product-one relations give $\prod Z_r = 1$. The identities

$$(Z_r - 1)N_r = P_r - N_r, \qquad (P_r - N_r)N_r^{-1} = Z_r - 1$$

give equality of ideals. Send both Cartan families in $\mathcal{A}/J$ to K_r, fixing the roots. Every relation becomes a defining relation of $\mathcal{T}$, including (9). Conversely, send K_r to the class of P_r and fix the roots. The maps are inverse on generators.

For the last assertion send K_r to the source K_{λ_r} and fix E_i, F_i. The inverse sends K_λ to $\prod_r K_r^{m_r}$ for $\lambda = \sum m_r\lambda_r$. Lemma 4.1 proves independence of the representative. The source K_{α_i} becomes L_i, so its commutator, weights and Serre relations coincide with those of $\mathcal{T}$. This comparison concerns that source-defined algebra; an additional convention identification is needed for the target in (3). □

Proposition 6.2. *The ideal* $J_{\mathrm{root}} = (A_i - B_i : 1 \le i < n)$ *is strictly smaller than* J.

Proof. We have $A_i B_i^{-1} = Z_i Z_{i+1}^{-1}$, so $J_{\mathrm{root}} \subseteq J$. Modulo J_{root}, all Z_r coincide with a central element z satisfying $z^n = 1$; this does not force $z = 1$. More precisely, there is a homomorphism

$$\begin{aligned} \mathcal{A} &\longrightarrow \mathbb{K}[\zeta]/(\zeta^n - 1), \\ P_r \longmapsto \zeta, \quad N_r &\longmapsto 1, \quad E_i, F_i \longmapsto 0. \end{aligned}$$

The element ζ is a unit, both determinant relations hold, and both A_i and B_i map to one, so every defining relation is respected. The map kills J_{root} but sends $Z_1 - 1$ to the nonzero class of $\zeta - 1$, whose degree is less than n. □

For example, at $n = 2$ write $P_1 = p = P_2^{-1}$ and $N_1 = m = N_2^{-1}$. Then

$$[E_1, F_1] = \frac{p^2 - m^{-2}}{d},$$

and the full quotient has $p = m = k$ with commutator $(k^2 - k^{-2})/d$. The weight Cartan k itself survives; its square is the simple-root Cartan. Although one of the two relations $Z_r = 1$ follows from the other, both belong to the specified list of diagonal relations.

7 Triangular matrices and explicit inverse maps

A matrix $T = (t_{ab})$ is a *q-quantum matrix* if every submatrix in increasing row and column order, $\left(\begin{smallmatrix} a & b \\ c & e \end{smallmatrix}\right)$, satisfies

$$ab = qba, \quad ac = qca, \quad be = qeb, \quad ce = qec, \quad bc = cb, \quad ae - ea = d\,bc. \tag{16}$$

Its quantum determinant is

$$\det_q T = \sum_{\sigma \in S_n} (-q)^{\ell(\sigma)} t_{1,\sigma(1)} \cdots t_{n,\sigma(n)},$$

where $\ell(\sigma)$ is the number of inversions of the permutation. Let Q_L be the algebra presented by a lower triangular q-quantum matrix of determinant one, and Q_U the corresponding upper triangular algebra. Upper or lower forbidden entries are set equal to zero, not removed from the relations before substitution.

In either triangular algebra the diagonal entries commute: a diagonal commutator in (16) contains a forbidden entry. Only the identity permutation survives in the determinant. Indeed, in the lower case a surviving term requires $\sigma(r) \le r$ for all r, and equality of the sums forces equality at every r; the upper case is the same with reversed inequalities. Thus if $p_r = t_{rr}$ then

$$\prod_r p_r = 1, \qquad p_r^{-1} = \prod_{u \ne r} p_u. \tag{17}$$

No localization is necessary to invert these entries.

For any root family X_i, define interval root vectors recursively by

$$\begin{aligned} R_{b+1,b}(X) &= X_b, \\ R_{ab}(X) &= R_{a-1,b}(X) X_{a-1} - q^{-1} X_{a-1} R_{a-1,b}(X) \quad (a > b+1). \end{aligned} \tag{18}$$

They are homogeneous of degree $a - b$ in the simple generators.

7.1 From entries to simple generators

Lemma 7.1. *There is a surjective homomorphism $\phi_+ : B_+ \to Q_L$ given by*

$$P_r \longmapsto p_r, \qquad E_i \longmapsto d^{-1}p_{i+1}^{-1}t_{i+1,i}. \tag{19}$$

For all $a > b$ it satisfies

$$\phi_+(R_{ab}(E)) = d^{-1}p_a^{-1}t_{ab}. \tag{20}$$

Proof. Set $u_i = t_{i+1,i}p_{i+1}^{-1}$. Same-row and same-column relations give $p_r t_{i+1,i} = q^{w_{ri}}t_{i+1,i}p_r$; other diagonal commutator corrections vanish by triangularity. Hence the proposed image of E_i is qu_i/d and has the required weights.

For adjacent indices, put

$$z = t_{i+1,i}, \quad w = t_{i+2,i+1}, \quad c = t_{i+2,i},$$
$$h = p_{i+1}, \quad k = p_{i+2}, \quad u = zh^{-1}, \quad v = wk^{-1}, \quad r = ck^{-1}.$$

The relevant relations are

$$zw - wz = dhc, \quad zc = qcz, \quad cw = qwc, \quad hc = ch, \quad zh = qhz,$$
$$hw = qwh, \quad kz = zk, \quad kw = q^{-1}wk, \quad kc = q^{-1}ck.$$

Moving the diagonal inverses to the right gives

$$quv - vu = dr, \qquad ur = qru, \qquad rv = qvr.$$

If $R = quv - vu$, the identities $uR = qRu$ and $Rv = qvR$ expand to the two ordered adjacent relations (6), after division by q. Distant u_i, u_j commute: their raw commutator correction contains a forbidden upper entry and the relevant diagonals commute with the other entry. Uniform scalar multiplication preserves these homogeneous relations. This proves that (19) is a homomorphism.

For $a > b + 1$, the matrix relations also give

$$t_{a-1,b}t_{a,a-1} - t_{a,a-1}t_{a-1,b} = d\,p_{a-1}t_{ab}. \tag{21}$$

Induction on $a - b$, using (17), proves generation by diagonals and adjacent entries, and hence surjectivity.

For the more precise formula, write $h = p_{a-1}$, $k = p_a$, $x = t_{a-1,b}$, $y = t_{a,a-1}$, $z = t_{ab}$. Besides (21), we have $kx = xk$ and $yh^{-1} = qh^{-1}y$. Induction in (18) gives

$$\phi_+(R_{ab}(E)) = d^{-2}h^{-1}k^{-1}(xy - yx) = d^{-1}k^{-1}z.$$

The initial case is (19). □

7.2 The inverse and its source input

The root identities needed to reverse this map can be stated without a basis assertion. Write $A = R_{ab}(E)$ and $B = R_{ce}(E)$, with $b < a$, $e < c$ and $(b, a) <_{\text{lex}} (e, c)$. They are

$$BA = \begin{cases} q^{-1}AB, & b = e < a < c \text{ or } b < e < a = c, \\ AB, & b < e < c < a \text{ or } b < a < e < c, \\ AB - dR_{cb}(E)R_{ae}(E), & b < e < a < c, \\ qAB - qR_{cb}(E), & b < a = e < c. \end{cases} \tag{22}$$

These are precisely the six configurations preceding [1, Theorem 3.1]. To check the normalization, set their parameter to $v = t^{n^2}$ and their rank to $n-1$. Then $v^2 = q$ and $v^8 \neq 1$, as required in their type-A discussion. Their generator x_{ba} is

$$x_{ba} = v^{a-b-1} R_{ab}(E),$$

by their recursive definition. In the crossing case the two product scalings coincide. In the touching case the source correction has one factor v and its long root has one extra factor v relative to the product; together these give the coefficient $v^2 = q$. This proves the precise specialization (22) of the cited identities in the Serre-presented algebra. It uses neither independence of root monomials nor a triangular-coordinate basis.

Proposition 7.2. *The map ϕ_+ is an isomorphism, with inverse*

$$t_{aa} \longmapsto P_a, \qquad t_{ab} \longmapsto dP_a R_{ab}(E) \quad (a > b). \tag{23}$$

There is also an isomorphism $\phi_- : B_- \to Q_U$. Writing y_{ab} for the entries of Q_U and $h_r = y_{rr}$, it is given by

$$N_r \longmapsto h_r^{-1}, \qquad F_i \longmapsto -d^{-1} y_{i,i+1} h_{i+1}^{-1}. \tag{24}$$

Its inverse sends, for $a > b$,

$$y_{aa} \longmapsto N_a^{-1}, \qquad y_{ba} \longmapsto (-1)^{a-b} d q^{a-b-1} R_{ab}(F) N_a^{-1}. \tag{25}$$

Proof. Let $T_{aa} = P_a$, $T_{ab} = dP_a R_{ab}(E)$ for $a > b$, and $T_{ab} = 0$ for $a < b$. Telescoping the simple weights gives

$$P_r R_{ab}(E) P_r^{-1} = q^{\delta_{rb} - \delta_{ra}} R_{ab}(E).$$

Thus the diagonal relations in (16) hold: common columns contribute q, common rows contribute q^{-1} when the diagonal is placed first, and disjoint diagonal commutators have a forbidden correction entry.

For completeness consider two distinct strict lower entries with intervals in the order used in (22). A shared start gives $T_{ab}T_{cb} = qT_{cb}T_{ab}$, since both products have common Cartan factor $d^2 P_a P_c$. A shared end gives $T_{ab}T_{ae} = qT_{ae}T_{ab}$, since both products have common factor $d^2 q P_a^2$. Nested intervals give northeast–southwest commutation. Disjoint intervals commute and their northwest–southeast correction is zero, because $T_{ae} = 0$.

For crossing intervals $b < e < a < c$, the forward and reverse products have common factor $d^2 P_a P_c$. Their difference is

$$d^3 P_a P_c R_{cb}(E) R_{ae}(E) = dT_{ae} T_{cb}.$$

Here the two correction roots are nested and commute, and their indicated Cartan reorderings have exponent zero. For touching intervals $b < a = e < c$, the two products are $d^2 P_a P_c AB$ and $d^2 q^{-1} P_a P_c BA$. Their difference is

$$d^2 P_a P_c R_{cb}(E) = dT_{aa} T_{cb}.$$

These are the remaining commutator relations. The six interval configurations exhaust distinct strict lower entries. If a northwest or southeast entry of a rectangle is forbidden, its northeast entry is also forbidden, so its commutator relation is zero on both sides. Relations with forbidden entries in a common row or column are also zero. Finally, the triangular determinant is $\prod P_a = 1$. Thus (23) defines a homomorphism $\theta_+ : Q_L \to B_+$.

Equation (20) shows that $\phi_+\theta_+$ fixes every matrix entry; the other composite fixes P_r, E_i immediately. This proves the first isomorphism.

Transposition $t_{ab} \mapsto y_{ba}$ preserves the six matrix relations and the triangular determinant. The presentation map

$$\chi : B_- \longrightarrow B_+, \qquad N_r \longmapsto P_r^{-1}, \quad F_i \longmapsto -q^{-1}E_i$$

is an isomorphism: its inverse sends P_r to N_r^{-1} and E_i to $-qF_i$, and the weights and homogeneous root relations are respected. Composing χ, ϕ_+ and transposition gives (24), using $h_{i+1}^{-1}y_{i,i+1} = qy_{i,i+1}h_{i+1}^{-1}$.

The homogeneity of R_{ab}, of degree $m = a - b$, now gives

$$\phi_-(R_{ab}(F)) = (-1)^m q^{1-m} d^{-1} y_{ba} h_a^{-1},$$

which proves (25). In particular its adjacent-entry formula is $y_{i,i+1} \mapsto -dF_iN_{i+1}^{-1}$; both the sign and the placement of the Cartan factor are fixed. □

8 A conditional comparison theorem

We now isolate precisely the matrix information needed from a skein algebra. The hypotheses concern an independently given algebra, and include linear independence through a factorization, not just a set of generator relations.

Theorem 8.1. *Let $\mathcal{H}$ be a unital $\mathbb{K}$-algebra with a lower triangular q-quantum matrix $X = (x_{ij})$ and an upper triangular q-quantum matrix $Y = (y_{kl})$, each of determinant one. Assume:*

1. *The entry assignments from Q_L, Q_U induce a vector-space bijection*

$$Q_L \otimes Q_U \longrightarrow \mathcal{H}, \qquad a \otimes b \longmapsto a(X)b(Y). \tag{26}$$

2. *For all $1 \le i, j, k, l \le n$ the mixed exchange identity is*

$$\begin{aligned} y_{kl}x_{ij} = {} & q^{\delta_{ik}-\delta_{jl}}x_{ij}y_{kl} + d\,\mathbf{1}_{\{k<i\}}q^{-\delta_{jl}}x_{kj}y_{il} \\ & - d\,\mathbf{1}_{\{j<l\}}q^{\delta_{ik}}x_{il}y_{kj} - d^2\mathbf{1}_{\{k<i\}}\mathbf{1}_{\{j<l\}}x_{kl}y_{ij}. \end{aligned} \tag{27}$$

   Here an indicator $\mathbf{1}_{\{C\}}$ equals one if C holds and zero otherwise, and forbidden triangular entries are zero.

Then there is an algebra isomorphism $\Phi : \mathcal{H} \to \mathcal{A} \cong \mathcal{D}$ with

$$\begin{aligned} &\Phi(x_{rr}) = P_r, \qquad \Phi(y_{rr}) = N_r^{-1}, \\ &\Phi(x_{ab}) = dP_aR_{ab}(E) \quad (a > b), \\ &\Phi(y_{ba}) = (-1)^{a-b}dq^{a-b-1}R_{ab}(F)N_a^{-1} \quad (a > b). \end{aligned} \tag{28}$$

If $I_{\mathcal{H}} = (x_{rr}y_{rr} - 1 : 1 \le r \le n)$, then $\Phi(I_{\mathcal{H}}) = J$ and $\mathcal{H}/I_{\mathcal{H}} \cong \mathcal{T}$.

Proof. Put $p_r = x_{rr}$, $h_r = y_{rr}$, $e_i = x_{i+1,i}$ and $f_i = y_{i,i+1}$. By (17), all p_r, h_r are units. The internal matrix weights are

$$p_re_i = q^{w_{ri}}e_ip_r, \qquad h_rf_i = q^{w_{ri}}f_ih_r.$$

Substitute a diagonal entry into (27). Every correction term contains a forbidden entry of X or Y, giving

$$p_rh_u = h_up_r, \qquad p_rf_i = q^{-w_{ri}}f_ip_r, \qquad h_re_j = q^{-w_{rj}}e_jh_r. \tag{29}$$

For example, with $x_{ij} = x_{rr}$, the three correction branches contain respectively x_{kr} with $k < r$, x_{rl} with $r < l$, and x_{kl} with $k < r < l$. The case of diagonal Y follows in the same way from its upper triangularity.

For adjacent entries (27) becomes

$$f_i e_j = q^{\delta_{i,j+1} - \delta_{i,j-1}} e_j f_i + \delta_{ij} d(p_i h_{i+1} - p_{i+1} h_i). \tag{30}$$

Indeed the two linear correction branches survive only for $i = j$; the quadratic branch requires $i = j$ and then contains $x_{i,i+1} = 0$.

Set $u_j = e_j p_{j+1}^{-1}$ and $v_i = f_i h_{i+1}^{-1}$. The internal weights and (29) yield

$$\begin{aligned} u_j v_i &= q^{\delta_{i,j+1} - \delta_{ij}} e_j f_i p_{j+1}^{-1} h_{i+1}^{-1}, \\ v_i u_j &= q^{\delta_{i,j-1} - \delta_{ij}} f_i e_j p_{j+1}^{-1} h_{i+1}^{-1}. \end{aligned}$$

Thus (30) implies $[u_j, v_i] = 0$ for $i \neq j$, whereas

$$[u_i, v_i] = -q^{-1} d(p_i p_{i+1}^{-1} - h_i h_{i+1}^{-1}).$$

It follows that

$$\begin{aligned} &\Psi(P_r) = p_r, \qquad \Psi(N_r) = h_r^{-1}, \\ &\Psi(E_i) = d^{-1} p_{i+1}^{-1} e_i = q u_i / d, \\ &\Psi(F_i) = -d^{-1} f_i h_{i+1}^{-1} = -v_i / d \end{aligned} \tag{31}$$

satisfy the mixed commutator (8). Equations (29) give the other mixed relations. Lemma 7.1 applied to X and to Y^{T} gives the internal weights and Serre relations. Hence $\Psi : \mathcal{A} \to \mathcal{H}$ is a homomorphism.

To prove bijectivity, multiplication $\nu : B_+ \otimes B_- \to \mathcal{A}$ is surjective by the word induction in Theorem 5.3, now exchanging adjacent negative–positive pairs. Solving (8) in that direction still decreases the inversion count or removes two root letters. The composite $\Psi\nu$ is the vector-space bijection (26) transported through the factor isomorphisms of Proposition 7.2. If $\nu(v) = 0$, this composite forces $v = 0$. If $\Psi(a) = 0$, choose v with $\nu(v) = a$; again $v = 0$. The composite also shows Ψ is surjective. Therefore Ψ is bijective, and its inverse is (28) by the explicit factor inverses. This proves the whole-algebra comparison without presuming that the formulas in the opposite direction respect unspecified additional relations.

Each diagonal generator of $I_{\mathcal{H}}$ maps to $P_r N_r^{-1} - 1 = Z_r - 1$. Thus $\Phi(I_{\mathcal{H}}) \subseteq J$. Conversely, in any finite sum $\sum a'_r (Z_r - 1) b'_r$ representing an element of J, lift both coefficients through the surjective map Φ. This gives a preimage in $I_{\mathcal{H}}$ and proves equality of the full two-sided ideals. Proposition 6.1 gives the quotient statement. □

9 Application to the original problem and remaining identifications

Theorem 8.1 reduces the original question to identifying its hypotheses with the actual skein algebra and its designated arcs. The distinction between a factorization before and after boundary reduction matters. The following elementary lemma supplies that bridge when the required ideal normality is known.

Lemma 9.1. *Let H_1, H_2 be unital subalgebras of an algebra H over a field, such that multiplication $H_1 \otimes H_2 \to H$ is a vector-space bijection. Suppose L_i is a two-sided ideal of H_i and $HL_i = L_iH$ for $i = 1, 2$. If K is the ideal they generate in H, multiplication induces a vector-space bijection*

$$(H_1/L_1) \otimes (H_2/L_2) \longrightarrow H/K.$$

Proof. Normality makes L_1H and HL_2 two-sided ideals, so $K = L_1H + HL_2$. Since H is spanned by H_1H_2, the internal ideal properties imply

$$K = L_1H_2 + H_1L_2.$$

Under the multiplication bijection this is $L_1 \otimes H_2 + H_1 \otimes L_2$. Extending bases of the L_i to bases of the H_i identifies that subspace with the kernel of the tensor quotient map. Taking quotients proves the assertion. The factors need not commute; this is not an isomorphism with an algebra tensor product. □

In [2], bad corner arcs have states $i < j$ in the local corner convention; their generated ideal defines the reduced algebra. A punctured surface is essentially bordered when every component has boundary. A disjoint system of boundary-to-boundary arcs is saturated when every complementary component contains exactly one removed point and the system is maximal with this property. For such systems, [2, Theorem 6.2] gives a multiplication factorization by arc-neighborhood algebras. Theorem 7.2, equations (128)–(129), gives normality of the corner ideals and expresses the total bad-arc ideal as their sum. These are possible inputs to Lemma 9.1; their application requires distinguishing reduction at one selected corner of a neighborhood bigon from reduction at both corners, and identifying the corresponding triangular quotients.

For the original problem, a sufficient missing geometric statement is the following: the designated c_{ij} are the entries x_{ij} of a lower triangular matrix, the designated b_{ij} are the entries y_{ij} of an upper triangular matrix, and these entries satisfy (26) and the exact identity (27), with the parameter convention (4). Rightward orientation alone does not determine these claims. Endpoint state order, framing, stacking, the chosen boundary reduction and the two crossing signs must all be checked. We do not supply that identification here.

If it were established, Theorem 8.1 would give the prospective original formulas

$$\begin{aligned} &c_{rr} \mapsto P_r, && b_{rr} \mapsto N_r^{-1}, \\ &c_{ab} \mapsto dP_aR_{ab}(E) && (a > b), \\ &b_{ba} \mapsto (-1)^{a-b}dq^{a-b-1}R_{ab}(F)N_a^{-1} && (a > b), \end{aligned}$$

with the forbidden entries zero. Since p_r and h_r commute by (29), the products $b_{rr}c_{rr}$ and $x_{rr}y_{rr}$ would coincide under this dictionary. The whole-ideal argument would then apply to every original diagonal generator. These are conditional formulas for the designated arcs, not an asserted map from the original skein algebra.

A second identification is needed on the quantum-group side. Proposition 5.1 uses the full-weight source presentation, passes to coopposite Borels, and constructs the ordered multiplication (13). Its generator pairing agrees numerically with (2). Identifying it with the convention intended in (1) still requires a dictionary for the original extended Cartans, coproduct, antipode, both skew rules and the order of the double. The comparison of $\mathcal{T}$ with the source presentation does not by itself provide that dictionary.

Finally, the base in (4) is an explicit generic choice, not a recovered hypothesis of the original problem. The matrix inverse above invokes the source condition $v^8 \neq 1$, with $v^2 = q$. No conclusion

for $q^4 = 1$, a divided-power form, or a restricted quantum group follows from that argument. A specialization to an algebraic value of t cannot be obtained merely by evaluating a $\mathbb{Q}(t)$-isomorphism: a nonzero polynomial vanishing at that value is invertible in $\mathbb{Q}(t)$. A result over a suitable localized coefficient ring, together with its geometric identifications, is needed for such a deduction.

Thus the pairing normalization, auxiliary inverse maps, injectivity mechanism and full diagonal quotient are available under the stated hypotheses. Establishing their correspondence with the original skein algebra, arcs, coefficients and quantum-group conventions remains necessary for both (1) and (3).

References

[1] Yuqun Chen, Hongshan Shao and K. P. Shum, *Rosso–Yamane Theorem on PBW basis of* $U_q(A_N)$, arXiv:0804.0954v1 (2008).

[2] Thang T. Q. Lê and Tao Yu, *Quantum traces for* SL_n*-skein algebras*, arXiv:2303.08082v2 (2025).

[3] Christian Voigt and Robert Yuncken, *Complex semisimple quantum groups and representation theory*, arXiv:1705.05661v4 (2020).