

Advanced Multivariable Calculus - Homework 4

Philip Warton

May 16, 2021

Problem 1

Let $G \subset \mathbb{R}^n$ be open and let $f : G \rightarrow \mathbb{R}$ and suppose that ∂_{x_i} exists and is bounded for every $1 \leq i \leq n$. Show that f is continuous.

Proof. Let $\vec{x} \in G$ be a point arbitrarily. The since G is open we know that some ϵ -neighborhood of \vec{x} is contained in G . Let $\epsilon > 0$ be arbitrary. To show that there exists some $\delta > 0$ such that $\vec{y} \in B_\delta(\vec{x}) \Rightarrow f(\vec{y}) \in B_\epsilon(f(\vec{x}))$ would suffice to show that f is continuous by the definition of continuity. Define

$$M = \max_{i \in \{1, 2, \dots, n\}} \{M_i\}$$

Where $M_i = \sup_{\vec{g} \in G} \{\partial_{x_i}\}$. Then define $0 < \delta < \frac{\epsilon}{nM}$ such that $B_\delta(\vec{x}) \subset G$. Write $\vec{x} = (x_1, x_2, \dots, x_n)$ and let $\vec{y} \in B_\delta(\vec{x})$. Let $\vec{u}_k = (y_1, \dots, y_k, x_{k+1}, \dots, x_n)$ for $k = 1, 2, \dots, n$. Then,

$$\begin{aligned} |f(\vec{x}) - f(\vec{y})| &= |f(\vec{x}) - f(\vec{u}_1) + f(\vec{u}_1) - f(\vec{u}_2) + f(\vec{u}_2) - \dots - f(\vec{u}_k) + f(\vec{u}_k) - f(\vec{y})| \\ &\leq |f(\vec{x}) - f(\vec{u}_1)| + |f(\vec{u}_1) - f(\vec{u}_2)| + \dots + |f(\vec{u}_k) - f(\vec{y})| \\ &\leq M|x_1 - y_1| + M|x_2 - y_2| + \dots + M|x_n - y_n| \\ &\leq nM\delta \\ &= nM \frac{\epsilon}{nM} = \epsilon \end{aligned}$$

We know that $|f(\vec{u}_{i+1}) - f(\vec{u}_{i+2})| \leq M|x_i - x_{i+1}|$ by the mean value theorem, applied to f in the i -th plane at our point \vec{x} . □

Problem 2

Let

$$f(x, y) = \begin{cases} \frac{x^3}{x^2 + y^2} & (x, y) \neq \mathbf{0} \\ 0 & (x, y) = \mathbf{0} \end{cases}$$

a)

Show that ∂_x, ∂_y exist and are bounded.

Suppose that $(x, y) \neq \mathbf{0}$. Then we write

$$\frac{\partial f}{\partial x} = \frac{(x^2 + y^2)(3x^2) - (x^3)(2x)}{(x^2 + y^2)^2} = \frac{x^4 + 3x^2y^2}{x^4 + 2x^2y^2 + y^4}$$

$$\frac{\partial f}{\partial y} = \frac{(x^2 + y^2)(0) - (x^3)(2y)}{(x^2 + y^2)^2} = \frac{-x}{x^2 + y^2} \cdot \frac{2xy}{x^2 + y^2}$$

Then if $(x, y) = \mathbf{0}$ simply write $\partial_x(\mathbf{0}) = 1, \partial_y(\mathbf{0}) = 0$ by computing $\lim_{h \rightarrow 0} \frac{f(h, 0) - f(0, 0)}{h}$ and $\lim_{h \rightarrow 0} \frac{f(0, 1) - f(0, 0)}{h}$. To show that these are bounded we write

$$\begin{aligned} \frac{\partial f}{\partial x} &= \frac{x^4}{x^4 + 2y^2x^2 + y^4} + \frac{3x^2y^2}{x^2 + 2x^2y^2 + y^4} \\ &\leq \frac{x^4}{x^4} + 3x^2y^2 2x^2y^2 \\ &\leq 1 + \frac{3}{2} \end{aligned}$$

Because every term is squared and positive, it follows that $0 \leq \partial_x \leq \frac{5}{2}$ and is bounded. For our y partial, we write

$$0 \leq \frac{x^2}{x^2 + y^2} \leq 1$$

Then we say that

$$\begin{aligned} (|y| - |x|)^2 &\geq 0 \\ y^2 - 2|y||x| + x^2 &\geq 0 \\ y^2 + x^2 &\geq 2|x||y| \\ 1 &\geq \frac{2|x||y|}{x^2 + y^2} \\ 1 &\geq \left| \frac{2xy}{x^2 + y^2} \right| \end{aligned}$$

So then we have $\partial_y = \frac{x^2}{x^2 + y^2} \cdot \frac{2xy}{x^2 + y^2}$ is a product of bounded functions, and is therefore bounded.

b)

Show that $\nabla_{\mathbf{u}}(f)$ exists at $\mathbf{0}$ for every unit vector \mathbf{u} .

Denote $\mathbf{u} = (u_1, u_2)$. We compute the directional derivative as

$$\begin{aligned} \lim_{h \rightarrow 0} \frac{f(\mathbf{0} + h\mathbf{u}) - f(\mathbf{0})}{h} &= \lim_{h \rightarrow 0} \frac{\frac{hu_1^3}{u_1^2 + u_2^2}}{h} \\ &= \lim_{h \rightarrow 0} \frac{u_1^3}{u_1^2 + u_2^2} \end{aligned}$$

We know that $u_1^2 + u_2^2 = 1$ since \mathbf{u} is a unit vector, and we know that $|u_1| \leq 1$ for the same reason. So it follows that $|\nabla_{\mathbf{u}}f(\mathbf{0})| < 1$.

c)

Choose $\mathbf{u} = (\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})$. Then we have

$$\nabla_{\mathbf{u}}f(\mathbf{0}) = \frac{\left(\frac{1}{\sqrt{2}}\right)^3}{\frac{1}{\sqrt{2}}^2 + \frac{1}{\sqrt{2}}^2} = \left(\frac{1}{\sqrt{2}}\right)^3 = 2^{-3/2}$$

However, note that

$$\nabla f(\mathbf{0}) \cdot \mathbf{u} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \cdot \begin{pmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{pmatrix} = \frac{1}{\sqrt{2}} \neq \frac{1}{\sqrt{2}^3}$$

Problem 3

Let $G \subset \mathbb{R}^n$ be an open set, and assume $f : G \rightarrow \mathbb{R}$ is differentiable on G . Also assume f has a local maximum at $x \in G$. Prove that $\nabla f(x) = \mathbf{0}$.

Proof. Write $\mathbf{x} = (a_1, a_2, \dots, a_n)$. Now write

$$f_i(t) = f(a_1, \dots, a_{i-1}, t, a_{i+1}, \dots, a_n)$$

For every $i = 1, 2, \dots, n$. If $t \in (-\epsilon + a_i, a_i + \epsilon)$ then $(a_1, \dots, a_{i-1}, t, a_{i+1}, \dots, a_n) \in B_\epsilon(\mathbf{x})$. So it follows that for each such t in the ϵ -neighborhood of a_i , $f_i(t) \leq f(\mathbf{x}) = f_i(a_i)$. Thus we have that $t = a_i$ is a local maximum for f_i , and therefore

$$f'_i(a_i) = \frac{\partial f}{\partial x_i}(\mathbf{x}) = \mathbf{0}$$

Since f is differentiable, we know that each partial derivative exists, and it must be the case that these partial derivatives for x_i are equal to the derivative of the restricted function f_i at the point \mathbf{x} by definition of the partial derivative. \square

Problem 4

Let $f(x, y) = (e^x \cos y, e^x \sin y)$.

a)

Compute the Jacobian J_f .

We are going to compute the appropriate derivatives as

$$\frac{\partial}{\partial x} e^x \cos y = e^x \cos y$$

$$\frac{\partial}{\partial y} e^x \cos y = -e^x \sin y$$

$$\frac{\partial}{\partial x} e^x \sin y = e^x \sin y$$

$$\frac{\partial}{\partial y} e^x \sin y = e^x \cos y$$

Then our matrix will clearly be

$$J_f = \begin{pmatrix} e^x \cos y & -e^x \sin y \\ e^x \sin y & e^x \cos y \end{pmatrix}$$

b)

Show that $J_f(x, y)^{-1}$ exists for every point in \mathbb{R}^2 .

Proof. We know that $J_f(x, y)^{-1}$ exists if and only if $\det(J_f(x, y)) \neq 0$. So we compute

$$\begin{aligned} \det(J_f(x, y)) &= (e^x \cos y)(e^x \cos y) - ((-e^x \sin y)(e^x \sin y)) \\ &= (e^x)^2(\cos^2 y) + (e^x)^2(\sin^2 y) \\ &= (e^x)^2(\cos^2 y + \sin^2 y) \\ &= (e^x)^2 \end{aligned}$$

Since e^x is strictly positive, so is its square, so obviously $\det(J_f(x, y)) = (e^x)^2 > 0$ and we say that the matrix is invertible. □

c)

Show that f is not invertible globally; that is, find different points of the domain that are mapped to the same value in the range.

Proof. Since e^x is a bijective map, we fix $x = 0$. Then we know that trigonometric functions are 2π -periodic, so pick the following two points $\mathbf{a} = (0, 0)$, $\mathbf{b} = (0, 2\pi)$. Then we know that $f(\mathbf{a}) = (1, 0)$ and $f(\mathbf{b}) = (1, 0)$. So we know that the pre-image of $(1, 0)$ includes both \mathbf{a} and \mathbf{b} so then f^{-1} is not globally invertible. □

Problem 5

Let $f_1(x, y_1, y_2) = 3y_1 + y_2^2 + 4x = 0$, $f_2(x, y_1, y_2) = 4y_1^3 + y_2 + x = 0$.

a)

Verify that $(x, y_1, y_2) = (-4, 0, 4)$ is a solution.

We have that

$$f_1(-4, 0, 4) = 3(0) + (4)^2 + 4(-4) = 16 - 16 = 0$$

And also that

$$f_2(-4, 0, 4) = 4(0)^3 + (4) + (-4) = 0$$

So then we know that $(-4, 0, 4)$ is a valid solution.

b)

Verify that $\exists g = (g_1, g_2)$ such that $g_1(x) = y_1, g_2(x) = y_2$ for all solutions near $(-4, 0, 4)$.

We write

$$\begin{pmatrix} \frac{\partial f_i}{\partial y_j} \end{pmatrix} = \begin{pmatrix} 3 & 2y_2 \\ 12y_1^2 & 1 \end{pmatrix}$$

Then because the determinant is equal to $3 - (12y_1^2)(2y_2)$, if we plug in $s = (-4, 0, 4)$ we will get $\det \begin{pmatrix} \frac{\partial f_i}{\partial y_j} \end{pmatrix} = 3 \neq 0$. So by the implicit function theorem it follows that $\exists g : \mathbb{R} \rightarrow \mathbb{R}^2$ such that for every $(x, y_1, y_2) \in S \cap B_\epsilon(-4, 0, 4)$ where S is the set of solutions to our system of equations $(x, y_1, y_2) = (x, g(x))^t$ therefore $g_1(x) = y_1$ and $g_2(x) = y_2$.

c)

So we use the final piece of the implicit function theorem for solutions around $(-4, 0, 4)$ since

$$\begin{pmatrix} 4 \\ 1 \end{pmatrix} + \begin{pmatrix} 3 & 8 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \frac{\partial g_1}{\partial x} \\ \frac{\partial g_2}{\partial x} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

So it follows that $\frac{\partial g_2}{\partial x} = -1$ and $\frac{\partial g_1}{\partial x} = \frac{4}{3}$.

Problem 6

Let $T = \{(x, y, u, v) \in \mathbb{R}^4 \mid x^2 + y^2 = 1, ux + vy = 0\}$. Then let

$$f(x_1, x_2, y_1, y_2) = (x_1^2 + y_1^2, x_2x_1 + y_2y_1)$$

Where $x_1 = x, y_1 = y, x_2 = u, y_2 = v$. Then we write $T = f^{-1}(1, 0)$.

a)

Compute $\frac{\partial f_i}{\partial y_j}$. We simply take each of our partial derivatives,

$$\begin{aligned} \begin{pmatrix} \frac{\partial f_i}{\partial y_j} \end{pmatrix} &= \begin{pmatrix} \frac{\partial f_1}{\partial y_1} & \frac{\partial f_1}{\partial y_2} \\ \frac{\partial f_2}{\partial y_1} & \frac{\partial f_2}{\partial y_2} \end{pmatrix} \\ &= \begin{pmatrix} 2y_1 & 0 \\ y_2 & y_1 \end{pmatrix} \end{aligned}$$

b)

We know that $\det \begin{pmatrix} \frac{\partial f_i}{\partial y_j} \end{pmatrix} = 2y_1^2$, so it follows that if $y_1 = y \neq 0$ then this determinant is non zero. So there exists by the implicit function theorem a continuously differentiable function $g : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that for all $(x_1, x_2, y_1, y_2) \in T$ we can write

$$(x_1, x_2, y_1, y_2) = (x_1, x_2, g(x_1, x_2))$$

which is entirely in terms of x and u since $x = x_1, u = x_2$.

c)

Find formulas for y and v in terms of x and u valid when $y \neq 0$.

If $y \neq 0$ then we can write $y = \pm\sqrt{1-x^2}$ and $v = \frac{ux}{y} = \frac{ux}{\pm\sqrt{1-x^2}}$. Then we know that our denominator is non-zero and we have a formula for y and v all in terms of u and x .

d)

If $x \neq 0$ then $x_1 \neq 0$ so we write $x = \pm\sqrt{1-x^2}$ and $u = \frac{vy}{x} = \frac{vy}{\sqrt{1-x^2}}$. Since the denominator is guaranteed to be non-zero we have written x, u in terms of y, v .